

11. Simple germs and moduli

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **25 (1979)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **11.07.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

- (iii) *The germs in Table 2c right simplify only to the germs in Table 2b and 2a.*
- (iv) *A germ in \mathcal{F} not right equivalent to a germ in Table 2a, b, or c right simplifies to a germ in Table 2c.*

11. SIMPLE GERMS AND MODULI

A germ $f \in \mathfrak{m}$ is said to be *right* (or *contact*) *simple* if there is a neighborhood of f in \mathfrak{m} intersecting only finitely many right (or contact) equivalence orbits. In the language of algebraic geometry, a germ f is contact simple if and only if the versal deformation of $f^{-1}(0)$ contains only finitely many isomorphism classes of analytic spaces.

The germs in Table 2a are right and contact simple by Proposition 10.1. The germs in Table 2b are not contact simple (and hence not right simple): \tilde{E}_6 is a family of cones over non-singular elliptic curves in $\mathbf{C}P^2$, \tilde{E}_7 is a family of four lines through the origin in \mathbf{C}^2 , and \tilde{E}_8 is a family of three parabolas [Arnold 1; Siersma]. Note that the germs of Table 2c form one-dimensional families under right equivalence, but all members of the family are contact equivalent [Laufer 4; Siersma p. 54]. Clearly if a germ g right simplifies to f and f is not right simple, then g is not right simple; the same applies to contact equivalence.

Characterization B3. The germ f is right simple.

Characterization B4. The germ f is contact simple.

The equivalence of Characterizations B1 and B3 follows from Proposition 10.1 and the above remarks [Arnold 1]. Characterization B3 implies Characterization B4 by definition. Conversely, a contact simple germ f which is not right simple right simplifies to a germ in Table 2b (by Proposition 10.1), but these are not contact simple. Hence f must be right simple.

The classification of simple germs has recently been extended to complete intersections [Giusti]. The *modality* of a germ f is defined in [Arnold 3]. A right-simple germ is zero-modal; all right equivalence classes of 1 and 2-modal germs have been listed [Arnold 2, 3, 5]. Moduli of resolutions of two-dimensional normal singularities are studied in [Laufer 3, 4]. The following result provides a connection between Characterizations A2 and B3.

THEOREM 11.1 [Randell]. *For almost all germs $f(x, y, z)$ (in the sense of the Newton diagram), the geometric genus p of $f^{-1}(0)$ is less than or equal to the modality of f .*