

# ON THE GENUS OF GENERALIZED FLAG MANIFOLDS

Autor(en): **Glover, Henry H. / Mislin, Guido**

Objektyp: **Article**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **27 (1981)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **14.09.2024**

Persistenter Link: <https://doi.org/10.5169/seals-51749>

## **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

# ON THE GENUS OF GENERALIZED FLAG MANIFOLDS

by Henry H. GLOVER and Guido MISLIN

## INTRODUCTION

Let  $X$  be a nilpotent space of finite type. We denote by  $G(X)$  the genus of  $X$ , i.e. the set of all homotopy types  $Y$  (nilpotent, of finite type) with  $p$ -localizations  $Y_p \simeq X_p$  for all primes  $p$ , (cf. [HMR]). The set  $G(X)$  has been studied extensively in case of  $X$  an  $H$ -space. In particular it is known that for the special unitary group  $SU(n)$  one has

$$|G(SU(n))| \geq \prod_{1 < m < n} (\phi(m!)/2)$$

where  $\phi$  is the Euler function [Z, p. 152]. We are interested in this note in finding non-trivial examples  $X$  with  $G(X) = \{[X]\}$  and we call such spaces *generically rigid*. A large family of such generically rigid spaces is provided by certain generalized flag manifolds. Let

$$G = U(n_1 + n_2 + \dots + n_k)$$

and

$$H = U(n_1) \times U(n_2) \times \dots \times U(n_k),$$

embedded in  $G$  in the obvious way. Then

$$M = M(n_1, n_2, \dots, n_k) = G/H$$

is a generalized flag manifold (generalizing the standard complex flag manifold  $U(n)/T^n$  which corresponds to  $M(1, 1, \dots, 1)$ ). We will show essentially that whenever the homotopy rigidity result for linear actions holds for  $M$  (cf. [L1], [L2], [EL]), then  $M$  is also generically rigid. These two seemingly unrelated rigidity results are tied up by certain results on  $E(X)$  and  $E(X_0)$ , the groups of homotopy classes of self equivalences of  $X$  and  $X_0$ ,  $X_0$  the rationalization of  $X$ .

To make our result more precise, we need some further notation. For

$$M = M(n_1, \dots, n_k) = G/H$$

as above, we write  $N(H)$  for the normalizer of  $H$  in  $G$ . The finite group  $N(H)/H$  acts on  $M$  in an obvious way and it is well known that through that action,  $N(H)/H$  is faithfully represented in  $H^*(M; \mathbf{Q})$ . We can therefore consider  $N(H)/H$  as a subgroup of  $E(M)$  or  $E(M_0)$ . By Theorem 1.1 of [GH2] the canonical map

$$E(M_0) \rightarrow \text{Aut}_{\text{alg}} H^*(M; \mathbf{Q})$$

is a group isomorphism. In particular, the grading automorphisms

$$g(q): H^*(M; \mathbf{Q}) \rightarrow H^*(M; \mathbf{Q})$$

defined by  $g(q)x = q^i x$  for  $x \in H^{2i}(M; \mathbf{Q})$  and  $q \in \mathbf{Q}^*$ , lift to unique self equivalences of  $M_0$  (which we denote also by  $g(q)$ ), and thus

$$\text{Gr}(M_0) = \{g(q) \mid q \in \mathbf{Q}^*\} \subset E(M_0)$$

is a central subgroup isomorphic to  $\mathbf{Q}^*$ .

In all cases of generalized flag manifolds for which  $E(M_0)$  has been computed, the subgroup generated by  $\text{Gr}(M_0)$  and  $N(H)/H$ ,

$$\langle \text{Gr}(M_0), N(H)/H \rangle \subset E(M_0)$$

is all of  $E(M_0)$ . The following conjecture is thus plausible.

*Conjecture C.* Let  $M = M(n_1, n_2, \dots, n_k)$  be a generalized flag manifold. Then

$$E(M_0) = \langle \text{Gr}(M_0), N(H)/H \rangle.$$

A similar conjecture appears in [L1, Conjecture C] but the relationship between the two conjectures is not entirely clear.

The Conjecture C has been verified in the following cases:

- 1)  $n_1 = n_2 = \dots = n_k = 1$  (compare the proof of Thm. 1 in [EL])
- 2)  $n_1 = n_2 = \dots = n_{k-1} = 1, n_k \geq k - 1$  (compare the proof of Theorem 9 in [L1])
- 3)  $n_1 = 2$  and  $k = 2$  (follows from [O])
- 4)  $n_2 > n_1$  and  $k = 2$  ([GH1], [Br])
- 5)  $n_1 = 1, n_2 > 1, n_3 \geq 2n_2^2 - 1$  and  $k = 3$  ([GH2])

The Conjecture C holds therefore for instance for all complex Grassmann manifolds  $G_p(\mathbf{C}^{p+q}) = M(p, q)$  with  $p \neq q$  (since  $M(p, q) \simeq M(q, p)$ ), and for the classical flag manifolds  $U(n)/T^n$ .

Our main theorem may be stated as follows.

THEOREM. Let  $M = M(n_1, \dots, n_k)$  be a generalized flag manifold for which the Conjecture C holds. Then

$$G(M) = \{[M]\}.$$

In particular the Grassmann manifolds  $G_p(\mathbb{C}^{p+q})$  for  $p \neq q$  and the flag manifolds  $U(n)/T^n$  are all generically rigid.

§1. GENUS AND SELF MAPS

Let  $P$  denote a fixed set of primes. Two  $P$ -sequences

$$S_1, S_2: P \rightarrow E(X_0)$$

are called *equivalent*, if there exist maps  $h(0) \in E(X_0)$  and

$$h(p) \in \text{im}(E(X_p) \xrightarrow{\text{can}} E(X_0))$$

such that for all  $p \in P$  one has

$$h(0) S_1(p) = S_2(p) h(p).$$

*Definition 1.1.* We denote by  $P\text{-Seq}(E(X_0))$  the set of equivalence classes of  $P$ -sequences in  $E(X_0)$ .

If  $P$  is a finite set of primes and  $X$  a nilpotent space of finite type, then there is a canonical map

$$\theta: G(X) \rightarrow P\text{-Seq}(E(X_0)).$$

It is defined as follows. Let  $Y \in G(X)$  and  $P = \{p_1, \dots, p_n\}$ . Then the localization  $Y_P$  is a pull-back of maps  $X_{p_i} \xrightarrow{\lambda_i} X_0$ , i.e.  $Y_P \simeq \text{hoinvlim} \{X_{p_i} \xrightarrow{\lambda_i} X_0\}$ . The maps  $\lambda_i$  induce equivalences  $\bar{\lambda}_i \in E(X_0)$  and we put

$$\theta(Y) = \{[\bar{\lambda}_1, \bar{\lambda}_2, \dots, \bar{\lambda}_n]\}.$$

If  $Y_P$  may also be represented by  $\text{hoinvlim} \{X_{p_i} \xrightarrow{\mu_i} X_0\}$ , then there exist maps  $h(0) \in E(X_0)$  and  $\tilde{h}(p_i) \in E(X_{p_i})$ ,  $i \in \{1, \dots, n\}$  rendering the diagrams

$$\begin{array}{ccc}
 X_{p_i} & \xrightarrow{\tilde{h}(p_i)} & X_{p_i} \\
 \lambda_i \downarrow & & \downarrow \mu_i \\
 X_0 & \xrightarrow{h(0)} & X_0
 \end{array}$$

homotopy commutative and thus inducing  $\text{hoinvlim } \{\lambda_i\} \simeq \text{hoinvlim } \{\mu_i\}$ . Hence

$$\{[\bar{\lambda}_1, \dots, \bar{\lambda}_n]\} = \{[\bar{\mu}_1, \dots, \bar{\mu}_n]\} \in P\text{-Seq}(E(X_0))$$

and therefore  $\theta$  is well defined.

LEMMA 1.2. Let  $X$  be a nilpotent space of finite type and let  $P$  denote a finite set of primes. Then

$$\theta: G(X) \rightarrow P\text{-Seq}(E(X_0))$$

is surjective with fibers of the form

$$\theta^{-1}(\theta(Y)) = \{Z \in G(X) \mid Z_P \simeq Y_P\}.$$

*Proof.* Let  $P = \{p_1, \dots, p_n\}$  and

$$\{[\bar{f}_1, \dots, \bar{f}_n]\} \in P\text{-Seq}(E(X_0)).$$

Let  $e_i: X_{p_i} \rightarrow X_0$  denote the canonical maps and put

$$f_i = \bar{f}_i \circ e_i: X_{p_i} \rightarrow X_0.$$

Define  $W = \text{hoinvlim } \{f_i\}$ ;  $W$  comes equipped with a canonical map  $f: W \rightarrow X_0$ . Let  $Z$  be the homotopy pull back of  $W \xrightarrow{f} X_0 \xleftarrow{\text{can}} X_{\bar{P}}$ , where  $\bar{P}$  denotes the set of primes complementary to  $P$ . Then  $Z \in G(X)$  and

$$\theta(Z) = \{[\bar{f}_1, \dots, \bar{f}_n]\};$$

thus  $\theta$  is surjective. It is clear from the definition of  $\theta$  that for  $Y, Z \in G(X)$  one has  $\theta(Y) = \theta(Z)$  if and only if  $Y_P \simeq Z_P$ .


The next lemma provides a sufficient condition for  $\theta$  to be monic "at the basepoint".

LEMMA 1.3. Let  $X$  be a nilpotent space of finite type. Suppose that there exists a finite set of primes  $P$  with complement  $\bar{P}$  such that


- a)  $Y \in G(X)$  implies  $Y_{\bar{P}} \simeq X_{\bar{P}}$
- b) every  $f \in E(X_0)$  can be written as  $f_1 \circ f_2$  with  $f_1 \in \text{im}(E(X_P) \xrightarrow{\text{can}} E(X_0))$  and  $f_2 \in \text{im}(E(X_{\bar{P}}) \rightarrow E(X_0))$ .

Then for  $\theta: G(X) \rightarrow P\text{-Seq}(E(X_0))$  as above, one has  $\theta^{-1}(\theta(X)) = \{X\}$ .

*Proof.* Let  $Y \in G(X)$  with  $\theta(Y) = \theta(X)$ . Then  $Y_P \simeq X_P$  by the definition of  $\theta$ , and  $Y_{\bar{P}} \simeq X_{\bar{P}}$  by assumption. Hence  $Y$  may be obtained as a homotopy pull back of the form




If  $\alpha$  induces  $\bar{\alpha} \in E(X_0)$  and if  $\gamma = \bar{\alpha}^{-1} \circ \beta$ , then  $Y$  is also a pull back of the form



Let  $\bar{\gamma} \in E(X_0)$  be the map induced by  $\gamma$  and write  $\bar{\gamma} = f_1 f_2$  with

$$f_1 \in \text{im}(E(X_P) \rightarrow E(X_0)), \quad f_2 \in \text{im}(E(X_{\bar{P}}) \rightarrow E(X_0)).$$

Choose a lift  $\tilde{f}_1^{-1} \in E(X_P)$  of  $f_1^{-1}$  and a lift  $\tilde{f}_2 \in E(X_{\bar{P}})$  of  $f_2$ . Then  $f_1^{-1} \bar{\gamma} = \text{can} \circ \tilde{f}_2$  and one can form a commutative diagram,



which shows that  $Y \simeq X$ .

§2. THE CASE OF GENERALIZED FLAG MANIFOLDS

The following result is an easy consequence of [F].

LEMMA 2.1. Let  $M$  be a generalized flag manifold. Then the following holds.

- a) If  $g(\lambda) \in Gr(M_0)$  is a grading map with  $\lambda \in \mathbf{Z}_Q^*$  for some (not necessarily finite) set of primes  $Q$ , then  $g(\lambda)$  lifts to a homotopy equivalence  $\tilde{g}(\lambda): M_Q \rightarrow M_Q$ .
- b) Let  $P$  be an arbitrary set of primes with complement  $\bar{P}$ . Then every

$$f \in \langle Gr(M_0), N(H)/H \rangle$$

may be written in the form  $f = f_1 \circ f_2$  with

$$f_1 \in \text{im}(E(M_P) \rightarrow E(M_0))$$

and

$$f_2 \in \text{im}(E(M_{\bar{P}}) \rightarrow E(M_0)).$$

*Proof.* Let  $\lambda = k/l$  with  $k$  and  $l$  relatively prime integers. Then  $g(k)$  and  $g(l)$  lift to equivalences

$$\tilde{g}(k), \tilde{g}(l): M_Q \rightarrow M_Q.$$

since necessarily  $k, l \in \mathbf{Z}_Q^*$  (compare [F]). Thus  $\tilde{g}(k) \tilde{g}(l)^{-1}$  is a lift of  $g(\lambda)$ . For b) we note that  $f = g(\rho) \circ \sigma$  for some  $\rho \in \mathbf{Q}^*$  and

$$\sigma \in N(H)/H.$$

If we write  $\rho = \rho_1 \cdot \rho_2$  with  $\rho_1 \in \mathbf{Z}_P^*$  and  $\rho_2 \in \mathbf{Z}_P^*$ , then

$$f = g(\rho_1) \cdot (g(\rho_2) \sigma)$$

and we may choose

$$f_1 = g(\rho_1), f_2 = g(\rho_2) \sigma.$$

Since  $\sigma$  lifts even to  $E(M)$ , we infer by using a) that  $f_1$  and  $f_2$  lift as desired.

A final step towards proving the Theorem formulated in the introduction consists in the following.

LEMMA 2.2. Let  $M$  be a generalized flag manifold for which Conjecture C holds. Then for every finite set of primes  $P$ ,

$$P\text{-Seq}(E(M_0)) = \{[1, 1, \dots, 1]\}.$$

*Proof.* Let  $\{[\mu_1, \dots, \mu_n]\} \in P\text{-Seq}(E(M_0))$ , where  $P = \{p_1, \dots, p_n\}$  and

$$\mu_i \in \text{im}(E(M_{p_i}) \rightarrow E(M_0))$$

for all  $i$ . Then  $\mu_i = g(\lambda_i) \circ \sigma_i$  with  $\lambda_i \in \mathbf{Q}^*$  and

$$\sigma_i \in N(H)/H \subset E(M_0).$$

Define  $\lambda \in \mathbf{Q}^*$  by  $\lambda = \prod p_i^{m_i}$ , where  $m_i \in \mathbf{Z}$  is such that  $p_i^{m_i} \lambda_i \in \mathbf{Z}_{p_i}^*$ . Then  $g(\lambda) \mu_i = g(\lambda \lambda_i) \sigma_i$  with  $\lambda \lambda_i \in \mathbf{Z}_{p_i}^*$ . By Lemma 2.1 a) we know that  $g(\lambda \lambda_i)$  lifts to  $M_{p_i}$ , and since  $\sigma_i$  lifts even to  $M$  we conclude that

$$h(p_i) = g(\lambda \lambda_i) \sigma_i \in \text{im}(E(M_{p_i}) \rightarrow E(M_0))$$

for all  $i$ . The equation

$$g(\lambda) \mu_i = h(p_i), i \in \{1, \dots, n\}$$

show that  $\{[\mu_1, \dots, \mu_n]\} = \{[1, \dots, 1]\} \in P\text{-Seq}(E(M_0))$ .

The proof of the main Theorem:

Let  $M$  be a generalized flag manifold for which the Conjecture C holds. Since  $M$  is a formal space we can find for every  $N \in G(M)$  a rational equivalence



$f(N): N \rightarrow M$ . Let  $P(M)$  denote the set of primes which appear in any of the orders of

$$\ker (f(N)_* : H_*(N; \mathbf{Z}) \rightarrow H_*(M; \mathbf{Z}))$$

or  $\text{coker } f(N)_*$ ,  $N$  ranging over  $G(M)$ . The set  $P(M)$  is finite, since each  $\ker f(N)_*$  and  $\text{coker } f(N)_*$  is finite and since  $G(M)$  is a finite set by [W]. Consider now the map

$$\theta: G(M) \rightarrow P\text{-Seq } E(M_0)$$

with respect to this finite set of primes  $P(M) = P$ . Since  $P$  is finite,

$$P\text{-Seq } (E(M_0))$$

consists of only one element (Lemma 2.2). It remains to show that

$$\theta^{-1}(\theta(M)) = \{M\}.$$

For this we apply Lemma 1.3. Note that  $N \in G(M)$  implies  $N_{\bar{P}} \simeq M_{\bar{P}}$  since  $f(N): N \rightarrow M$  is a  $\bar{P}$ -equivalence. Moreover, the condition b) of 1.3 is satisfied in view of Lemma 2.1 b). Therefore we conclude that  $G(M) = \{[M]\}$  and the proof is completed.

*Note added in proof.* Since this paper went to press, we have been informed that Conjecture C has been proved for the case  $k = 2, n_1 = n_2$ , by M. Hoffman: "Cohomology endomorphisms of complex flag manifolds", Ph.D. dissertation, MIT 1981. As a consequence, it follows that all complex Grassmann manifolds are generically rigid.

## REFERENCES

- [B] BREWSTER, S. *Automorphisms of the cohomology ring of finite Grassmann manifolds*. Ph.D. dissertation, Ohio State University, 1978.
- [EL] EWING, J. and A. LIULEVICIUS. Homotopy rigidity of linear actions on friendly homogeneous spaces. *J. of Pure and Appl. Algebra* 18 (1980), 259-267.
- [F] FRIEDLANDER, E. Maps between localized homogeneous spaces. *Top.* 16 (1977), 205-216.
- [GH1] GLOVER, H. and W. HOMER. Endomorphisms of the cohomology ring of finite Grassmann manifolds. *Lecture Notes in Math. Springer-Verlag*, 657 (1978), 170-193.
- [GH2] ——— Self maps of flag manifolds I. *Trans. of the Amer. Math. Soc.* 267 (1981), 423-434.
- [HMR] HILTON, P. G. MISLIN and J. ROITBERG. *Localization of nilpotent groups and spaces*. North Holland, 1975.
- [L1] LIULEVICIUS, A. Homotopy rigidity of linear actions: characters tell all. *Bull. Amer. Math. Soc.* 84 (1978), 213-221.
- [L2] ——— Equivariant K-theory and homotopy rigidity. *Springer-Verlag Lecture Notes in Math.*, 788 (1980), 340-358.
- [O] O'NEILL, L. *The fixed point property for Grassmann manifolds*. Ph.D. dissertation, Ohio State University, 1974.
- [W] WILKERSON, C. Applications of minimal simplicial groups *Top.* 15 (1976), 111-130.
- [Z] ZABRODSKY, A. *Hopf Spaces*. North Holland, 1976.

(Reçu le 17 décembre 1980)

Henry H. Glover

Ohio State University  
Columbus, Ohio 43210

Guido Mislin

ETH  
CH-8092 Zürich

**Vide-leer-empty**