

Appendix B

Objektyp: **Appendix**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **29 (1983)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **08.08.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

APPENDIX B

Let K be a separable quadratic extension of a field k . We denote $x \mapsto \bar{x}$ the non trivial element $\text{Gal}(K/k)$. Let D be a simple algebra with dimension n^2 over its center K . We will check the criterion of the text, for the existence of an involution of the second kind on D , i.e. of an anti-involution $*$ of D , inducing $x \mapsto \bar{x}$ on K . The criterion is that $N_{K/k} \text{cl}(D) = 0$ in $\text{Br}(k)$.

Let us localize, for the étale topology, over $\text{Spec}(k)$. This means making large enough étale extensions of scalars $\otimes_k k'$, and keeping track of the functoriality in k' . The field K becomes the separable quadratic extension $K' = K \otimes_k k'$ of k' . The algebra D becomes $D' = D \otimes_k k'$, and is of the form $D' = \text{End}_{K'}(V')$, for V' a free module K' . The module V' is not determined uniquely by D' , only up to homotheties (the corresponding projective space is uniquely determined).

For any K -module M , let M^- be the module deduced from M by the extension of scalars $\bar{} : K \rightarrow K$, i.e. the module, unique up to unique isomorphism, provided with an anti-linear isomorphism $x \mapsto \bar{x} : M \xrightarrow{\sim} M^-$. Similarly for K' -modules. If $D' = \text{End}(V')$, then $D'^- = \text{End}(V'^-)$, and

$$(D \otimes_k D^-)' = \text{End}(V' \otimes V'^-).$$

Let W' be the fixed subspace of the anti-linear automorphism of $V' \otimes V'^-$ defined by $v \otimes \bar{w} \mapsto w \otimes \bar{v}$. It is the space of Hermitian forms on the dual of V' . One has $W' \otimes_{k'} K' = V' \otimes V'^-$. If $D_1 \subset D \otimes_k D^-$ is the fixed subspace of the anti-linear automorphism of $D \otimes_k D^-$ defined by $x \otimes \bar{y} \mapsto y \otimes \bar{x}$, then D'_1 is the k' -form of the K' -algebra $(D \otimes_k D^-)' = \text{End}(V' \otimes V'^-)$ deduced from the k' -form W' of the K' -module $V' \otimes V'^- : D'_1 = \text{End}_{k'}(W')$.

Involutions of the second kind on D' correspond one to one to non degenerate Hermitian forms on V' , taken up to a factor (in k'^*). Those, in turn, by the “dual form” construction, correspond to “non degenerate” elements of W' . Again, one has to take them up to a factor. The projective space $\mathbf{P}(W')$ over k' is determined up to unique isomorphism by D' . It is hence (this is the point of localisation) defined over $k : \mathbf{P}(W') = P \otimes_k k'$, functorially in k' . The k -points of P (rather, the non degenerate points) parametrize the involutions of the second kind on D .

The functorial isomorphism $D'_1 = \text{End}_{k'}(W')$ shows that P is the form of projective space (Severi-Brauer variety) attached to D_1 . It has a rational point, and is then the ordinary projective space, if and only if D_1 is a matrix algebra.

This shows that D has involutions of the second kind if and only if the class of D_1 in $\text{Br}(k)$ is trivial. This class is the required norm $N_{K/k}(\text{cl}(D))$. In the localization spirit, this can be deduced from the fact that the homothety by $\lambda \in K'^*$ of V' induces on W' the homothety by $N_{K'/k'}(\lambda) \in k'^*$.

APPENDIX C

For $n \geq 3$, examples can be obtained as follows: take $k' = \mathbf{Q}[\zeta]$, with $\zeta = \exp(2\pi i/n)$, and $k = k' \cap \mathbf{R}$. Fix $a, b \in k'^*$ and let D be the k' -algebra generated by X, Y , subject to

$$\begin{aligned} X^n &= a, & Y^n &= b \\ XY &= \zeta YX. \end{aligned}$$

It admits the anti-involution $*$, inducing complex conjugation on k' , defined by $\zeta^* = \zeta^{-1}$, $X^* = X$, $Y^* = Y$. The algebra D is of the type we require, provided it is a division algebra. This happens already with $a, b \in \mathbf{Z}$: take for a a prime congruent to 1 mod n , and for b an integer whose residue mod a has in the cyclic group of order n $(\mathbf{Z}/(a))^*/(\mathbf{Z}/(a))^{*n}$ an image of exact order n . For instance $n = 3$, $a = 7$, $b = 2$. For $n = 2$, one proceeds similarly with $k' = \mathbf{Q}[i]$, $\zeta = -1$, a congruent to 1 mod 4 and b not a square mod a . For instance, $a = 5$, and $b = 2$. In each case, the assumption on a ensures that k' embed in the a -adic completion \mathbf{Q}_a of \mathbf{Q} , and the fact that D is a division algebra can be seen locally at a : $D \otimes_{k'} \mathbf{Q}_a$ is a division algebra with center \mathbf{Q}_a .