

§1. The Bergman-Roseblade Theorem

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **30 (1984)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **06.08.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

§ 1. THE BERGMAN-ROSEBLADE THEOREM

We introduce some notation. Let A denote a finitely generated free abelian group. Let k be a field and have \underline{A} designate an isomorphic copy of A inside the multiplicative group of some field extension of k . We will write $k[\underline{A}]$ for the k -algebra generated by \underline{A} and $k(\underline{A})$ for its field of fractions. The reader is cautioned that $k[\underline{A}]$ is not the group algebra; distinct elements of \underline{A} need not be linearly independent over k . (It's even possible for \underline{A} to be contained in k^* .)

We will reserve the notation $k[A]$, without the underbar, for the group algebra. There is an obvious relation between the free object $k[A]$ and $k[\underline{A}]$. Indeed, the given isomorphism $A \simeq \underline{A}$ induces a k -algebra isomorphism $k[A]/P \simeq k[\underline{A}]$ where P is a prime ideal. The ideal P is "faithful with respect to A ."

Suppose that G is a group which acts on A . Set

$$D = \{a \in A \mid a \text{ has a finite } G\text{-orbit}\}.$$

It is sometimes called the orbital subgroup or relative finite conjugate subgroup.

We are primarily interested in a group G which acts as a group of k -automorphisms of $k[\underline{A}]$. (The slight awkwardness of language allows us to include possibly nonfaithful actions.) We say that G acts multiplicatively on $k[\underline{A}]$ if G stabilizes \underline{A} . Thus if $k[\underline{A}] = k[A]/P$ as described above, then P is a G -stable ideal under the corresponding action on $k[A]$.

The fundamental theorem in multiplicative invariant theory is Roseblade's Theorem D ([10]). Roseblade based his arguments on profound insights of G. Bergman ([3]).

BERGMAN-ROSEBLADE THEOREM. *Assume that G acts multiplicatively on $k[\underline{A}] = k[A]/P$. Then $P = (P \cap k[D])k[A]$.*

To understand the implications of this theorem, we take a closer look at D . Obviously D is a finitely generated abelian group. Since each generator has a finite G -orbit, D is centralized by a subgroup of finite index in G . In other words, G acts like a finite group of automorphisms when restricted to D .

It is easy to see that if a power of an element in A has a finite orbit, then so does the original element. Hence there is a splitting $A = D \times B$. (Unfortunately, there may be no choice of B which is stabilized by G .) The

conclusion of the Bergman-Roseblade Theorem can be rewritten—every element in P has a unique representation $\sum f(b)b$ where $b \in B$ and $f(b) \in P \cap k[D]$. Thus $k[A]$ is the group ring $(k[D])[B]$ for a finitely generated free abelian group B .

Roseblade proves that the fixed ring $(k[A])^G$ lies in $k[D]$ ([10], Lemma 10). This will also be a consequence of the first lemma in the next section. In any event, it has a remarkable consequence.

THEOREM 1. *Assume that G is an arbitrary group acting multiplicatively on $k[A]$. Then $k[A]^G$ is finitely generated.*

Proof. As we have remarked, $(k[A])^G = (k[D])^G$. But G acts like a finite group of automorphisms on the affine algebra $k[D]$. Noether's Theorem ([11]), states that, in this case, the algebra of invariants is a finitely generated algebra. \square

This is an unexpected surprise. In contrast to the situation for linear actions, Hilbert's 14th problem holds for multiplicative actions without any restriction on the group!

The theme of the paper has emerged. A theory of invariants for multiplicative actions is ultimately a theory for finite groups.

§ 2. GALOIS THEORY

We begin this section by establishing an analogue to the "finiteness" phenomenon of the previous section, for a multiplicative action of G on $k(A)$. Notation is taken from § 1.

LEMMA 2. *Suppose that G acts multiplicatively on $k(A)$. Then $k(A)^G \subset k(D)$.*

Proof. The crucial fact is that $k(D)[B]$ is a unique factorization domain. If ${}^g f = f$ for $f \in k(A)$ then we can write $f = \alpha/\beta$ where α and β in $k(D)[B]$ have no common factors. The invariance of f becomes

$$({}^g \alpha)\beta = ({}^g \beta)\alpha \quad \text{for all } g \in G.$$

Hence $\alpha \mid {}^g \alpha$ and ${}^g \alpha \mid \alpha$; we have $({}^g \alpha)\alpha^{-1}$ a unit in $k(D)[B]$. A similar result holds for β .

$${}^g \alpha = u(g)\alpha \quad \text{and} \quad {}^g \beta = w(g)\beta$$

for $u(g), w(g) \in k(D)^* \cdot B$.