

1. Introduction

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **31 (1985)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **13.07.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

A GEOMETRIC PROOF OF BIEBERBACH'S THEOREMS ON CRYSTALLOGRAPHIC GROUPS

by Peter BUSER

Pour Ariane et Georges

1. INTRODUCTION

In 1910 Bieberbach proved two celebrated theorems in response to Hilbert's 18th problem.

THEOREM I. *Every discrete group of isometries acting on the n -dimensional euclidean space \mathbf{R}^n with compact fundamental domain contains n linearly independent translations.*

Groups which satisfy the hypothesis of Theorem I are called n -dimensional crystallographic groups.

THEOREM II. *For each fixed n there are only finitely many isomorphism classes of n -dimensional crystallographic groups.*

Bieberbach's original proof of Theorem I is based on Minkowski's Theorem on simultaneous rational approximation and is difficult to read. Shortly after it came out, Frobenius gave a more accessible proof which is based on an argument using the commutativity of unitary matrices. Frobenius's method has, in one form or another, become standard in the contemporary literature.

In this note we present a completely different approach to Theorem I which has its origins in Gromov's work on almost flat manifolds [5]. The new idea is to start with those rigid motions which have a very small rotation part (cf. § 2 for notation), and then proceed to show that, in fact, these motions are pure translations. The simplification which results from this approach is striking.

We also give a new proof of Theorem II which does not run via the usual algebraic characterization of a crystallographic group. Instead we shall

use a method which is more in the spirit of Minkowski's geometry of numbers, from where Bieberbach's original arguments departed.

Since the material is standard, the exposition will be condensed. Yet some efforts have been made not to frustrate the reader by omitting details.

I would like to express my thanks to Leon Charlap, Bernhard Ruh, Han Sah and Klaus Dieter Semmler for many stimulating conversations.

2. RIGID MOTIONS

In this section we fix the notation and collect the necessary (and hopefully sufficient) rudiments from Linear Algebra.

We consider \mathbf{R}^n as an euclidean vector space with the standard inner product. We use $|x|$ to denote the length of a vector $x \in \mathbf{R}^n$, and $\sphericalangle(x, y) \in [0, \pi]$ to denote the angle between two vectors. A rigid motion α (isometry of \mathbf{R}^n) will be expressed in the form

$$x \mapsto \alpha x = Ax + a \quad (x \in \mathbf{R}^n)$$

where $A = \text{rot } \alpha \in O(n)$ is an orthogonal map, called the *rotation part* of α , and $\alpha = \text{trans } \alpha \in \mathbf{R}^n$ is a vector, called the *translation part*.

2.1. *The commutator* $[\alpha, \beta]$ of two rigid motions $x \mapsto \alpha x = Ax + a$ and $x \mapsto \beta x = Bx + b$ is defined as $[\alpha, \beta] = \alpha\beta\alpha^{-1}\beta^{-1}$. The following formulae are easily checked:

$$\text{rot } [\alpha, \beta] = [A, B]$$

$$\text{trans } [\alpha, \beta] = (A - id)b + (id - [A, B])b + A(id - B)A^{-1}a.$$

2.2. *Rotations.* For $A \in O(n)$ we define

$$m(A) = \max \{ |Ax - x| / |x| \mid x \in \mathbf{R}^n \setminus \{0\} \}.$$

Note that $|Ax - x| \leq m(A)|x|$ for $x \in \mathbf{R}^n$. The set

$$(i) \quad E_A = \{ x \in \mathbf{R}^n \mid |Ax - x| = m(A)|x| \}$$

is a non trivial A -invariant subspace. This is immediately checked except perhaps for the part " $x, y \in E_A$ implies $x \pm y \in E_A$ ". This part follows from the equation

$$2m^2(A)(|x|^2 + |y|^2) = 2(|Ax - x|^2 + |Ay - y|^2) = |A(x+y) - (x+y)|^2 + |A(x-y) - (x-y)|^2 \leq m^2(A)(|x+y|^2 + |x-y|^2) = 2m^2(A)(|x|^2 + |y|^2)$$