

V. Necessary conditions in D for $L_c = L_{\{\bot\}}$

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **31 (1985)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **16.07.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

IV.3. THE CLASS \mathcal{D} OF DEFINITE SPACES. Positive definite forms over ordered fields satisfy the triangle inequality as well as the Cauchy-Schwarz inequality. We therefore set down

Definition 15. A definite space is a nondegenerate hermitean space $(\mathfrak{E}; \langle \cdot, \cdot \rangle)$ over an involutorial division ring $(k, *)$, $\text{char } k \neq 2$, that is equipped with a $*$ -valuation φ that has $\varphi(2) = 0$ (cf. Remark 35) and that satisfies one (and hence all) of the four statements in Lemma 14. A definite space \mathfrak{E} will always be considered as a topological vector space, the topology being given by the zero-neighbourhood basis $\mathcal{U}_\gamma := \{\eta \in \mathfrak{E} \mid \varphi\langle \eta \rangle \geq \gamma\}$, $\gamma \in \Gamma$. If $(e_\iota)_{\iota \in I}$ is any family over vectors in \mathfrak{E} such that the net of all finite ("partial") sums $\sum e_\iota$ has a limit x in \mathfrak{E} then we write $x = \sum_{\iota \in I} e_\iota$ and call $(e_\iota)_{\iota \in I}$ summable.

LEMMA 16. Let $(e_\iota)_{\iota \in I}$ be an orthogonal family in the definite space $(\mathfrak{E}; \langle \cdot, \cdot \rangle)$ and \mathfrak{F} its span. For each x in the topological closure of \mathfrak{F} we have $x = \sum_{\iota \in I} \langle x, e_\iota \rangle \langle e_\iota \rangle^{-1} e_\iota$.

Proof. Let \mathcal{P} be the set of all finite subsets of I . For $V \in \mathcal{P}$ we set $x_V := \sum_{\iota \in V} \langle x, e_\iota \rangle \langle e_\iota \rangle^{-1} e_\iota$. We have to prove that for each $\gamma \in \Gamma$ there is $U \in \mathcal{P}$ such that $\varphi\langle x - x_V \rangle \geq \varepsilon$ for all V with $U \subset V \in \mathcal{P}$. Now there is $\eta \in \mathfrak{F}$ with $\varphi\langle x - \eta \rangle \geq \varepsilon$. Pick $U \in \mathcal{P}$ with $\eta \in \text{span}\{e_\iota \mid \iota \in U\}$. If $U \subset V \in \mathcal{P}$ then $x - x_V \perp x_V - \eta$, so by "Pythagoras" (Lemma 14 (ii)) we obtain $\varepsilon \leq \varphi\langle x - \eta \rangle = \min\{\varphi\langle x - x_V \rangle, \varphi\langle x_V - \eta \rangle\} \leq \varphi\langle x - x_V \rangle$. \square

V. NECESSARY CONDITIONS IN \mathcal{D} FOR $L_c = L_{\perp\perp}$

The principal result of this section is

THEOREM 17 ([20]). Let \mathfrak{E} be an infinite dimensional definite space carrying an admissible topology i.e., the topology mentioned in Definition 15 is admissible in the sense of Definition 1; let furthermore $(e_\iota)_{\iota \in I}$ be an orthogonal family in \mathfrak{E} such that $(\varphi\langle e_\iota \rangle)_{\iota \in I}$ has a lower bound in Γ . Then $\sum_{\iota \in I} e_\iota$ exists.

Proof. Let $\mathfrak{F} := \text{span}\{\langle e_\iota \rangle^{-1} e_\iota - \langle e_0 \rangle^{-1} e_0 \mid \iota \in I\}$. We first wish to show that $\langle e_0 \rangle^{-1} e_0$ is not an element of the topological closure $\overline{\mathfrak{F}}$. Indeed,

if γ is a lower bound of $(\varphi\langle e_i \rangle)_{i \in I}$ and if we let $x := \sum_{i \in U} \lambda_i \langle e_i \rangle^{-1} e_i - \langle e_0 \rangle^{-1} e_0$ be a typical vector of \mathfrak{F} (U some finite nonvoid subset of $I \setminus \{0\}$) then we get the inequalities

$$\begin{aligned} \varphi\langle x - \langle e_0 \rangle^{-1} e_0 \rangle &= \varphi\langle (-1 - \sum_U \lambda_i) \langle e_0 \rangle^{-1} e_0 + \sum_U \lambda_i \langle e_i \rangle^{-1} e_i \rangle \\ &= \min_{i \in U} \{2\varphi(-1 - \sum_U \lambda_i) - \varphi\langle e_0 \rangle, 2\varphi(\lambda_i) - \varphi\langle e_i \rangle\} \\ &\leq 2 \min_{i \in U} \{\varphi(-1 - \sum_U \lambda_i), \varphi(\lambda_i)\} - \gamma \leq \varphi(-1) - \gamma = -\gamma. \end{aligned}$$

Thus $\overline{\mathfrak{F}} \neq \mathfrak{E}$.

Since $\mathfrak{F}^{\perp\perp} = \overline{\mathfrak{F}}$ we have $\mathfrak{F}^{\perp} \neq (0)$. Pick a non-zero $x \in \mathfrak{F}^{\perp}$; so $\langle x, e_i \rangle \langle e_i \rangle^{-1} = \langle x, e_0 \rangle \langle e_0 \rangle^{-1}$. If we assume that $(e_i)_{i \in I}$ is a maximal orthogonal family then by $L_c = L_s$ and Lemma 16 $x = \sum_I \langle x, e_i \rangle \langle e_i \rangle^{-1} e_i = \langle x, e_0 \rangle \langle e_0 \rangle^{-1} \sum_I e_i$ and thus $\sum_I e_i \in \mathfrak{F}^{\perp}$. If $(e_i)_{i \in I}$ is not maximal then we write it as a difference of two maximal bounded families: Complete the given family to a maximal orthogonal bounded family $(e_i)_{i \in J}$, $J \supset I$, by Zorn's Lemma. For $i \in J$ let $\alpha_i := 1 \in k$ when $i \in I$ and $\alpha_i := 2$ when $i \in J \setminus I$. The two families $(2e_i)_{i \in J}$, $(\alpha_i e_i)_{i \in J}$ are bounded maximal families to which the previous result may be applied. We get $\sum_{i \in I} e_i = \sum_{i \in I} (2e_i) - \sum_{i \in I} \alpha_i e_i \in \mathfrak{E}$. \square

COROLLARY 18. If \mathfrak{E} and $(e_i)_{i \in I}$ are as in Theorem 17 then $(e_i)_{i \in I}$ converges to $0 \in \mathfrak{E}$. \square

COROLLARY 19. If \mathfrak{E} is as in Theorem 17 then the cofinality type of Γ is ω_0 . In particular, the topology on \mathfrak{E} satisfies the first countability axiom. \square

COROLLARY 20. If \mathfrak{E} is as in Theorem 17 then all orthogonal families of non-zero vectors are countable.

Proof. Let $(e_i)_{i \in I}$ be such a family; by multiplying e_i by a suitable scalar, if necessary, we may assume $(\varphi\langle e_i \rangle)_{i \in I}$ to be bounded below. Since $\sum_{i \in I} e_i$ exists by Theorem 17, the sets $I_\gamma = \{i \in I \mid \varphi\langle e_i \rangle \leq \gamma\}$ are finite for all $\gamma \in \Gamma$. Let $(\gamma_i)_{i \in \mathbb{N}}$ be confinal in Γ . Then $I = \cup \{I_{\gamma_i} \mid i \in \mathbb{N}\}$ is countable. \square

Definition 21. The elements of the group $\Gamma/2\Gamma$ are called *types*. Let $T: \Gamma \rightarrow \Gamma/2\Gamma$ be the canonical projection. $T \circ \varphi$ is constant on the square classes of k (elements of k/k^2) and $T \circ \varphi \circ \langle \rangle$ is constant on the "punctured"

straight lines in E . A family $(e_l)_{l \in I}$ of vectors in \mathfrak{E} is said to satisfy the *type-condition* iff for all $(\alpha_l)_{l \in I} \in k^I$ the following holds: if $(\varphi\langle\alpha_l e_l\rangle)_{l \in I}$ is bounded (below) then $(\alpha_l e_l)_{l \in I}$ converges to $0 \in E$.

COROLLARY 22. *Let \mathfrak{E} be as in Theorem 17. $\Gamma/2\Gamma$ is infinite. Each orthogonal family in \mathfrak{E} satisfies the type-condition, equivalently, $\Gamma/2\Gamma$ satisfies (8) below. \square*

COROLLARY 23. *Let \mathfrak{E} be as in Theorem 17. Then k is complete.*

Proof. By Corollary 19 it suffices to show that a sequence $(\alpha_i)_{i \in \mathbb{N}}$ with limit $0 \in k$ is summable. Let $(e_i)_{i \in \mathbb{N}}$ be maximal orthogonal in \mathfrak{E} with $(\varphi\langle e_i \rangle)_{i \in \mathbb{N}}$ bounded below. If $(\lambda_i)_{i \in \mathbb{N}} \in k^{\mathbb{N}}$ has $(\varphi(\lambda_i))_{i \in \mathbb{N}}$ bounded below then $(\lambda_i e_i)_{i \in \mathbb{N}}$ is summable and by continuity of $\langle \cdot, \cdot \rangle$ we obtain

$$\langle \sum_{\mathbb{N}} \lambda_i e_i, \sum_{\mathbb{N}} e_i \rangle = \sum_{\mathbb{N}} \lambda_i \langle e_i \rangle.$$

Thus, all families $(\lambda_i \langle e_i \rangle)_{i \in \mathbb{N}}$ with bounded $(\lambda_i)_{i \in \mathbb{N}}$ are summable.

Pick a strictly monotonic sequence $(n_i)_{i \in \mathbb{N}} \in \mathbb{N}^{\mathbb{N}}$ with $u_0 = 0$ and for all $i \in \mathbb{N}^+$ and all $m \geq n_i$: $\varphi(\alpha_m) > \varphi\langle e_i \rangle$, and set $A_i := \sum \{\alpha_j \mid n_i \leq j < n_{i+1}\}$. The family $(A_i)_{i \in \mathbb{N}}$ is summable if and only if $(\alpha_i)_{i \in \mathbb{N}}$ summable and, if the sums exist, these must be equal. If we set $\lambda_i := A_i \langle e_i \rangle^{-1}$ then, by what we have shown, the family of the $A_i = \lambda_i \langle e_i \rangle$ is summable. \square

COROLLARY 24. *Let \mathfrak{E} be as in Theorem 17. Then \mathfrak{E} is complete.*

Proof. Let $(x_i)_{i \in \mathbb{N}}$ be a Cauchy sequence (Corollary 19). For each fixed $\eta \in \mathfrak{E}$ the map $x \mapsto \langle \eta, x \rangle$ is uniformly continuous. Hence by Cor. 23 the map $f: \eta \mapsto \lim_i \langle \eta, x_i \rangle$ is well-defined. As it is a continuous linear map, its kernel is a closed hyper-plane and so $(L_c(\mathfrak{E}) = L_{\perp\perp}(\mathfrak{E}))$ there is $a \in \mathfrak{E}$ such that $f(\eta) = \langle x, a \rangle$. Let $N \subseteq \mathbb{N}$ be infinite. Because $\lim \varphi\langle \eta, a - x_i \rangle = \infty$ for all $\eta \in \mathfrak{E}$ it follows by systematic use of the Cauchy-Schwarz inequality that $\{\varphi\langle a - x_i \rangle \mid i \in N\}$ is not bounded above by any $\gamma \in \Gamma$. Therefore $(x_i)_{i \in \mathbb{N}}$ converges to a .

VI. SUFFICIENT CONDITIONS IN \mathscr{D} FOR $L_c = L_{\perp\perp}$

VI.1. ASSUMPTIONS. In this chapter $(\mathfrak{E}; \langle \cdot, \cdot \rangle)$ is a definite space in the sense of Definition 15. Of the base field k we shall furthermore assume