

Zeitschrift: L'Enseignement Mathématique
Band: 35 (1989)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: SIMPLE NASH-MOSER IMPLICIT FUNCTION THEOREM
Anhang: Appendix : Construction of the Smoothing Operators in Sobolev Spaces
Autor: Raymond, Xavier Saint
DOI: <https://doi.org/10.5169/seals-57374>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 17.11.2024

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

If $\phi \in H^\infty(\Omega)$ is valued in $\mathbf{R}^{n(n+1)/2}$, let us consider it as a function valued in $\mathbf{R}^{n(n+3)/2}$ by adding n components $\phi_j = 0$ for $1 \leq j \leq n$, and define $\psi(u)\phi$ as a continuous extension to \mathbf{R}^n of the function

$$(16) \quad v = -\frac{1}{2} A(u)^{-1} \phi$$

where $A(u)$ is the $n(n+3)/2$ square matrix the rows of which are $\partial_j u$ for $1 \leq j \leq n$ and $\partial_j \partial_k u$ for $1 \leq j \leq k \leq n$; thanks to our choice of u_0 , the matrix $A(u_0)$ is invertible on Ω , and so is $A(u)$ for any u close enough to u_0 . Since $A(u)^{-1}$ is an algebraic function of derivatives of u up to order 2, estimates such as (3) are again classical.

Finally, we have to prove that this operator ψ inverts ϕ' (formula (2)). Applying $A(u)$ to the function v in (16), one gets

$$\begin{aligned} \langle \partial_j u, v \rangle &= -\frac{1}{2} \phi_j = 0 & 1 \leq j \leq n \\ \langle \partial_j \partial_k u, v \rangle &= -\frac{1}{2} \phi_{jk} & 1 \leq j \leq k \leq n. \end{aligned}$$

The x_k derivative of the first equation gives $\langle \partial_j \partial_k u, v \rangle + \langle \partial_j u, \partial_k v \rangle = 0$, and one gets also $\langle \partial_j \partial_k u, v \rangle + \langle \partial_k u, \partial_j v \rangle = 0$ so that the second equation and (15) give $\phi'(u)v = \phi$ in Ω .

Thus all the assumptions of the theorem are fulfilled, and it follows that we can get a solution if $\phi(u_0)$ is sufficiently small in some $H^s(\Omega)$ norm; but according to (14), $\phi(u_0) = g^0 - g$, and the result is that (13) can be solved for any metric g close enough to g^0 , as required.

APPENDIX:

CONSTRUCTION OF THE SMOOTHING OPERATORS IN SOBOLEV SPACES

Let us recall that $v \in H^s(\mathbf{R}^n)$ means $v \in \mathcal{S}'(\mathbf{R}^n)$ and

$$|v|_s^2 = (2\pi)^{-n} \int (1 + |\xi|^2)^s |\hat{v}(\xi)|^2 d\xi < \infty.$$

Let $\chi: \mathbf{R}^n \rightarrow [0, 1]$ be a C^∞ function taking the value 1 in a neighborhood of 0 and vanishing for $|\xi| \geq \sqrt{3}$. For $v \in H^\infty(\mathbf{R}^n)$ and $\theta > 1$ one sets

$$\widehat{S_\theta v}(\xi) = \chi(\xi/\theta) \hat{v}(\xi).$$

Then, if $s \geq t$,

$$\begin{aligned} (1 + |\xi|^2)^s |\widehat{S_\theta v}(\xi)|^2 &\leq \theta^{2(s-t)} (1 + |\xi/\theta|^2)^{s-t} |\chi(\xi/\theta)|^2 (1 + |\xi|^2)^t |\hat{v}(\xi)|^2 \\ &\leq (2\theta)^{2(s-t)} (1 + |\xi|^2)^t |\hat{v}(\xi)|^2 \end{aligned}$$

since $|\chi| \leq 1$ and $|\xi/\theta| \leq \sqrt{3}$ for $(\xi/\theta) \in \text{supp } \chi$; this gives the first estimate (4) with $C_{s,t} = 2^{s-t}$.

Similarly, for $s \leq t$,

$$(1 + |\xi|^2)^s |\hat{v}(\xi) - \widehat{S_\theta v}(\xi)|^2 = |1 - \chi(\xi/\theta)|^2 (1 + |\xi|^2)^s |\hat{v}(\xi)|^2;$$

a Taylor formula gives $|1 - \chi(\xi/\theta)| \leq C_k |\xi/\theta|^k$ with $C_k = \sup |\chi^{(k)}|/k!$ for any $k \in \mathbb{N}$ since $\chi(0) = 1$ and $\chi^{(j)}(0) = 0$ for $j > 0$, so that for $t = s + k$

$$\begin{aligned} (1 + |\xi|^2)^s |\hat{v}(\xi) - \widehat{S_\theta v}(\xi)|^2 &\leq C_{t-s}^2 |\xi/\theta|^{2(t-s)} (1 + |\xi|^2)^s |\hat{v}(\xi)|^2 \\ &\leq C_{t-s}^2 \theta^{2(s-t)} (1 + |\xi|^2)^t |\hat{v}(\xi)|^2 \end{aligned}$$

whence the second estimate (4) with $C_{s,t} = C_{t-s} = \sup |\chi^{(t-s)}|/(t-s)!$

REFERENCES

- [1] HAMILTON, R. The inverse function theorem of Nash-Moser. *Bulletin of the A.M.S.* 7 (1982), 65-222.
- [2] HÖRMANDER, L. The boundary problems of physical geodesy. *Arch. Rat. Mech. Anal.* 62 (1976), 1-52.
- [3] ——— *Implicit function theorems*. Lectures at Stanford University, Summer Quarter 1977.
- [4] ——— On the Nash-Moser implicit function theorem. *Annales Acad. Sci. Fenniae, Series A.I. Math.* 10 (1985), 255-259.
- [5] MOSER, J. A new technique for the construction of solutions of nonlinear differential equations. *Proc. Nat. Acad. Sci.* 47 (1961), 1824-1831.
- [6] ——— A rapidly convergent iteration method and nonlinear partial differential equations I and II. *Ann. Scuola Norm. Sup. di Pisa* 20 (1966), 265-315 and 499-533.
- [7] NASH, J. The imbedding problem for Riemannian manifolds. *Ann. of Math.* 63 (1956), 20-63.
- [8] SCHWARTZ, J. T. *Nonlinear functional analysis, Chap. II.A*. Gordon & Breach, New York 1969.
- [9] SERGERAERT, F. Une généralisation du théorème des fonctions implicites de Nash. *C. R. Acad. Sci. Paris, 270A* (1970), 861-863.
- [10] ——— Un théorème des fonctions implicites sur certains espaces de Fréchet et quelques applications. *Ann. Sci. Ec. Norm. Sup. Paris 4^e série*, 5 (1972), 599-660.
- [11] ZEHNDER, E. Generalized implicit function theorems with applications to some small divisor problems I and II. *Comm. in Pure and Appl. Math.* 28 (1975), 91-140; 29 (1976), 49-111.

(Reçu le 14 juin 1989)

Xavier Saint Raymond

Purdue University and C.N.R.S.