

6. The analytic proof

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **36 (1990)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **09.08.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

that the image of $t - s$ in $J(C)$ is a constant section which completes the proof. \square

6. THE ANALYTIC PROOF

In this section we will suppose $K = \mathbf{C}$.

a. *The Poincaré Lemma*

Suppose (\mathcal{S}, ∇) is a sheaf on S^{an} with integrable connection. Then by the Poincaré lemma for integrable connections, it follows that the complex of sheaves

$$\mathcal{S} \xrightarrow{\nabla} \Omega_{San}^1 \otimes \mathcal{S} \xrightarrow{\nabla} \Omega_{San}^2 \otimes \mathcal{S} \xrightarrow{\nabla} \dots$$

is a resolution of the sheaf \mathcal{S}^∇ . Hence,

PROPOSITION 1.6.1. $H^i(\mathcal{S}, \nabla)$ is naturally isomorphic to $H^i(S, \mathcal{S}^\nabla)$.

Remark. As in Proposition 1.1.1, $H^1(\mathcal{S}, \nabla)$ is isomorphic to $\text{Ext}(\mathcal{S}^\vee, \mathcal{O}_{San})$. We can describe the isomorphism from $H^1(\mathcal{S}, \nabla)$ to $H^1(S, \mathcal{S}^\nabla)$ explicitly as follows: Let h be an element of $H^1(\mathcal{S}, \nabla)$. Let \mathcal{L} is a covering of S by open disks. Suppose \mathcal{E} is an extension of \mathcal{S}^\vee by \mathcal{O}_{San} corresponding to h . Then \mathcal{E}^\vee is an extension of \mathcal{O}_{San} by \mathcal{S} . For each $U \in \mathcal{L}$, there exists an $s_U \in \mathcal{E}^\vee(U)^\nabla$ which maps to 1 in $\mathcal{O}_{San}(U)$. Then the image h in $H^1(S, \mathcal{S}^\nabla)$ is the class of the cocycle $\{(U, V) \rightarrow s_U - s_V\}$.

Suppose, X is a smooth proper S -scheme and Z is a subscheme of X which is either empty or finite over S . We will define the Betti homology sheaf $\mathcal{H}_i(X/S, Z, \mathbf{Z})$ on S^{an} as follows. If Z is smooth over S , we define $\mathcal{H}_i(X/S, Z, \mathbf{Z})$ to be the sheaf associated to the presheaf

$$U \rightarrow H_i(f^{-1}(U), f^{-1}(U) \cap Z, \mathbf{Z}),$$

(this latter group is the Betti homology of $f^{-1}(U)$ relative to $f^{-1}(U) \cap Z$). More generally, let S' be a non-empty affine open subset of S such that $Z' = Z \times_S S'$ is étale over S' . Let $X' = X \times_S S'$ and let ι denote the inclusion morphisms $X' \rightarrow X, Z' \rightarrow Z$ and $S' \rightarrow S$. We set

$$\mathcal{H}_i(X/S, Z, \mathbf{Z}) = \iota_* \mathcal{H}_i(X'/S', Z', \mathbf{Z}).$$

This is independent of the choice of S' . We also set

$$\mathcal{H}_i(X/S, \mathbf{Z}) = \mathcal{H}_i(X/S, \emptyset, \mathbf{Z}) \text{ and } \mathcal{H}_1(X/S, Z, \mathbf{C}) = \mathcal{H}_1(X/S, Z, \mathbf{Z}) \otimes \underline{\mathbf{C}}.$$

Suppose s and t are two distinct sections of X/S and $Z = s \cup t$. Suppose S' is an affine open of S such that Z' is étale over S' in the notation of the previous paragraph. We have exact sequences

$$0 \rightarrow \mathcal{H}_1(X/S, \mathbf{Z}) \rightarrow \mathcal{H}_1(X/S, \mathbf{Z}, \mathbf{Z}) \rightarrow \iota_* \mathcal{H}_0(Z'/S', \mathbf{Z}) \rightarrow \mathcal{H}_0(X/S, \mathbf{Z}) \rightarrow 0 .$$

and

$$0 \rightarrow \mathcal{H}_0(S'/S', \mathbf{Z}) \rightarrow \mathcal{H}_0(Z'/S', \mathbf{Z}) \rightarrow \mathcal{H}_0(X'/S', \mathbf{Z}) ,$$

where the first map is $t_* - s_*$. From which we derive the short exact sequence

$$0 \rightarrow \mathcal{H}_1(X/S, \mathbf{Z}) \rightarrow \mathcal{H}_1(X/S, \mathbf{Z}, \mathbf{Z}) \rightarrow \underline{\mathbf{Z}} \rightarrow 0 .$$

since $\iota_* \underline{\mathbf{Z}}|_{S'^{an}} \cong \underline{\mathbf{Z}}$. In particular, if U is an open disk in S^{an} , we have an exact sequence

$$0 \rightarrow \mathcal{H}_1(X/S, \mathbf{Z})(U) \rightarrow \mathcal{H}_1(X/S, \mathbf{Z}, \mathbf{Z})(U) \rightarrow \mathbf{Z} \rightarrow 0$$

We define the Betti cohomology sheaf $\mathcal{H}^1(X/S, \mathbf{Z}, \mathbf{C})$ in the same way and it is easy to see that $\mathcal{H}^1(X/S, \mathbf{Z}, \mathbf{C}) \cong \text{Hom}(\mathcal{H}_1(X/S, \mathbf{Z}, \mathbf{C}), \underline{\mathbf{C}})$. Also, it is known that if Z is étale over S then $\mathcal{H}^1(X/S, \mathbf{Z}, \mathbf{C}) \cong R_{f*}^1 \mathcal{I}_Z$ where \mathcal{I}_Z is the subsheaf of $\underline{\mathbf{C}}$ whose sections vanish on Z .

Suppose X is proper over S with connected fibers. Let

$$(\mathcal{H}_{DR}^1(X/S, \mathbf{Z}), \nabla) = \mathcal{O}_{S^{an}} \otimes_{\mathcal{O}_S} (H_{DR}^1(X/S, \mathbf{Z}), \nabla) .$$

We claim, for $Z \subseteq X$ finite over S .

$$(\mathcal{H}_{DR}^1(X/S, \mathbf{Z}), \nabla) \cong (\mathcal{O}_{S^{an}} \otimes \mathcal{H}^1(X/S, \mathbf{Z}, \mathbf{C}), d \otimes id)$$

This follows from the relative Poincaré lemma above on S' and hence on all of S since both sides are integrable connections. Hence,

LEMMA 1.6.2. *There is a natural isomorphism*

$$(\mathcal{H}_{DR}^1(X/S, \mathbf{Z})^\vee, \check{\nabla}) \cong (\mathcal{O}_{S^{an}} \otimes \mathcal{H}^1(X/S, \mathbf{Z}, \mathbf{C}), d \otimes id) .$$

In particular

$$H^i(\mathcal{H}_{DR}^1(X/S, \mathbf{Z})^\vee, \check{\nabla}) \cong H^i(S^{an}, \mathcal{H}^1(X/S, \mathbf{Z}, \mathbf{C})) .$$

We conclude, using this, Proposition 1.1.1 and GAGA that

THEOREM 1.6.3. *There exists a natural isomorphism*

$$\beta: \text{Ext}(H_{DR}^1(X/S), \mathcal{O}_S) \rightarrow H^1(S^{an}, \mathcal{H}^1(X/S, \mathbf{C})) .$$

b. *End of Analytic Proof*

Now suppose X is an Abelian scheme over S . We have an exact sequence of sheaves over S^{an} ,

$$0 \rightarrow \mathcal{H}_1(X/S, \mathbf{Z}) \rightarrow \mathcal{L}ie_{X^{an}/S^{an}} \rightarrow \underline{X^{an}} \rightarrow 0 .$$

From the corresponding long exact sequence of cohomology groups we obtain an exact sequence

$$\mathcal{L}ie_{X^{an}/S^{an}}(S^{an}) \rightarrow X^{an}(S^{an}) \xrightarrow{\delta} H^1(S^{an}, \mathcal{H}_1(X/S, \mathbf{Z})) .$$

We may describe $\delta(s)$ as follows: Suppose $e \neq s$. Let $Z = e \cup s$. Then as $f_*(\Omega_{X^{an}/S^{an}}^1)$ maps into $\mathcal{H}_{DR}^1(X/S)$, $\mathcal{H}_1(X/S, \mathbf{Z}, \mathbf{Z})$ maps into

$$f_*(\Omega_{X^{an}/S^{an}}^1)^\vee = \mathcal{L}ie_{X^{an}/S^{an}}$$

so that the diagram

$$\begin{array}{ccc} \mathcal{H}_1(X/S, \mathbf{Z}, \mathbf{Z}) & & \\ \uparrow & \searrow & \\ \mathcal{H}_1(X/S, \mathbf{Z}) & \rightarrow & \mathcal{L}ie_{X^{an}/S^{an}} \end{array}$$

commutes. Let \mathcal{U} be an ordered covering of S by open disks. For each $U \in \mathcal{U}$ let $\gamma_U \in \mathcal{H}_1(X/S, \mathbf{Z}, \mathbf{Z})(U)$ such that $\gamma_U \rightarrow 1$ under the map $\mathcal{H}_1(X/S, \mathbf{Z})(U) \rightarrow \mathbf{Z}$. Then the image of γ_U in $X(U)$ is $s(U)$. Hence $\delta(s)$ is represented by the one cocycle $\{(U, V) \rightarrow \gamma_U - \gamma_V\}$.

Now, it follows from this and the remark after Proposition 1.6.1 that $\beta \circ M$ is equal to the composition of δ and the natural map

$$H^1(S^{an}, \mathcal{H}_1(X/S, \mathbf{Z})) \rightarrow H^1(S^{an}, \mathcal{H}_1(X/S, \mathbf{C})) \cong H^1(S^{an}, \mathcal{H}_1(X/S, \mathbf{Z})) \otimes \mathbf{C} .$$

Hence, if $s \in X^{an}(S^{an})$, $M(s) = 0$ iff there exists a positive integer n such that $\delta(ns) = n\delta(s) = 0$. Hence ns is in the image of $\mathcal{L}ie_{X^{an}/S^{an}}(S^{an}) \rightarrow X^{an}(S^{an})$ and so is an infinitely divisible element of $X^{an}(S^{an})$.

Suppose $s \in X(S)$. We claim ns is an infinitely divisible element of $X(S)$. Let m be a positive integer. Let $t \in X^{an}(S^{an})$ such that $mt = ns$. There exists a finite étale Galois covering \tilde{S} of S such that $t \in X(\tilde{S})$. If $\sigma \in \text{Gal}(\tilde{S}/S)$, then $t^\sigma = t$ because $t^\sigma(x) = t(\sigma^{-1}(x))$ for $x \in \tilde{S}(\mathbf{C})$. It follows that $t \in X(S)$. This establishes our claim.

Finally, it follows from the function field Mordell-Weil Theorem [LN] that the image of ns in $X_{\mathbf{C}(S)}(\mathbf{C}(S))$ is a constant section X/S . Theorem 1.4.3 now follows immediately. \square