

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 37 (1991)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON A THEOREM OF SIKORAV
Autor: Brunella, Marco
DOI: <https://doi.org/10.5169/seals-58732>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 15.01.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

ON A THEOREM OF SIKORAV

by Marco BRUNELLA

The aim of this note is to give a new proof of the following theorem of J. C. Sikorav:

THEOREM ([Sik]). *Let M be a closed manifold, let*

$$\phi_t: T^*M \rightarrow T^*M, t \in [0, 1],$$

*be a hamiltonian isotopy and let $L \subset T^*M$ be an immersed lagrangian submanifold with a generating function $S: M \times \mathbf{R}^k \rightarrow \mathbf{R}$ quadratic at infinity, then also $\phi_1(L)$ has a generating function quadratic at infinity.*

Recall that a *generating function* of an immersed lagrangian submanifold $L \subset T^*M$ is a function $S: M \times \mathbf{R}^k \rightarrow \mathbf{R}, (q, \lambda) \mapsto S(q, \lambda)$, such that:

a) $\frac{\partial S}{\partial \lambda}: M \times \mathbf{R}^k \rightarrow \mathbf{R}^k$ is transverse to $0 \in \mathbf{R}^k$, so $\left\{ \frac{\partial S}{\partial \lambda} = 0 \right\} \subset M \times \mathbf{R}^k$

is a submanifold;

b) $L = \{ \xi \in T^*M \mid \exists \lambda \in \mathbf{R}^k: \frac{\partial S}{\partial \lambda}(\pi(\xi), \lambda) = 0, \xi = d(S(\cdot, \lambda))(\pi(\xi)) \}$

where $\pi: T^*M \rightarrow M$ is the canonical projection.

The generating function S is said to be *quadratic at infinity* if for some $R > 0$

$$S(q, \lambda) = Q(\lambda) \quad \forall (q, \lambda) \in M \times \mathbf{R}^k, \|\lambda\| > R$$

where Q is some non-degenerate quadratic form.

A *hamiltonian isotopy* of a symplectic manifold is a smooth curve of symplectic diffeomorphisms $\{ \phi_t \}_{t \in [0, 1]}$ such that

$$\phi_0 = id \quad \text{and} \quad \dot{\phi}_t \stackrel{\text{def}}{=} \frac{d}{ds} \Big|_{s=0} \phi_{t+s} \circ \phi_t^{-1}$$

is a hamiltonian vector field $\forall t \in [0, 1]$.

The above theorem is important in the intersection theory of lagrangian submanifolds.

Our proof starts with the following remark, contained in [Gir] under the name of "Chekanov trick". Let $i: M \rightarrow \mathbf{R}^N$ be any embedding and let \mathbf{R}^N , M be equipped with riemannian metrics such that i is an isometric embedding. These metrics induce isomorphisms $T^*M \simeq TM$ and $T^*\mathbf{R}^N \simeq T\mathbf{R}^N$, so the embedding $Ti: TM \rightarrow T\mathbf{R}^N$ induces an embedding

$$j: T^*M \rightarrow T^*\mathbf{R}^N$$

which is a symplectic embedding. Remark that $j(T^*M)$ is a subbundle of $(T^*\mathbf{R}^N)|_{i(M)}$ and there is a canonical decomposition

$$(T^*\mathbf{R}^N)|_{i(M)} = j(T^*M) \oplus N_{i(M)}^*$$

where $N_{i(M)}^*$ is the conormal bundle of $i(M)$ in \mathbf{R}^N . This decomposition is the dual version of the decomposition $(T\mathbf{R}^N)|_{i(M)} = (Ti)(TM) \oplus N_{i(M)}, N_{i(M)}$ being the normal bundle of $i(M)$ in \mathbf{R}^N .

We want to extend hamiltonian isotopies and lagrangian submanifolds from $T^*M (\simeq j(T^*M))$ to $T^*\mathbf{R}^N$.

LEMMA 1. Let $\phi_t: T^*M \rightarrow T^*M, t \in [0, 1]$, be a hamiltonian isotopy and let $j: T^*M \rightarrow T^*\mathbf{R}^N$ as above, then there exists a hamiltonian isotopy $\psi_t: T^*\mathbf{R}^N \rightarrow T^*\mathbf{R}^N, t \in [0, 1]$, such that $\forall t \in [0, 1]$:

$$1) j \circ \phi_t = \psi_t \circ j$$

$$2) \psi_t \text{ leaves invariant } (T^*\mathbf{R}^N)|_{i(M)}$$

moreover, if $V \subset \mathbf{R}^N$ is any neighborhood of $i(M)$ then we may choose every ψ_t with support contained in $(T^*\mathbf{R}^N)|_V$.

Proof. Let $pr: (T^*\mathbf{R}^N)|_{i(M)} = j(T^*M) \oplus N_{i(M)}^* \rightarrow j(T^*M)$ be the projection on the first factor; $(T^*\mathbf{R}^N)|_{i(M)}$ is a coisotropic submanifold of $T^*\mathbf{R}^N$ and the fibres of pr are its characteristic leaves. For every $x \in j(T^*M)$ let E_x be the antiorthogonal complement of $T_x(j(T^*M))$; E_x is transverse to $T_x(j(T^*M))$ and intersects $T_x((T^*\mathbf{R}^N)|_{i(M)})$ along $T_x(pr^{-1}(x))$. This implies that we may find a tubular neighborhood of $j(T^*M)$ in $T^*\mathbf{R}^N$, $p_0: U \rightarrow j(T^*M)$, such that the fibre $p_0^{-1}(x)$ intersects $(T^*\mathbf{R}^N)|_{i(M)}$ along $pr^{-1}(x)$:

$$p_0^{-1}(x) \cap (T^*\mathbf{R}^N)|_{i(M)} = pr^{-1}(x) \cap U \quad \forall x \in j(T^*M).$$

Let now $\{H_t\}_{t \in [0, 1]}$ be hamiltonians of the vector fields $\{\dot{\phi}_t\}_{t \in [0, 1]}$, define $\forall t \in [0, 1] K_t: U \cup (T^*\mathbf{R}^N)|_{i(M)} \rightarrow \mathbf{R}$ by:

$$K_t(x) = \begin{cases} H_t(j^{-1}(p_0(x))) & \text{if } x \in U \\ H_t(j^{-1}(pr(x))) & \text{if } x \in (T^*\mathbf{R}^N)|_{i(M)}. \end{cases}$$

The relation between p_0 and pr guarantees that K_t are well defined; now extend smoothly the family $\{K_t\}_{t \in [0,1]}$ on all $T^*\mathbf{R}^N$, in such a way that K_t are constant outside $(T^*\mathbf{R}^N)|_V$ (this is possible choosing U such that its projection on \mathbf{R}^N has closure contained in V). Then the hamiltonian isotopy $\psi_t: T^*\mathbf{R}^N \rightarrow T^*\mathbf{R}^N, t \in [0, 1]$, generated by $\{K_t\}_{t \in [0,1]}$ satisfies the conclusions of the lemma. \square

LEMMA 2. *Let $L \subset T^*M$ be an immersed lagrangian submanifold with generating function $S: M \times \mathbf{R}^k \rightarrow \mathbf{R}$ quadratic at infinity, then there exists an immersed lagrangian submanifold $\tilde{L} \subset T^*\mathbf{R}^N$ with a generating function $\tilde{S}: \mathbf{R}^N \times \mathbf{R}^k \rightarrow \mathbf{R}$ quadratic at infinity and such that $\tilde{L} \cap (T^*\mathbf{R}^N)|_{i(M)} = j(L)$ (transversally); moreover, if $V \subset \mathbf{R}^N$ is any neighborhood of $i(M)$ then we may choose \tilde{L} and \tilde{S} such that \tilde{L} is equal to the null section outside $(T^*\mathbf{R}^N)|_V$ and \tilde{S} is equal to Q (= quadratic form associated to S) outside $V \times \mathbf{R}^k$.*

Proof. Let $W \xrightarrow{q_0} i(M)$ be a tubular neighborhood of $i(M)$ in \mathbf{R}^N , with fibres orthogonal to $i(M)$ and $\bar{W} \subset V$; define

$$\tilde{S}: W \times \mathbf{R}^k \rightarrow \mathbf{R} \quad \text{by} \quad \tilde{S}(x, \lambda) = S(i^{-1}(q_0(x)), \lambda)$$

and extend \tilde{S} to all $\mathbf{R}^N \times \mathbf{R}^k$ preserving the quadraticity at infinity and in such a way that $\tilde{S}(x, \lambda) = Q(\lambda)$ outside $V \times \mathbf{R}^k$; a transversality argument allows to find such a \tilde{S} such that it generates a lagrangian submanifold $\tilde{L} \subset T^*\mathbf{R}^N$, and clearly \tilde{L}, \tilde{S} satisfy the conclusions of the lemma. \square

Conversely:

LEMMA 3. *Let $\tilde{L} \subset T^*\mathbf{R}^N, L \subset T^*M$ be immersed lagrangian submanifolds such that $j(L) = \tilde{L} \cap (T^*\mathbf{R}^N)|_{i(M)}$ transversally; if $\tilde{S}: \mathbf{R}^N \times \mathbf{R}^k \rightarrow \mathbf{R}$ is a generating function for \tilde{L} , then $S: M \times \mathbf{R}^k \rightarrow \mathbf{R}, S(x, \lambda) = \tilde{S}(i(x), \lambda)$, is a generating function for L . \square*

Remark that if \tilde{S} is quadratic at infinity, so is S .

Let now $L \subset T^*M, S: M \times \mathbf{R}^k \rightarrow \mathbf{R}, \phi_t: T^*M \rightarrow T^*M, t \in [0, 1]$, as in the hypotheses of the theorem. Let $\psi_t: T^*\mathbf{R}^N \rightarrow T^*\mathbf{R}^N, t \in [0, 1]$, be an extension of ϕ_t as in lemma 1, and let $\tilde{L} \subset T^*\mathbf{R}^N$ be an extension of L as in lemma 2, with generating function $\tilde{S}: \mathbf{R}^N \times \mathbf{R}^k \rightarrow \mathbf{R}$. We have:

$$j(\phi_1(L)) = \psi_1(j(L)) = \psi_1(\tilde{L}) \cap (T^*\mathbf{R}^N)|_{i(M)}$$

and so, by lemma 3, we have only to prove that $\psi_1(\tilde{L})$ has a generating function, quadratic at infinity. If we choose V with compact closure, then the support properties of ψ_t outside $(T^*\mathbf{R}^N)|_V$ allows to “compactify” $\{\psi_t\}_{t \in [0,1]}$, that is there exists a hamiltonian isotopy $\{\tilde{\psi}_t\}_{t \in [0,1]}$ of $T^*\mathbf{R}^N$ with compact support such that $\tilde{\psi}_1(\tilde{L}) = \psi_1(\tilde{L})$ (remark that the quadraticity at infinity of \tilde{S} implies that $\tilde{L} \cap (T^*\mathbf{R}^N)|_V$ is contained in a compact set, and so is $[\cup_{t \in [0,1]} \psi_t(\tilde{L})] \cap (T^*\mathbf{R}^N)|_V$). Observe that \tilde{S} is equal to a quadratic form $Q = Q(\lambda)$ outside a compact set.

We may decompose $\tilde{\psi}_1$ as a product $\tilde{\psi}_1 = g_1 \circ \dots \circ g_l$, where each g_j is a symplectic diffeomorphism with compact support and with a *generating function* $F_j: \mathbf{R}^N \times \mathbf{R}^N \rightarrow \mathbf{R}$:

$$(P, Q) = g_j(p, q) \Leftrightarrow P = p + \frac{\partial F_j}{\partial Q}(Q, p), q = Q + \frac{\partial F_j}{\partial p}(Q, p)$$

(we use here the standard symplectic coordinates on $T^*\mathbf{R}^N$). We may suppose that each F_j has compact support.

The proof of the theorem is achieved by an iteration of the following lemma:

LEMMA 4. *Let $\tilde{L} \subset T^*\mathbf{R}^N$ be an immersed lagrangian submanifold with generating function $\tilde{S}: \mathbf{R}^N \times \mathbf{R}^k \rightarrow \mathbf{R}$ such that outside a compact set $\tilde{S}(x, \lambda) = Q(\lambda) =$ non-degenerate quadratic form, let $g: T^*\mathbf{R}^N \rightarrow T^*\mathbf{R}^N$ be a symplectic diffeomorphism with a generating function $F: \mathbf{R}^N \times \mathbf{R}^N \rightarrow \mathbf{R}$ with compact support; then $g(\tilde{L})$ has a generating function equal to a non-degenerate quadratic form outside a compact set.*

Proof. A computation shows that $T: \mathbf{R}^N \times \mathbf{R}^k \times \mathbf{R}^N \times \mathbf{R}^N \rightarrow \mathbf{R}$ defined by

$$T(x, \lambda, \xi, \eta) = F(x, \xi) + \tilde{S}(x + \eta, \lambda) - \xi \cdot \eta$$

is a generating function for $g(\tilde{L})$ (with λ, ξ, η as parameters). Let $\rho: [0, +\infty) \rightarrow [0, 1]$ be a smooth function s.t. $\rho(t) = 1 \forall t \leq 1$ and $\rho(t) = 0 \forall t \geq 2$, then for an appropriate choice of positive constants K_1, K_2, K_3, K_4 the function

$$\begin{aligned} \hat{T}(x, \lambda, \xi, \eta) &= \rho\left(\frac{\|\lambda\|}{K_1}\right) \rho\left(\frac{\|\eta\|}{K_2}\right) F(x, \xi) \\ &+ \rho\left(\frac{\|\xi\|}{K_3}\right) \rho\left(\frac{\|x\|}{K_4}\right) [\tilde{S}(x + \eta, \lambda) - Q(\lambda)] + Q(\lambda) - \xi \cdot \eta \end{aligned}$$

is again a generating function for $g(\tilde{L})$, and it is equal to the non degenerate quadratic form $Q(\lambda) - \xi \cdot \eta$ outside a compact set. A possible choice of the constants K_j is the following:

$$\text{if } S_0(x, \lambda) \stackrel{\text{def}}{=} \tilde{S}(x, \lambda) - Q(\lambda),$$

$$\text{if } \text{supp } S_0 \subset \{\|x\| < R, \|\lambda\| < R\}, \text{supp } F \subset \{\|x\| < R, \|\xi\| < R\},$$

and

$$\text{if } a = \sup F, \quad b = \sup S_0, \quad c = \sup \left\| \frac{\partial F}{\partial \xi} \right\|, \quad d = \sup \left\| \frac{\partial S_0}{\partial x} \right\|,$$

$$e = \sup \left\| \frac{\partial S_0}{\partial \lambda} \right\|, \quad \beta = \sup \left| \frac{dp}{dt} \right|,$$

then define:

$$K_1 \text{ s.t. } K_1 > R \quad \text{and} \quad \|\lambda\| \geq K_1 \Rightarrow \left\| \frac{\partial Q}{\partial \lambda}(\lambda) \right\| > \frac{1}{K_1} \beta a + e$$

$$K_2 = K_3 = K \text{ s.t. } K > R, \quad K > \frac{1}{K} \beta b + c, \quad K > \frac{1}{K} \beta a + d$$

$$K_4 \text{ s.t. } K_4 > K + R. \quad \square$$

REFERENCES

- [Gir] GIROUX, E. Formes génératrices d'immersions lagrangiennes dans un espace cotangent. *Coll. Int. du Sémin. Sud-Rhodanien de Géométrie 1988*, Springer Lecture Notes 1416 (1990), 139-145.
- [Sik] SIKORAV, J. C. Problèmes d'intersections et de points fixes en géométrie hamiltonienne. *Comm. Math. Helv.* 62 (1987), 62-73.

(Reçu le 27 novembre 1990)

Marco Brunella

S.I.S.S.A.

Strada Costiera 11

34014 Trieste, Italy

Vide-leer-empty