

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 39 (1993)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: JACOBI FORMS AND SIEGEL MODULAR FORMS: RECENT RESULTS AND PROBLEMS
Autor: Kohnen, Winfried
Kapitel: 1.1. Siegel modular forms of genus 2
DOI: <https://doi.org/10.5169/seals-60416>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 22.12.2024

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

JACOBI FORMS AND SIEGEL MODULAR FORMS: RECENT RESULTS AND PROBLEMS

by Winfried KOHNEN

INTRODUCTION

In the present paper we would like to describe some recent developments how Jacobi forms can be used to study Siegel modular forms of genus 2 and what problems arise in this way. After a preliminary section on Siegel modular forms and Jacobi forms (§ 1) which mainly serves to fix some notation, we shall discuss the so called Maass space in § 2. We shall then study relations between Jacobi forms and spinor zeta functions of Hecke eigenforms of genus 2 (§ 3) and finally in § 4 will indicate how Jacobi forms can be used to give estimates for Fourier coefficients of Siegel cusp forms.

Sections 2-4 are divided into two parts: part one describes known results while part two gives some open problems.

We do not go here into any more intrinsic properties of Jacobi forms (as e.g. the trace formula or relations to modular forms of integral weight) nor discuss any representation-theoretic aspects of the theory. For good surveys, we refer to [33, 36] for the first and to [3] for the second topic.

§ 1. PRELIMINARIES ON SIEGEL MODULAR FORMS AND JACOBI FORMS

1.1. SIEGEL MODULAR FORMS OF GENUS 2

We write \mathcal{H}_2 for the Siegel upper half-space of genus 2. The natural action of $\mathrm{Sp}_2(\mathbf{R})$ on \mathcal{H}_2 is denoted by

$$(M, Z) \mapsto M \langle Z \rangle := (AZ + B)(CZ + D)^{-1}$$

$$\left(M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \mathrm{Sp}_2(\mathbf{R}), \quad Z \in \mathcal{H}_2 \right).$$

We put $\Gamma_2 := \mathrm{Sp}_2(\mathbf{Z})$ and for $k \in \mathbf{Z}$ denote by $M_k(\Gamma_2)$ the space of Siegel modular forms of weight k on Γ_2 , i.e. the space of holomorphic func-

tions $F: \mathcal{H}_2 \rightarrow \mathbf{C}$ satisfying $F(M \langle Z \rangle) = \det(CZ + D)^k F(Z)$ for all $M = \begin{pmatrix} \cdot & \cdot \\ C & D \end{pmatrix} \in \Gamma_2$. Such a function has a Fourier expansion

$$F(Z) = \sum_{T = T' \geq 0} a(T) e^{2\pi i \operatorname{tr}(TZ)}$$

where T runs over all positive semi-definite half-integral $(2, 2)$ -matrices. We write $S_k(\Gamma_2)$ for the subspace of cusp forms (require $a(T) = 0$ for $T \not\geq 0$).

For $F, G \in S_k(\Gamma_2)$ we denote by

$$\langle F, G \rangle = \int_{\Gamma_2 \backslash \mathcal{H}_2} F(Z) \overline{G(Z)} (\det Y)^{k-3} dX dY \quad (X = \operatorname{Re}(Z), Y = \operatorname{Im}(Z))$$

the Petersson scalar product of F and G .

For basic facts on Siegel modular forms we refer to [12, 17].

1.2. JACOBI FORMS

We write \mathcal{H} for the complex upper half-plane. We let $H(\mathbf{R})$ be the Heisenberg group, i.e. the set of triples $((\lambda, \mu), \kappa) \in \mathbf{R}^2 \times \mathbf{R}$ with group law $((\lambda, \mu), \kappa) ((\lambda', \mu'), \kappa') = ((\lambda + \lambda', \mu + \mu'), \kappa + \kappa' + \lambda\mu' - \lambda'\mu)$, and denote by $G^J := SL_2(\mathbf{R}) \times H(\mathbf{R})$ the Jacobi group where $SL_2(\mathbf{R})$ operates on $H(\mathbf{R})$ from the right by $((\lambda, \mu), \kappa)M = ((\lambda, \mu)M, \kappa)$. The group G^J acts on $\mathcal{H} \times \mathbf{C}$ by

$$\left(\begin{pmatrix} a & b \\ c & d \end{pmatrix}, ((\lambda, \mu), \kappa) \right) \circ (\tau, z) = \left(\frac{a\tau + b}{c\tau + d}, \frac{z + \lambda\tau + \mu}{c\tau + d} \right).$$

We set $\Gamma_1 := SL_2(\mathbf{Z})$, $\Gamma_1^J := \Gamma_1 \times H(\mathbf{Z})$ and for $k \in \mathbf{Z}$ and $m \in \mathbf{N}_0$ denote by $J_{k,m}$ the space of Jacobi forms of weight k and index m on Γ_1^J , i.e. the space of holomorphic functions $\phi: \mathcal{H} \times \mathbf{C} \rightarrow \mathbf{C}$ satisfying the transformation formula

$$\phi(\gamma \circ (\tau, z)) = (c\tau + d)^k \exp \left(2\pi i m \left(\frac{c(z + \lambda\tau + \mu)^2}{c\tau + d} - \lambda^2\tau - 2\lambda z \right) \right) \phi(\tau, z)$$

for all $\gamma = \left(\begin{pmatrix} a & b \\ c & d \end{pmatrix}, ((\lambda, \mu), \kappa) \right) \in \Gamma_1^J$ and having a Fourier expansion

$$\phi(\tau, z) = \sum_{n, r \in \mathbf{Z}, r^2 \leq 4mn} c(n, r) q^n \zeta^r$$

where $q = e^{2\pi i \tau}$, $\zeta = e^{2\pi i z}$. We write $J_{k,m}^{\text{cusp}}$ for the subspace of cusp forms (require $c(n, r) = 0$ for $r^2 = 4mn$). Note that the coefficients $c(n, r)$ depend