

6. The number of proper 4-colorings of a graph

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **40 (1994)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **12.07.2024**


Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

For example, all elements of weight 3 in $K_{m,n}$ can be represented (up to proper size and location) by the following picture:



or its vertical analogue. This picture represents a codeword of the form

$$E(k, l; r_1, r_2) + E(k, l; r_1, r_3) + E(k, l; r_2, r_3) .$$

Thus, the number of codewords of weight 3 in $K_{m,n}$ is equal to

$$w_3(K_{m,n}) = \binom{m}{2} \binom{n}{3} + \binom{m}{3} \binom{n}{2} .$$

Similarly, one can show that

$$w_4(K_{m,n}) = 3 \binom{m}{2} \binom{n}{4} + 9 \binom{m}{3} \binom{n}{3} + 3 \binom{m}{4} \binom{n}{2} ;$$

$$w_5(K_{m,n}) = 12 \binom{m}{2} \binom{n}{5} + 72 \binom{m}{3} \binom{n}{4} + 72 \binom{m}{4} \binom{n}{3} + 12 \binom{m}{2} \binom{n}{5} + 9 \binom{m}{3} \binom{n}{3} .$$

As a last remark, note that an upper bound for the weights in the associated code L_f is given by $\frac{1}{8}n^3(n-1)$, and that this bound is actually attained for some n if and only if there exists a Hadamard matrix of order n . This follows from, say, Corollary 3.

6. THE NUMBER OF PROPER 4-COLORINGS OF A GRAPH

Let $G = (V, E)$ be a simple graph (no loops, no multiple edges) with vertex set V and edge set E . We will identify V with $\{1, \dots, n\}$, and denote the cardinality of E by e .

A 4-coloring of G is the assignment to every vertex of one among four fixed colors; such a coloring is *proper* if the colors assigned to the end vertices of any edge are distinct. For a survey on the 4-colorings of planar graphs, see [SK].

We will count the number of proper 4-colorings of G , in terms of the weight enumerator of a certain code of length $3e$.

STEP 1. *The defining equations for proper 4-colorings.*

As our palette of colors, we will choose the 4-set $\{1, -1\}^2$. The space of all 4-colorings of G can thus be identified with $\{1, -1\}^{2n}$, for example as follows:

Convention. If $p = (p_1, \dots, p_{2n}) \in \{1, -1\}^{2n}$, then the pair (p_i, p_{i+n}) represents the color assigned to vertex i , for $i = 1, \dots, n$.

Consider now $2n$ variables x_1, \dots, x_{2n} , and define

$$g_{i,j} := (1 + x_i x_j) (1 + x_{i+n} x_{j+n})$$

for $1 \leq i, j \leq n$.

If p is a 4-coloring of G , then $g_{i,j}(p) = 0$ if and only if the colors assigned to vertices i and j are distinct.

Thus, a 4-coloring p of G is proper if and only if

$$g_{i,j}(p) = 0 \quad \text{for all } (i,j) \in E .$$

STEP 2. *Reduction to a single equation.*

Let

$$g := \sum_{(i,j) \in E} g_{i,j}^2 .$$

By construction, g satisfies the following properties:

- (1) $g(p) \geq 0$ for all 4-colorings p ;
- (2) $g(p) = 0$ if and only if p is proper.

Developing the expression for g , we obtain:

$$\begin{aligned} g &= \sum_E g_{i,j}^2 \\ &= \sum_E (1 + x_i x_j + x_{i+n} x_{j+n} + x_i x_j x_{i+n} x_{j+n})^2 \\ &= 4 \sum_E (1 + x_i x_j + x_{i+n} x_{j+n} + x_i x_j x_{i+n} x_{j+n}) \\ &= 4e + 4f, \end{aligned}$$

where

$$f := \sum_E (x_i x_j + x_{i+n} x_{j+n} + x_i x_j x_{i+n} x_{j+n}) .$$

(Here again, the computation was performed modulo $x_i^2 = 1$ for all i .)

Obviously, f satisfies the following properties:

- (1) $f(p) \geq -e$ for all 4-colorings p ;
- (2) $f(p) = -e$ if and only if p is proper.

STEP 3. *The code associated with f .*

Let $K_G := L_f^\perp$ be the dual of the code L_f associated with f . To describe it, we consider the map

$$\begin{aligned}\phi_G: \mathbf{F}_2^{3e} &\rightarrow \mathbf{F}_2^{2n} \\ E_{i,j} &\mapsto e_i + e_j \\ E_{i+n,j+n} &\mapsto e_{i+n} + e_{j+n} \\ E_{i+2n,j+2n} &\mapsto e_i + e_j + e_{i+n} + e_{j+n} \quad ((i,j) \in E)\end{aligned}$$

where $\{E_{i,j}, E_{i+n,j+n}, E_{i+2n,j+2n}\}_{(i,j) \in E}$ and $\{e_i\}_{i \in V}$ denote the standard bases of the left and right spaces, respectively. By construction, $K_G = \text{Ker}(\phi_G)$.

Here again, as a direct consequence of Theorem 4 and of the stated properties of f , we have:

THEOREM 7. *Let K_G be the code of length $3e$ defined above, and let $P_G(T)$ denote its weight enumerator. Then, the number $\chi_G(4)$ of proper 4-colorings of G , is given by*

$$\chi_G(4) = \frac{1}{4^{e-n} e!} P_G^{(e)}(-1),$$

where $P_G^{(e)}(-1)$ denotes the value at -1 of the e -th derivative of $P_G(T)$.

Proof. Apply the formula of Theorem 4 by replacing

- N , the length of the code, by $3e$;
- v , a lower bound for the range of f , by $-e$;
- n , the number of variables in f , by $2n$. □

Note that there are some obvious elements of weight 3 in K_G , namely

$$E_{i,j} + E_{i+n,j+n} + E_{i+2n,j+2n} \quad ((i,j) \in E).$$

More interestingly, every cycle of length r in G gives rise to at least 3 elements of weight r in K_G . Indeed, if (i_1, \dots, i_r) is a cycle, then

$$E_{i_1+kn, i_2+kn} + E_{i_2+kn, i_3+kn} + \dots + E_{i_r+kn, i_1+kn} \in K_G,$$

for $k = 0, 1, 2$.

One can go a little bit further. A somewhat technical computation shows that $P_G(T)$ can be decomposed as follows:

$$P_G(T) = (1 + T^3)^e P_{\bar{K}} \left(\frac{T}{1 - T + T^2} \right),$$

where \tilde{K} is the \mathbf{F}_4 -code defined by the exact sequence

$$0 \rightarrow \tilde{K} \rightarrow \mathbf{F}_4^e \rightarrow \mathbf{F}_4^n,$$

with weight enumerator $P_{\tilde{K}}(T)$. (The map on the right sends the basis vector corresponding to an edge, to the formal sum of its two endvertices.)

With the above formula, we find that $P_G^{(e)}(-1) = 3^e e! P_{\tilde{K}}\left(-\frac{1}{3}\right)$. Plugging this into Theorem 7, we obtain

$$\chi_G(4) = \frac{3^e}{4^{e-n}} P_{\tilde{K}}\left(-\frac{1}{3}\right).$$

In particular, G is 4-colorable if and only if $P_{\tilde{K}}\left(-\frac{1}{3}\right) \neq 0$. See also [E, Theorem 5.7], for a different formulation and proof of this formula.

REFERENCES

- [B] BARKER, R. H. Group synchronizing of binary digital systems. In *Communication Theory*, W. Jackson, Ed., London: Butterworth, 1953, 273-287.
- [E] ELIAHOU, S. An algebraic criterion for a graph to be four-colourable. *Aportaciones Matemáticas, Notas de Investigación 6* (1992), 3-27.
- [EKS] ELIAHOU, S., M. KERVAIRE and B. SAFFARI. On Golay polynomial pairs. *Adv. Appl. Math.* 12 (1991), 235-292.
- [G] GOLAY, M. J. E. Static multislit spectrometry and its application to the panoramic display of infrared spectra. *J. Opt. Soc. Amer.* 41 (1951), 468-472.
- [H] HADAMARD, J. Résolution d'une question relative aux déterminants. *Bull. Sci. Math. (2)* 17 (1893), 240-246.
- [M] MACWILLIAMS, F. J. A theorem on the distribution of weights in a systematic code. *Bell Syst. Tech. J.* 42 (1963), 79-94.
- [MS] MACWILLIAMS, F. J. and N. J. A. SLOANE. *The Theory of Error-Correcting Codes*. North-Holland, 1977.
- [SK] SAATY, T. L. and P. C. KAINEN. *The Four-Color Problem, Assaults and Conquest*. McGraw-Hill, New York, 1977.
- [SY] SEBERRY, J. and M. YAMADA. Hadamard matrices, Sequences, and Block Designs. In *Contemporary Design Theory: A Collection of Surveys*, J. H. Dinitz and D. R. Stinson, Eds., Wiley-Interscience, New York, 1992, 431-560.

(Reçu le 16 novembre 1993)

Shalom Eliahou

Section de Mathématiques
 Université de Genève
 C.P. 240
 1211 Genève 24, Switzerland
 e-mail: shalom@sc2a.unige.ch