

4. Very Low Dimensions

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **42 (1996)**

Heft 3-4: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **09.08.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

2. If M is an irreducible noncompact globally symmetric space G/K , with $G = \text{Isom}(M)$ and K a maximal compact subgroup, then one can say more about the bottom of the spectrum. If $\text{rk}(G) = \text{rk}(K)$ then $\text{Ker} \left(\Delta_{\frac{\dim(M)}{2}} \right)$ is infinite-dimensional and the spectrum of Δ is bounded away from zero otherwise. If $\text{rk}(G) > \text{rk}(K)$ then $\text{Ker}(\Delta) = 0$ and $0 \in \sigma(\Delta_p)$ if and only if

$$p \in \left[\frac{\dim(M)}{2} - \frac{\text{rk}(G) - \text{rk}(K)}{2}, \frac{\dim(M)}{2} + \frac{\text{rk}(G) - \text{rk}(K)}{2} \right]$$

[19, Section VIIB].

Finally, we state a result about uniformly contractible Riemannian manifolds.

DEFINITION 6 [15, p. 29]. *A metric space Z has finite asymptotic dimension if there is an integer n such that for any $r > 0$, there is a covering $Z = \bigcup_{i \in I} C_i$ of Z by subsets of uniformly bounded diameter so that no metric ball of radius r in Z intersects more than $n + 1$ elements of $\{C_i\}_{i \in I}$. The smallest such integer n is called the asymptotic dimension $\text{asdim}_+(Z)$ of Z .*

PROPOSITION 8 (Yu [33]). *If M is a uniformly contractible Riemannian manifold with finite asymptotic dimension then $0 \in \sigma(\Delta_p)$ for some p .*

The proof of Proposition 8 uses methods of coarse index theory [28].

4. VERY LOW DIMENSIONS

In this section we show that the answer to the zero-in-the-spectrum question is “yes” for one-dimensional simplicial complexes and two-dimensional Riemannian manifolds.

4.1 ONE DIMENSION

As a one-dimensional manifold is either S^1 or \mathbf{R} , zero is clearly in the spectrum.

A more interesting problem is to consider a connected one-dimensional simplicial complex K . Let V be the set of vertices of K and let E be the set of oriented edges of K . That is, an element e of E consists of an edge of K and an ordering (s_e, t_e) of ∂e . We let $-e$ denote the same edge with the

reverse ordering of ∂e . For $x \in V$, let m_x denote the number of unoriented edges of which x is a boundary. We assume that $m_x < \infty$ for all x . Put

$$C^0(K) = \{f : V \rightarrow \mathbf{C} \text{ such that } \sum_{x \in V} m_x |f(x)|^2 < \infty\},$$

$$(4.1) \quad C^1(K) = \{F : E \rightarrow \mathbf{C} \text{ such that } F(-e) = -F(e) \text{ and } \frac{1}{2} \sum_{e \in E} |F(e)|^2 < \infty\}.$$

Then $C^0(K)$ and $C^1(K)$ are Hilbert spaces. The weighting used to define $C^0(K)$ is natural in certain respects [8].

There is a bounded operator $d : C^0(K) \rightarrow C^1(K)$ given by $(df)(e) = f(t_e) - f(s_e)$. Define the Laplace-Beltrami operators by $\Delta_0 = d^*d$ and $\Delta_1 = dd^*$. An element of $\text{Ker}(\Delta_1)$ is an $F \in C^1(K)$ such that for each vertex x the total current flowing into x vanishes, i.e. $\sum_{e \in E: t_e=x} F(e) = 0$.

The next proposition is essentially due to Gromov [15, p. 236], who proved it in the case when $\{m_x\}_{x \in V}$ is bounded.

PROPOSITION 9. $0 \in \sigma(\Delta_0)$ or $0 \in \sigma(\Delta_1)$.

Proof. As the nonzero spectra of d^*d and dd^* are the same, for our purposes it suffices to consider $\sigma(\Delta_0)$ and $\text{Ker}(\Delta_1)$. We argue by contradiction. Suppose that $0 \notin \sigma(\Delta_0)$ and $\text{Ker}(\Delta_1) = 0$. First, K must be infinite, as otherwise $\text{Ker}(\Delta_0) \neq 0$. Second, K must be a tree, as if K had a loop then we could create a nonzero element of $\text{Ker}(\Delta_1)$ by letting a current of unit strength flow around the loop.

We now show that K has lots of branching. For $x, y \in V$, let $[x, y]$ be the geodesic arc from x to y and let (x, y) be its interior. Let $d(x, y)$ be the number of edges in $[x, y]$.

LEMMA 5. *There is a constant $L > 0$ such that if $d(x, y) > L$ then there is an infinite subtree of K which intersects (x, y) but does not contain x or y .*

Proof. If the lemma is not true then for all $N > 1$, there are vertices x and y such that $d(x, y) > N$ but there are no infinite subtrees as in the statement of the lemma. In other words, the connected component C of $K - \{x\} - \{y\}$ which contains (x, y) is finite. As K is a tree, x is only connected to the vertices in C by a single edge, and similarly for y (see Fig. 5). Define $f \in C^0(K)$ by

$$(4.2) \quad f(v) = \begin{cases} 1 & \text{if } v \in C, \\ 0 & \text{otherwise.} \end{cases}$$

Then

$$(4.3) \quad \frac{\langle df, df \rangle}{\langle f, f \rangle} \leq \frac{2}{2(d(x, y) - 1)} \leq \frac{1}{N}.$$

As N can be taken arbitrarily large, this contradicts the assumption that $0 \notin \sigma(\Delta_0)$. \square




FIGURE 5

It follows that K contains a subtree K' which is topologically equivalent to an infinite triadic tree, with the distances between branchings at most L (see Fig. 6). We can create a nonzero square-integrable harmonic 1-cochain F' on K' by letting a unit current flow through it, as in Fig. 6. Let $F \in C^1(K)$ be the extension of F' by zero to K . If x is a vertex of K' then the total current flowing into x is still zero, as no new current comes in along the edges of $K - K'$. Thus $\text{Ker}(\Delta_1) \neq 0$, which is a contradiction. \square




FIGURE 6

4.2 TWO DIMENSIONS

PROPOSITION 10 (Lott, Dodziuk). *The answer to the zero-in-the-spectrum question is “yes” if M is a two-dimensional manifold.*

Proof. The Hodge decomposition gives

$$(4.4) \quad \Lambda^0(M) = \text{Ker}(\Delta_0) \oplus \Lambda^0(M) / \text{Ker}(d),$$

$$(4.5) \quad \Lambda^1(M) = \text{Ker}(\Delta_1) \oplus \overline{d\Lambda^0(M)} \oplus \overline{*d\Lambda^0(M)},$$

$$(4.6) \quad \Lambda^2(M) = *\text{Ker}(\Delta_0) \oplus *(\Lambda^0(M) / \text{Ker}(d)).$$

Thus it is enough to look at

$$\text{Ker}(\Delta_0), \quad \text{Ker}(\Delta_1) \quad \text{and} \quad \sigma(\Delta_0 \text{ on } \Lambda^0(M) / \text{Ker}(d)).$$

We argue by contradiction. Assume that zero is not in the spectrum. By Proposition 4, $\text{Im}(\mathbb{H}_c^1(M) \rightarrow \mathbb{H}^1(M)) = 0$. Thus M must be planar, in the sense of either of the following two equivalent conditions:

1. Any simple closed curve in M separates it into two pieces.
2. M is diffeomorphic to the complement of a closed subset of S^2 .

As $\text{Ker}(\Delta_0) = 0$, M cannot be S^2 . By Proposition 5, the possible existence of nonzero square-integrable harmonic 1-forms on M only depends on the underlying Riemann surface coming from the Riemannian metric on M .

We recall some notions from Riemann surface theory [1]. A function $f \in C^\infty(M)$ is *superharmonic* if $\Delta_0 f > 0$. (This is a conformally-invariant statement.) The Riemann surface underlying M is *hyperbolic* if it has a positive superharmonic function and *parabolic* otherwise. If M is planar and hyperbolic then there is a nonconstant harmonic function $f \in C^\infty(M)$ such that $\int_M df \wedge *df < \infty$ [1, p. 208]. Then df would be a nonzero element of $\text{Ker}(\Delta_1)$. Thus M must be parabolic.

Put $\lambda_0 = \inf(\sigma(\Delta_0))$. Choose some λ such that $0 < \lambda < \lambda_0$. Then there is a positive $f \in C^\infty(M)$ (not square-integrable!) such that $\Delta_0 f = \lambda f$ [31, Theorem 2.1]. However, this contradicts the parabolicity of M . \square

We do not know of any result analogous to Proposition 10 for general two-dimensional simplicial complexes, say uniformly finite. See, however, Subsection 5.2.