

7. A RECTANGULAR LATTICE

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **42 (1996)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **09.08.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

To satisfy (3.11) we *redefine*

$$(6.18) \quad \pi_i(t) = a \mathcal{P}(t - c_i/2),$$

and set

$$(6.19) \quad G_0(z_1, z_2) = a^4 G(z_1/a, z_2/a),$$

so that

$$(6.20) \quad \overline{G_0(z_1, z_2)} = G_0(\bar{z}_2, \bar{z}_1).$$

In summary we have

PROPOSITION 6.1. *Let $\Lambda = \mathbf{C}/\Lambda$ have the holomorphic involutions (6.1) intertwined by the anti-holomorphic involution (5.6). Then (Γ, ρ, τ_i) is realized by the map (6.5), (6.18) onto the quartic curve $G_0(z_1, z_2) = 0$ given by (6.13), (6.14), (6.19). If the fixed-point set of ρ is non-empty, then this is the complexification of the real curve $G_0(z, \bar{z}) = 0$.*

7. A RECTANGULAR LATTICE

We consider the special case of Λ, ρ, τ_i as given in (5.19), (5.6), (6.1), with

$$(7.1) \quad a = +1, b = 0, \bar{c}_2 = c_1 = c'_1 + ic''_1, c = ic''_1.$$

From (6.16), (6.15) it follows that g_2, g_3, β are real, and β' is purely imaginary. Thus, the coefficients $\beta_1, \beta_2, \beta_3$ of $G(z_1, z_2)$ are real. With $t = t' + it''$, we have

$$(7.2) \quad FP(\rho) = \{t'' = 0\} \cup \{t'' = \omega''/2\},$$

$$(7.3) \quad \tau_1\{t'' = 0\} = \{t'' = c''_1\}, \tau_1\{t'' = \omega''/2\} = \{t'' = c''_1 + \omega''/2\}.$$

Let us assume that $0 < c''_1 < \omega''/2$. Then the torus Λ is divided into four annuli

$$A_1 = \{0 < t'' < c''_1\}, A_2 = \{c''_1 < t'' < \omega''/2\},$$

$$A_3 = \{\omega''/2 < t'' < c''_1 + \omega''/2\}, A_4 = \{c''_1 + \omega''/2 < t'' < \omega''\}.$$

The fixed points of τ_1 are, by (5.22),

$$(7.4) \quad c_1/2, (c_1 + 1)/2 \in A_1,$$

$$(7.5) \quad (c_1 + i\omega'')/2, (c_1 + 1 + i\omega'')/2 \in A_3.$$

For the map $z = \pi_1(t)$ (6.3), we get

$$(7.6) \quad D_j = \pi_1(A_j), C_j = \partial D_j, 1 \leq j \leq 4.$$

Then the z -plane is the disjoint union of $D_1, D_2 = D_4, D_3, C_1$, and C_3 . π_1 maps each of A_2 and A_4 biholomorphically onto D_2 , which is topologically an annulus with boundary $C_2 = C_1 - C_3$. π_1 also gives twofold branched coverings of A_i onto D_i , $i = 1, 3$, branched at (7.4), (7.5). In particular, D_1 is unbounded, containing $\pi_1(c_1/2) = \infty$ in its interior, and C_1 and C_3 are symmetric with respect to the real axis. $\pi_1(t)$ is real on the two horizontal lines through (7.4) and (7.5). It is also real on the two vertical lines $\{t' = c'_1/2\}$, $\{t' = (c'_1 + 1)/2\}$, which intersect ∂A_1 in the points

$$(7.7) \quad a_1 = \frac{1}{2} c'_1, a_2 = \frac{1}{2} c'_1 + ic''_1,$$

and

$$(7.8) \quad b_1 + 1 = \frac{1}{2} (c'_1 + 1), b_2 = \frac{1}{2} (c'_1 + 1) + ic''_1.$$

$\pi_1(a_1) = \pi_1(a_2) \in C_1 \cap \mathbf{R}$ and $\pi_1(b_1 + 1) = \pi_1(b_2) \in C_1 \cap \mathbf{R}$ are the "vertices" of C_1 .

The "annular" domain D_2 has the same conformal type as A_2 , which is determined by $\frac{1}{2} \omega'' - c''_1$. This depends on both Λ and τ_1 .

Finally we consider the Riemann map, $\zeta = f(z)$, of D_1 onto the right half plane H , which takes $\pi_1(a_1)$ to zero and $\pi_1(b_1 + 1)$ to ∞ . We extend $f \circ \pi_1(t)$ to the entire t -plane by reflection in the lines $\{t'' = 0\}$ and $\{t'' = c''_1\}$. This gives a doubly periodic meromorphic function $\hat{\phi}$ with period module

$$(7.9) \quad \hat{\Lambda} = \{n_1 \cdot 1 + n_2 \cdot 2c''_1 i \mid n_1, n_2 \in \mathbf{Z}\}.$$

$\hat{\phi}$ has the representation in terms of the sigma function \hat{S} relative to $\hat{\Lambda}$,

$$(7.10) \quad \hat{\phi} = \frac{\hat{S}(t - a_1) \hat{S}(t - a_2)}{\hat{S}(t - b_1) \hat{S}(t - b_2)}.$$

The invariance $\hat{\phi} \circ \tau_1 = \hat{\phi}$ follows as in section 4, since $\tau_1(a_1) = a_2$, $\tau_1(b_1) = b_2$. Since $z = \mathcal{P}(t - c_1/2)$, we have

THEOREM 7.1. *Let Λ be the lattice with periods (5.19), and let $D_1 \subset \mathbf{P}_1$ be the simply connected domain above. The Riemann map, $\zeta = f(z)$, of D_1 onto the right half ζ -plane H is given by*

$$(7.11) \quad \zeta = \hat{\phi}(\mathcal{P}^{-1}(z) + c_1/2),$$

where $\hat{\phi}$ is the sigma quotient (7.10) relative to the lattice (7.9), and $\mathcal{P}^{-1}(z)$ is the elliptic integral of the first kind, in Weierstrass normal form, relative to Λ .

REMARK. We have seen that double valued reflection places a severe restriction on a real algebraic curve in the complex plane. In fact our results should provide the basis for a complete and explicit classification. We have also seen how double valued reflection may be used to explicitly determine Riemann maps. Apparently, all known such examples can be so explained. The result in the above theorem seems to be new. It would be interesting to work out more examples in the genus one case.

REFERENCES

- [1] CARATHEODORY, C. *Conformal Representation*. Cambridge University Press (1952).
- [2] COURANT, R. and D. HILBERT. *Methods of Mathematical Physics*, vol. I. Interscience, New York (1953).
- [3] DU VAL, P. *Elliptic Functions and Elliptic Curves*. Cambridge University Press (1973).
- [4] FARKAS, H. and I. KRA. *Riemann Surfaces*. Springer-Verlag (1980).
- [5] FRICKE, R. *Elliptische Functionen*. B.G. Teubner, Leipzig und Berlin (1916).
- [6] HURWITZ, A. and R. COURANT. *Functionentheorie*. Springer-Verlag (1964).
- [7] KELLOGG, O.D. *Foundations of Potential Theory*. Dover Publications, New York (1954).
- [8] MOSER, J.K. and S.M. WEBSTER. Normal forms for real surfaces in \mathbb{C}^2 near complex tangents and hyperbolic surface transformations. *Acta Math.* 150 (1983), 255-296.
- [9] NEHARI, Z. *Conformal Mapping*. Dover Publications, New York (1975).
- [10] SCHWARZ, H.A. *Gesammelte Mathematische Abhandlungen*, vol. II. Berlin (1890).

(Reçu le 15 mai 1995)

S.M. Webster

University of Chicago
 Department of Mathematics
 5734 University Ave.
 Chicago, IL 60637
 U.S.A.