

4. Obtaining Elements of N

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **43 (1997)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **30.06.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

For $l \in L, v \in \mathbf{Q}^d$ we have $vF(ln^{-1})^{tr} = va^{-1}nFl^{tr}$ and hence $(L')^\# = L^\#an^{-1}$.

Since $L' \in \pi(L)$ one has $L' = L^\# \cap a^{-1}L$. Using this one obtains

$$Lan^{-2} = L'an^{-1} = L^\#an^{-1} \cap Ln^{-1} = (L')^\# \cap L' = L,$$

since $(L')^\#/L$ is the orthogonal complement of L'/L in $L^\#/L$ with respect to the induced quadratic form with values in \mathbf{Q}/\mathbf{Z} . So $a^{-1}n^2 \in G$.

Finally we check that $n \in N$. Let $g \in G$, then $n^{-1}gn$ is in $G = \text{Aut}(F)$ since $Ln^{-1}gn = L'gn = L'n = L$ and

$$n^{-1}gnFn^{tr}g^{tr}n^{-tr} = n^{-1}agFg^{tr}n^{-tr} = F. \quad \square$$

4. OBTAINING ELEMENTS OF N

Now we give examples as to how one may construct elements n of the normaliser N . To obtain similarities we are interested in $n \in N$ of determinant $\pm p^{d/2}$ for some (squarefree) natural number p such that $p^{-1}n^2 \in G$. The first method is an application of the normaliser principle to the situation (iii) described in Section 2:

PROPOSITION 4. *Let $U \trianglelefteq G$ be a normal subgroup of G and assume that the commuting algebra $K := C_{M_d(\mathbf{Q})}(U)$ is isomorphic to a number field. If $c \in K$ satisfies $c^2 = p \in \mathbf{Q}^*I_d$, then c lies in N .*

Proof. Since G normalises U , it acts by conjugation (and hence as Galois automorphisms) on the abelian number field K . Now let $c \in K$, with $c^2 =: p \in \mathbf{Q}^*I_d$ and $g \in G$. Then g stabilises the subfield $\mathbf{Q}[c]$ and hence $g^{-1}cg = \pm c$, which is equivalent to $c^{-1}gc = \pm g \in G$. Therefore $c \in N$, since we assumed that $-I_d \in G$. \square

The following construction described in [PIN 95] Proposition (II.4) also allows us to find elements of N .

For $i = 1, 2$ let $G_i \leq GL_{d_i}(\mathbf{Q})$ be finite rational irreducible matrix groups with commuting algebras $A_i \subseteq M_{d_i}(\mathbf{Q})$. Also let Q be a maximal common subalgebra of dimension z of A_1 and A_2 . Let $d := \frac{d_1d_2}{z}$ and view the G_i as subgroups of $G_1 \otimes_Q G_2 \leq GL_d(\mathbf{Q})$. If there exist elements $a_i \in N_{GL_d(\mathbf{Q})}(G_i)$

centralising G_j and a_j ($1 \leq i \neq j \leq 2$) and a squarefree natural number $p \neq 0$ such that $p^{-1}a_i^2 \in G_i$, the group

$$G := \langle G_1 \otimes_{\mathbf{Q}} G_2, p^{-1}a_1a_2 \rangle,$$

generated by the elements of $G_1 \otimes_{\mathbf{Q}} G_2$ and $p^{-1}a_1a_2$, is a finite subgroup of $GL_d(\mathbf{Q})$ containing $G_1 \otimes_{\mathbf{Q}} G_2$ as a subgroup of index 2.

For $d \leq 31$ and $p > 1$ we only need the case where a_2 is an element of the enveloping algebra of G_2 . Then G is denoted by $G_1 \otimes_{\mathbf{Q}}^{2(p)} G_2$ (or $G_1 \boxtimes_{\mathbf{Q}}^{2(p)} G_2$) according to whether a_1 is (or is not) a rational linear combination of elements of G_1 .

Using this notation one immediately has the following proposition.

PROPOSITION 5. *For $i = 1, 2$ the matrix a_i is an element of determinant $\pm p^{d/2}$ in the normaliser N of G .*

A common feature of the situations in Propositions 4 and 5 is that we extend the natural representation of G to a projective representation which is realisable as a linear representation over a quadratic extension of \mathbf{Q} .

PROPOSITION 6. *Let $G \trianglelefteq E$ be a supergroup containing G of index 2. Assume that $C_{M_d(\mathbf{Q})}(G) \cong \mathbf{Q}$ and that the natural character of G extends to E with character field $\mathbf{Q}[\sqrt{p}]$, where $p \in \mathbf{Z}$ is not a square. Then there exists $n \in N$ of determinant $\pm p^{d/2}$ with $p^{-1}n^2 \in G$.*

Proof. By Clifford theory one may extend the natural representation Δ of G to a representation $\delta_1 \otimes \delta_2 : E \rightarrow (\mathbf{Q}[\sqrt{p}] \otimes M_d(\mathbf{Q}))^*$, where δ_1 and δ_2 are projective representations $\delta_1(G) = \{1\}$ and $(\delta_2)|_G = \Delta$. Let $e \in E \setminus G$. Then

$$(\delta_1(e) \otimes \delta_2(e))^2 = \delta_1(e)^2 \otimes \delta_2(e)^2 = 1 \otimes \Delta(e^2),$$

since $e^2 \in G$. Therefore $\delta_1(e)^2 \in \mathbf{Q}$. Replacing $\delta_1(e)$ by a suitable rational multiple (and multiplying $\delta_2(e)$ by the inverse) one may assume that $\delta_1(e)^2 = p^{-1}$. Then $n := \delta_2(e)$ is an element of the normaliser N with the desired properties. \square

	$\text{Aut}(L)$	$\det(L)$	$\min(L)$	$ L_{\min} $	lattice sparse
4	$F_4 \tilde{\otimes} F_4 = [2_+^{1+8} \cdot O_8^+(2)]_{16}$	2^8	4	4320	+
6	$[(SL_2(9) \otimes_{\infty,3}^{2(3)} SL_2(9)) \cdot 2]_{16}$	3^8	4	720	+
9	$[(Sp_4(3) \circ C_3) \boxtimes_{\sqrt{-3}}^2 SL_2(3)]_{16}$	$2^8 \cdot 3^8$	6	960	+
14	$[2 \cdot \text{Alt}_{10}]_{16}$	5^8	6	2400	+
16	$[SL_2(5) \otimes_{\infty,2}^{2(2)} 2^{1+4'} \cdot \text{Alt}_5]_{16}$	$2^8 \cdot 5^8$	8	1200	+
19	$[SL_2(5) \otimes_{\infty,3}^{2(3)} SL_2(9)]_{16}$	$3^8 \cdot 5^8$	10	1440	+
21	$[SL_2(5) \otimes_{\infty,3}^{2(3)} (SL_2(3) \square^2 C_3)]_{16}$	$2^8 \cdot 3^8 \cdot 5^8$	12	480	+
25	$[2 \cdot \text{Alt}_7 \otimes_{\sqrt{-7}}^{2(3)} \tilde{S}_3]_{16}$	$3^8 \cdot 7^8$	12	1680	+
26	$[SL_2(7) \otimes_{\sqrt{-7}}^{2(3)} \tilde{S}_3]_{16}$	$3^8 \cdot 7^8$	10	336	$p \neq 2$
3	$[2 \cdot \text{Co}_1]_{24}$	1^{24}	4	196560	+
6	$[6 \cdot U_4(3) \cdot 2 \boxtimes_{\sqrt{-3}}^2 SL_2(3)]_{24}$	2^{12}	4	3024	$p \neq 3$
16	$[6 \cdot L_3(4) \cdot 2 \otimes^{2(2)} D_8]_{24}$	$2^{12} \cdot 3^{12}$	8	3024 + 7560	+
17	$[(SL_2(3) \circ C_4) \cdot 2 \boxtimes_{\sqrt{-1}}^{2(3)} U_3(3)]_{24}$	$2^{12} \cdot 3^{12}$	8	4536 + 6048	+
18	A_{24}	1^{24}	2	600	$p \neq 5$
22	$[2 \cdot J_2 \square^2 SL_2(5)]_{24}$	5^{12}	8	37800	+
35	$[L_2(7) \otimes^{2(2)} F_4]_{24}$	7^{12}	8	1008 + 3024	$p \neq 2$
40	$[SL_2(13) \square^{2(2)} SL_2(3)]_{24}$	13^{12}	12	$2 \cdot 2184 + 8736$	$p \neq 2$
42	$[6 \cdot \text{Alt}_7 : 2]_{24}$	2^{12}	4	3024	+
43	$[3 \cdot M_{10} \otimes_{\sqrt{-3}}^{2(2)} SL_2(3)]_{24}$	$2^{12} \cdot 5^{12}$	8	1080	$p \neq 3$
44	$[\text{Alt}_5 \otimes_{\sqrt{5}}^2 (C_3 \otimes^{2(2)} D_8)]_{24}$	$2^{12} \cdot 3^{12} \cdot 5^{12}$	16	$360 + 2 \cdot 720$	$p \neq 2$
45	$[3 \cdot M_{10} \otimes^{2(2)} D_8]_{24}$	$2^{12} \cdot 3^{12} \cdot 5^{12}$	16	$1080 + 1080$	+
64	$[SL_2(11) \otimes_{\sqrt{-11}}^{2(2)} SL_2(3)]_{24}$	$2^{12} \cdot 11^{12}$	12	1320	$p \neq 2$