

## 4. $SL(n,R)$ -ACTIONS ON $\mathbb{R}^n$ FOR $n \geq 3$

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **43 (1997)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **30.06.2024**

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

### **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

4.  $SL(n, \mathbf{R})$ -ACTIONS ON  $\mathbf{R}^n$  FOR  $n \geq 3$ 

Let  $n \geq 3$ . We first give examples of  $C^0$ -actions of  $SL(n, \mathbf{R})$  on  $\mathbf{R}^n$ . Consider the canonical projective action of  $SL(n, \mathbf{R})$  on  $S^{n-1}$ . Let  $\Delta^+$  be the radial half-line through the first basis element  $e_1$  and let  $H$  denote the subgroup of  $SL(n, \mathbf{R})$  that fixes  $\Delta^+$ . So  $SL(n, \mathbf{R})/H \cong S^{n-1}$ . Consider the homomorphism

$$\psi: (A_{ij}) \in H \mapsto \ln A_{11} \in \mathbf{R}.$$

Notice that one obtains a linear action of  $H$  on  $\mathbf{R}_*^+ = (0, \infty)$  by setting  $h(x) = e^{\psi(h)}x$ , for all  $h \in H$ ,  $x \in \mathbf{R}_*^+$ . Obviously this is conjugate to the  $H$ -action on  $\Delta^+$ . It follows from Lemma 3.7 that the action of  $SL(n, \mathbf{R})$  obtained by suspension of this action of  $H$  on  $\mathbf{R}_*^+$  is the canonical linear action of  $SL(n, \mathbf{R})$  on  $\mathbf{R}^n \setminus \{0\}$ . In fact, the map

$$\psi: [g \cdot x] \in (SL(n, \mathbf{R}) \times \mathbf{R}_*^+)/H \mapsto g(xe_1) \in \mathbf{R}^n \setminus \{0\}$$

is an isomorphism. We now deform the action of  $H$ . Choose a topological flow  $(\phi^t)_{t \in \mathbf{R}}$  on  $\mathbf{R}^+ = [0, \infty)$ , fixing 0. This defines an action of  $H$  on  $\mathbf{R}_*^+$  by setting  $h(x) = \phi^{\psi(h)}(x)$ , for all  $h \in H$ ,  $x \in \mathbf{R}_*^+$ . Now suspend this action of  $H$  and let  $\Phi$  denote the resulting action of  $SL(n, \mathbf{R})$  on the space  $M = (SL(n, \mathbf{R}) \times \mathbf{R}_*^+)/H$ . The space  $M$  fibres over  $S^{n-1}$ , with fibre  $\mathbf{R}_*^+$ , and the structure group is orientation preserving. So topologically,  $M$  is  $\mathbf{R}_*^+ \times S^{n-1}$ . Thus, identifying  $S^{n-1} \times \{0\}$  to a point, we obtain an  $SL(n, \mathbf{R})$ -action on  $\mathbf{R}^n$ . The fixed points of the flow  $\phi$  correspond to orbits in  $\mathbf{R}^n$  which are spheres of dimension  $n-1$ . In general, an  $n$ -dimensional orbit is either all of  $\mathbf{R}^n \setminus \{0\}$ , as in the linear case, or it is a spherical shell, bounded by  $S^{n-1}$  orbits, or a punctured ball bounded by an  $S^{n-1}$  orbit, or the exterior of an  $S^{n-1}$  orbit. In all cases, the  $n$ -dimensional orbits are conjugate to the canonical linear one on  $\mathbf{R}^n \setminus \{0\}$ , by Theorem 3.5(c).

**THEOREM 4.1.** *For all  $n \geq 3$ , every non-trivial  $C^0$ -action of  $SL(n, \mathbf{R})$  on  $(\mathbf{R}^n, 0)$  is conjugate to one of the above actions  $\Phi$ .*

*Proof.* Suppose that we have a non-trivial  $C^0$ -action of  $SL(n, \mathbf{R})$  on  $(\mathbf{R}^n, 0)$ . First use Proposition 3.8 to linearize the  $SO(n)$ -action. Then by Lemma 3.9, the  $SL(n, \mathbf{R})$ -action preserves the radial lines. Hence the radial projection  $\mathbf{R}^n \setminus \{0\} \rightarrow S^{n-1}$  is equivariant, where the action of  $SL(n, \mathbf{R})$  on  $S^{n-1}$  is the canonical projective one. Let  $H$  be the stabilizer of the radial half-line  $\Delta^+$  through  $e_1$ , as above. So the action of  $SL(n, \mathbf{R})$  on  $\mathbf{R}^n \setminus \{0\}$  is induced by some action of  $H$  on  $\mathbf{R}$ . Notice that this action is trivial when restricted to

$SO(n-1)$ . It remains to consider all actions of  $H$  on  $\mathbf{R}$  which are trivial on  $SO(n-1)$ . Again, by Lie [23, *ibid.*], these are given by homomorphisms from  $H$  to  $\mathbf{R}$ ,  $\mathbf{Aff}$ , or (some cover of)  $PSL(2, \mathbf{R})$ . We have the homomorphism  $\psi: (A_{ij}) \in H \mapsto \ln A_{11} \in \mathbf{R}$ . Note that  $\ker \psi = SL(n-1, \mathbf{R}) \ltimes \mathbf{R}^{n-1}$ . But it is easy to see that there are no non-trivial homomorphisms of  $\ker \psi$  to  $\mathbf{R}$  or  $\mathbf{Aff}$ . There are no non-trivial homomorphisms of  $\ker \psi$  to  $SL(2, \mathbf{R})$ , except in the case  $n=3$ , and in this case there are no such homomorphisms which are trivial on  $SO(n-1)$ . So the only possibility left is that  $H$  acts on  $\mathbf{R}$  by some flow. Finally, we put back the origin, as in the proof of Proposition 3.8. This completes the proof of the theorem.  $\square$

We now prove Theorem 1.1 for  $n \geq 3$ .

**THEOREM 4.2.** *For all  $n \geq 3$  and  $k = 1, \dots, \infty$ , every  $C^k$ -action of  $SL(n, \mathbf{R})$  on  $(\mathbf{R}^n, 0)$  is  $C^k$ -linearizable.*

*Proof.* Let  $n \geq 3$  and  $k = 1, \dots, \infty$  and suppose that we have a non-trivial  $C^k$ -action of  $SL(n, \mathbf{R})$  on  $(\mathbf{R}^n, 0)$ . By Remark 3.4, we may assume that the differential of the action at the origin is either the identity or the map  $g \mapsto (g^{-1})^t$ . We will assume that it is the identity; the other possibility can be handled using the same argument.

Linearizing the  $SO(n)$ -action, using the Bochner-Cartan theorem, one may assume that the  $SO(n)$ -action is the canonical one. Then by Lemma 3.9, the  $SL(n, \mathbf{R})$ -action preserves the radial lines. Let  $\Delta$  denote the radial line through the first of the canonical basis elements,  $e_1$ . Consider  $H = \text{Stab}_{SL(n, \mathbf{R})}(\Delta)$ , as before. So, as we saw in the proof of Theorem 4.1,  $H$  defines a  $C^k$ -flow on  $\Delta$ . This flow is hyperbolic, by the first paragraph. Hence by Theorem 2.5, this flow is linearizable by some local  $C^k$ -diffeomorphism  $f$  of  $\Delta(\cong \mathbf{R})$ . So, after conjugacy, we may assume that  $H$  acts linearly on  $\Delta$ . Now define the local  $C^k$ -diffeomorphism  $F$  of  $\mathbf{R}^n$  by the formula:

$$(2) \quad F(x) = \begin{cases} \frac{f(\|x\|)}{\|x\|} x, & x \neq 0 \\ 0, & x = 0. \end{cases}$$

To see that  $F$  is of class  $C^k$ , the key point is to verify that  $f$  is a  $C^k$  odd function on  $\mathbf{R}$ . This follows easily from the fact that the flow on  $\Delta$  commutes with  $\text{Stab}_{SO(n, \mathbf{R})}(\Delta)$ , and the  $SO(n)$ -action is linear.

Now notice that  $F$  agrees with  $f$  on  $\Delta^+ = \{te_1 \in \Delta : t \geq 0\}$ , and as  $F$  commutes with the  $SO(n)$ -action, the  $SO(n)$ -action is unchanged by conjugation by  $F$ . In particular, the  $SO(n)$ -action still commutes with dilations.

It follows that after conjugation by  $F$ , the  $SL(n, \mathbf{R})$ -action commutes with dilations. Indeed, consider the conjugated  $SL(n, \mathbf{R})$ -action. If  $f \in SL(n, \mathbf{R})$ ,  $x \in \mathbf{R}^n$  and  $\lambda > 0$ , then choose  $a, b \in SO(n)$  such that  $ax \in \Delta^+$  and  $bf(\lambda x) \in \Delta^+$ . Provided  $x$  is sufficiently close to 0,  $ax$  and  $bf(\lambda x)$  will lie in the domain of  $f$ . Then  $bfa^{-1} \in H$  and so

$$\begin{aligned} f(\lambda x) &= b^{-1}bfa^{-1}a(\lambda x) = b^{-1}(bfg^{-1})\lambda a(x) \\ &= b^{-1}\lambda(bfa^{-1})a(x) = \lambda b^{-1}(bfa^{-1})a(x) \\ &= \lambda f(x). \end{aligned}$$

The proof of the theorem is then completed by the following well known result (cf. [17, Lemma 2.1.4]).  $\square$

LEMMA 4.3. *Every  $C^1$  map commuting with dilations is linear.*

*Proof.* Suppose that  $f$  is a  $C^1$ -diffeomorphism of  $\mathbf{R}^n$  which commutes with dilations. By comparing the differential of  $\lambda \cdot f$  and  $f \circ \lambda$  at  $x$  we have  $\lambda df|_x = \lambda df|_{\lambda x}$ , for each  $\lambda > 0$  and every  $x \in \mathbf{R}^n$ . Hence  $df|_x = df|_{\lambda x}$  and so  $df$  is constant on the radial lines. Thus  $df|_x = df|_0$  for all  $x$  and so  $f$  is linear.  $\square$

## 5. THE ADJOINT REPRESENTATION OF $SL(2, \mathbf{R})$

Let us recall some facts concerning the linear representations of  $SL(2, \mathbf{R})$ . Let  $P_l(\mathbf{R}^2)$  denote the space of real valued homogeneous polynomials, of two variables, of degree  $l$ . As a vector space,  $P_l(\mathbf{R}^2) \cong \mathbf{R}^{l+1}$ , and the action of  $SL(2, \mathbf{R})$  on  $\mathbf{R}^2$  defines a linear action on  $P_l(\mathbf{R}^2)$ : up to isomorphism, this is the (unique) irreducible representation of  $SL(2, \mathbf{R})$  in dimension  $l + 1$ . In dimension 3, there is another useful realization of the polynomial representation, called the *adjoint representation*. Notice that the group  $SL(2, \mathbf{R})$  acts by the adjoint representation on its Lie algebra  $\mathfrak{sl}(2, \mathbf{R})$ . Of course,  $\mathfrak{sl}(2, \mathbf{R})$  is the space of  $2 \times 2$  real traceless matrices; so as a vector space,  $\mathfrak{sl}(2, \mathbf{R}) \cong \mathbf{R}^3$ . The adjoint representation  $Ad: SL(2, \mathbf{R}) \rightarrow GL(3, \mathbf{R})$ , defined by

$$Ad(g): h \mapsto ghg^{-1}, \quad \forall g \in SL(2, \mathbf{R}), \quad h \in \mathfrak{sl}(2, \mathbf{R}),$$

is an irreducible linear representation. In fact, an explicit equivariant isomorphism  $\psi: \mathfrak{sl}(2, \mathbf{R}) \rightarrow P_2(\mathbf{R}^2)$  is obtained by taking  $\psi(h)$ , as a function of variables  $x$  and  $y$ , to be the area of the parallelogram spanned by  $(x, y)$  and  $h(x, y)$ . That is,