

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 44 (1998)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: AN ASYMPTOTIC FREIHEITSSATZ FOR FINITELY GENERATED GROUPS
Autor: Cherix, Pierre-Alain / SCHAEFFER, Gilles
Kapitel: 1. Introduction
DOI: <https://doi.org/10.5169/seals-63893>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 19.01.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

AN ASYMPTOTIC FREIHEITSSATZ
FOR FINITELY GENERATED GROUPS

by Pierre-Alain CHERIX*) and Gilles SCHAEFFER

ABSTRACT. Given two fixed integers $k \geq 2$ and $l \geq 3$, let $\Gamma = \langle X \mid R \rangle$ be a presentation of the group Γ with $k = \#X$ generators and $l = \#R$ relations. We show that the following property of presentations of groups is generic in the sense of Gromov: for any $y \in X$, the subgroup of Γ generated by $X - \{y\}$ is free of rank $k - 1$. This gives some generic estimates for the spectral radius of the adjacency operator in the Cayley graph of Γ relative to the generating system $S = X \cup X^{-1}$.

1. INTRODUCTION

The existence of free subgroups in some finitely generated group Γ gives some information about the structure of Γ . For example, it implies that Γ is non-amenable, and in particular that Γ has exponential growth. There are several results which ensure that various groups do have non-abelian free subgroups. For example:

THEOREM (Tits's alternative [15]). *Let Γ be a finitely generated linear group. Then either Γ is almost solvable or Γ contains a free subgroup on two generators.*

*) Le premier auteur est supporté par une bourse "Jeune chercheur" du Fonds National Suisse de la Recherche Scientifique et a effectué ce travail à l'université de New South Wales.

THEOREM (Magnus's Freiheitssatz [12]). *Let $\Gamma = \langle X|r \rangle$ be a one relator group, $x_0 \in X$ be a generator of Γ that appears in the relation r and r be a cyclically reduced word in the free group \mathbf{F}_X generated by X ; then $X - \{x_0\}$ freely generates a free group in Γ .*

Our purpose in this work is to measure in some sense how frequent it is for a presentation $\Gamma = \langle X|R \rangle$ to be such that a proper subset of X is free in Γ . We prove the following result:

THEOREM 1.1. *Let $\Gamma = \langle X|R \rangle$ be a finite presentation with k generators, l relations and any fixed x_0 in X . Then the fact that $X - \{x_0\}$ freely generates a free group in Γ is generic in the sense of Gromov.*

The key idea is contained in proposition 4.1. Roughly speaking, if you choose at random l long relations and if the presentation satisfies a Dehn algorithm, then every generator x_0 will appear in every sufficiently long subword of every relation and hence it will appear in every product of conjugates of relations. So $X - \{x_0\}$ generates a free group in Γ .

In [6], the first author has shown that “ X generates a free semi-group” is generic and that this implies bounds on the spectrum of the adjacency operator associated to the oriented Cayley graph of Γ relative to X . In section 5 below, we consider the adjacency operator h_S of the Cayley graph of Γ relative to $S = X \cup X^{-1}$, and we prove similarly estimates on the norm of h_S .

After finishing this paper we discovered that a result similar to Theorem 1.1 has been proved, using different methods, by G. Arzhantseva and A. Ol'shanskii in [1]. They employed a slightly different definition of the genericity and they proved that the small cancellation condition $C'(\lambda)$ is generic with respect to this new definition.

We thank A. Valette for his useful remarks and for the proofreading of this paper.