

## 8.2 The case of multiple roots

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **44 (1998)**

Heft 3-4: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **09.08.2024**

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

### **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Clearly  $v > 0$ ; moreover

$$u = \frac{\frac{p_{h-1}}{q_{h-1}} - \frac{q_h}{q_{h-1}} \frac{p_h}{q_h}}{1 - \frac{q_h}{q_{h-1}}} = \frac{p_{h-1} - p_h}{q_{h-1} - q_h} > 0.$$

It follows that this circle is entirely contained in the right half plane. A root  $x_j$  different from  $x_0$  (see Fig. 3) lies outside the circle (8.3) if  $h$  is large enough to have  $\Delta > |b - a|$ . It follows that  $\operatorname{Re} \mathcal{F}(x_j) < 0$ , and hence it is external to the circle (8.6) corresponding to the value  $h + 1$ . Hence the condition

$$F_{h-1} F_{h-2} \Delta > 1$$

ensures that the polynomial  $f_{h+1}$  is reduced.

## 8.2 THE CASE OF MULTIPLE ROOTS

Obreschkoff's Lemma 8.1 yields the following

**COROLLARY 8.3.** *Let  $f(x) = (x - x_1)(x - x_2) \cdots (x - x_r)$ , where  $x_i \in \mathbf{R}^+$ . Then*

$$f_1(x) = (x^2 + 2\rho x \cos \varphi + \rho^2) f(x), \quad \rho > 0, \quad |\varphi| < \frac{\pi}{r+2}$$

*has exactly  $r$  variations. More generally, a polynomial having  $r$  positive real roots and all its other roots in the sector*

$$S = \left\{ x \mid x = -\rho(\cos \varphi + i \sin \varphi), \quad \rho > 0, \quad |\varphi| < \frac{\pi}{r+2} \right\}$$

*has exactly  $r$  variations.*

This allows us to extend Vincent's theorem to the case of multiple roots. Suppose the polynomial  $f(x)$  has multiple roots, and let  $\Delta$  be their least distance. If  $h$  is sufficiently large to verify

$$F_h F_{h-1} \Delta > 1,$$

at most one root  $x_0$  lies in  $(a, b)$ , but since this root may have multiplicity  $r$ ,  $f_h$  has 0 or at least  $r$  variations. It will have exactly  $r$  variations if we can ensure that  $x_0 \in (a, b)$  and that the other transformed roots lie in the sector

$$S = \{y \mid \operatorname{Re} y < 0, \quad |\operatorname{Im} y| < |\tan \varphi| \cdot |\operatorname{Re} x|\},$$

where  $\varphi = \frac{\pi}{r+2}$ . Let  $s = \tan \frac{\pi}{r+2}$  and let us make the appropriate substitutions into (8.4). We have proved

THEOREM 8.4. *Let  $f(x)$  be a real polynomial of degree  $n$  whose roots are of multiplicity smaller than  $r$ . Let  $\gamma = [c_0, c_1, c_2, \dots]$  and, maintaining the previous notation, consider the polynomials*

$$f_{h+1} = (q_{h-1} + q_h x) f\left(\frac{p_{h-1} + p_h x}{q_{h-1} + q_h x}\right).$$

Let  $s = \tan \frac{\pi}{r+2}$ . If  $h$  satisfies

$$F_h F_{h-1} \Delta > \sqrt{1 + \frac{1}{s^2}} = \frac{1}{\sin \frac{\pi}{r+2}},$$

then the number of variations of  $f_{h+1}$  equals the multiplicity of the root in  $\left(\frac{p_{h-1}}{q_{h-1}}, \frac{p_h}{q_h}\right)$ .

REMARK 9. Obviously, letting  $s = \tan \frac{\pi}{n+2}$ , we can implement an algorithm to isolate the roots, without being forced to reduce the polynomial  $f(x)$  to one with simple roots.

REMARK 10. We conclude our paper by showing that our estimate of the size of  $h$  is asymptotically better than Chen's. Suppose we consider a polynomial whose roots are of multiplicity  $\leq r$  (which necessarily has degree  $n \geq r$ ). We have proved that the isolation of a root can be carried out in  $p$  steps, where  $p$  verifies

$$(8.7) \quad F_p F_{p-1} \Delta > \sqrt{1 + \frac{1}{\tan^2 \frac{\pi}{r+2}}} = \frac{1}{\sin \frac{\pi}{r+2}}.$$

We want to compare this integer with that needed by Chen's theorem, that is the smallest integer  $m = h + k$ , where  $h$  and  $k$  satisfy

$$(8.8) \quad F_h F_{h-1} \Delta > 1 \quad \text{and} \quad k > \frac{1}{2} \log_{\phi} r.$$

We know that

$$\sqrt{5} F_k \approx \phi^{k+1},$$

hence (8.7) becomes

$$\frac{\Delta}{5} \phi^{2p+1} > \frac{1}{\sin \frac{\pi}{r+2}}.$$

On the other hand, by (8.8) we have

$$(8.9) \quad \frac{\Delta}{5} \phi^{2h+1} > 1 \quad \text{and} \quad k > \frac{1}{2} \log_{\phi} r.$$

The second equality may be rewritten as

$$(8.10) \quad \phi^{2k} > r.$$

From the first inequality of (8.9) and (8.10) it follows that

$$\frac{\Delta}{5} \phi^{2h+1} \phi^{2k} = \frac{\Delta}{5} \phi^{2(h+k)+1} > r.$$

Hence

$$\frac{\Delta}{5} \phi^{2m+1} > r.$$

Since

$$r > \frac{1}{\sin \frac{\pi}{r+2}} \quad \text{for } r \geq 2,$$

$m \geq p$  for  $r$  sufficiently large and the proof is concluded.  $\square$

ACKNOWLEDGMENTS. We are indebted to Xu Kang, Alberto Setti and Giancarlo Travaglini, for the help they gave us in preparing this paper.

#### REFERENCES

- [1] AKRITAS, A. G. Vincent's theorem in algebraic manipulation. Ph. D. Thesis, Operation Research Program, North Carolina State University, Raleigh, N.C., 1978.
- [2] ——— A new method for polynomial real root isolation<sup>20</sup>). *Proceedings of the 16<sup>th</sup> annual southeast regional ACM conference*, Atlanta, Georgia, April 1978, 39–43.
- [3] ——— A correction on a theorem by Uspensky. *Bull. Soc. Math. Grèce (N.S.)* 19 (1978), 278–285.
- [4] ——— Reflections on a pair of theorems by Budan and Fourier. *Math. Mag.* 55 (1982), 292–298.
- [5] ——— There is no "Uspensky's method". Extended abstract. *Proceedings of the 1986 Symposium on Symbolic and Algebraic Computation*, Waterloo, Ontario, Canada, 1986, 88–90.
- [6] ——— The role of the Fibonacci sequence in the isolation of the real roots of polynomial equations. *Applications of Fibonacci Numbers*, vol. 3, 1988.
- [7] ——— *Elements of Computer Algebra with Applications*. John Wiley & Sons, New York, 1989.
- [8] AKRITAS, A. G. and S. D. DANIELOPOULOS. On the forgotten theorem of Mr. Vincent. *Historia Math.* 5 (1978), 427–435.
- [9] AKRITAS, A. G. and S. D. DANIELOPOULOS. A converse rule of signs for polynomials. *Computing* 34 (1985), 283–286.

<sup>20</sup>) This paper won the first prize in the student paper competition.