

Zeitschrift: L'Enseignement Mathématique
Band: 44 (1998)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THE ORIGIN OF REPRESENTATION THEORY
Kapitel: 1. Introduction
Autor: CONRAD, Keith
DOI: <https://doi.org/10.5169/seals-63909>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 20.11.2024

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

THE ORIGIN OF REPRESENTATION THEORY

by Keith CONRAD

ABSTRACT. Representation theory was created by Frobenius about 100 years ago. We describe the background that led to the problem which motivated Frobenius to define characters of a finite group and show how representation theory solves the problem. The first results about representation theory in characteristic p are also discussed.

1. INTRODUCTION

Characters of finite abelian groups have been used since Gauss in the beginning of the 19th century, but it was only near the end of that century, in 1896, that Frobenius extended this concept to finite nonabelian groups [21]. Frobenius' approach to group characters is not in common use today, although some of his ideas that were overlooked for a long period have recently been revived [30].

Here we trace the development of the problem whose solution led Frobenius to introduce characters of finite groups, show how this problem can be solved using classical representation theory of finite groups, and indicate some relations between this problem and modular representations.

Other surveys on the origins of representation theory are by Curtis [7], Hawkins [24, 25, 26, 27], Lam [32], Ledermann [35] and van der Waerden [38]. While Curtis describes the development of modular representation theory focusing on the work of Brauer, we examine the earlier work in characteristic p of Dickson.