

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 45 (1999)  
**Heft:** 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** HARMONIC ANALYSIS ON VECTOR BUNDLES OVER  $Sp(1,n)/Sp(1) \times Sp(n)$   
**Autor:** van Dijk, G. / PASQUALE, A.  
**Kapitel:** 2. The convolution algebra  $D(G; \lambda_I)$   
**DOI:** <https://doi.org/10.5169/seals-64447>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

**Download PDF:** 03.01.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

for  $f \in C_c(G)$ . We adopt the usual notation  $C_c(G)$  for the space of continuous functions on  $G$  with compact support. In the above formulas,  $dn = dw dz$  ( $n = n(w, z)$ ) and  $dk$  is the normalized Haar measure on  $K$ .

Let

$$K_1 = \left\{ \begin{bmatrix} u & 0 \\ 0 & I \end{bmatrix} : u \in \text{Sp}(1) \right\}, \quad K_2 = \left\{ \begin{bmatrix} 1 & 0 \\ 0 & U \end{bmatrix} : U \in \text{Sp}(n) \right\}.$$

Then every  $g \in G$  can be written as  $g = k_1 k_2 a_t k'_2$  for uniquely determined  $k_1 \in K_1$ ,  $t \geq 0$  and for some  $k_2, k'_2 \in K_2$ . Writing  $g = [g_{ij}]_{i,j=0}^n$ , we have

$$(1.5) \quad k_1 = \frac{g_{00}}{|g_{00}|} \quad \text{and} \quad \cosh t = |g_{00}|.$$

If  $g \notin K$ , then  $t > 0$  and  $k_2, k'_2$  are uniquely determined modulo the subgroup

$$\left\{ \begin{bmatrix} 1 & 0 & 0 \\ 0 & V & 0 \\ 0 & 0 & 1 \end{bmatrix} : V \in \text{Sp}(n-1) \right\}.$$

After  $dg$  is normalized according to (1.3), the corresponding integral formula is

$$(1.6) \quad \int_G f(g) dg = \frac{1}{2} \left(\frac{\pi}{4}\right)^{2n} \frac{1}{\Gamma(2n)} \int_{K_1} \int_{K_2} \int_0^\infty \int_{K_2} f(k_1 k_2 a_t k'_2) \Delta(t) dk_1 dk_2 dt dk'_2$$

where

$$(1.7) \quad \Delta(t) := 2^{2\rho} (\sinh t)^{4n-1} (\cosh t)^3.$$

## 2. THE CONVOLUTION ALGEBRA $\mathcal{D}(G; \chi_l)$

Let  $\mathbf{N}/2$  be the set of nonnegative half-integers  $\{0, 1/2, 1, 3/2, \dots\}$ . Since  $K_1 \cong \text{Sp}(1)$  is isomorphic to  $\text{SU}(2)$ ,  $\mathbf{N}/2$  parametrizes the set of equivalence classes of unitary irreducible representations of  $K_1$ . We denote with the same symbol  $\tau_l$  either the equivalence class corresponding to the parameter  $l$  or a fixed representative for it. Thus  $\tau_l$  is a unitary irreducible representation of  $K_1$  in a Hilbert space  $V_l$  of dimension  $d_l = 2l + 1$ . We extend  $\tau_l$  to a representation of  $K$  by setting  $\tau_l \equiv \mathbf{1}$  on  $K_2$ . Each  $\tau_l$  is self-dual, i.e. unitarily equivalent to its contragredient representation. It follows in particular that the character  $\chi_l = \text{tr } \tau_l$  of  $\tau_l$  satisfies  $\chi_l(k^{-1}) = \chi_l(k)$ ,  $k \in K$ .

We denote by  $\mathcal{D}(G; \tau_l)$  the convolution algebra of the compactly supported  $C^\infty$  maps  $F: G \rightarrow \text{End}(V_l)$  satisfying

$$(2.8) \quad F(kxk') = \tau_l(k)F(x)\tau_l(k') \quad (k, k' \in K, x \in G).$$

Let  $\mathcal{D}(G)$  be the convolution algebra of the  $C^\infty$  compactly supported complex valued functions on  $G$ . Then  $\mathcal{D}(G; \tau_l)$  is isomorphic to the subalgebra  $\mathcal{D}(G; \chi_l)$  of  $\mathcal{D}(G)$  consisting of the functions  $f \in \mathcal{D}(G)$  satisfying

$$(2.9) \quad f^0 = f$$

and

$$(2.10) \quad f * d_l \chi_l = f,$$

where

$$(2.11) \quad f^0(x) := \int_K f(kxk^{-1}) dk.$$

The isomorphism is given by  $F \mapsto d_l \operatorname{tr} F$  (see e.g. [Dij], Theorem 1.1).

The commutativity of the algebra  $\mathcal{D}(G; \chi_l)$  can be deduced from the fact that the restriction  $\tau_l|_M$  of  $\tau_l$  to  $M$  is multiplicity free (according to the general criterion by Deitmar, cf. [Dei] Theorem 3, the commutativity of  $\mathcal{D}(G; \chi_l)$  and the multiplicity freeness of  $\tau_l|_M$  are in fact equivalent). An elementary direct argument by Takahashi proves the commutativity of a convolution algebra which is slightly bigger than  $\mathcal{D}(G; \chi_l)$ . Let  $\mathcal{D}_1(G)$  be the subalgebra of  $\mathcal{D}(G)$  consisting of the functions  $f \in \mathcal{D}(G)$  satisfying

$$(2.12) \quad f(k_2 k_1 g k_1^{-1} k_2') = f(g), \quad g \in G, k_1 \in K_1, k_2 \in K_2.$$

Clearly  $\mathcal{D}(G; \chi_l) \subset \mathcal{D}_1(G)$ . Moreover  $\mathcal{D}_1(G) = \oplus_l \mathcal{D}(G; \chi_l)$ . Showing that

$$(2.13) \quad f(g^{-1}) = f(g) \quad \text{for all } f \in \mathcal{D}_1(G) \text{ and } g \in G,$$

one proves the following lemma.

2.1. LEMMA ([T2], Proposition 1). *The algebra  $\mathcal{D}_1(G)$  is commutative.*

2.2. LEMMA (cf. [T2], Lemma 2). *For every function  $f \in \mathcal{D}(G; \chi_l)$ ,*

$$(2.14) \quad f(g) = f(k_1 a_t) = \frac{1}{d_l} \chi_l(k_1) f(a_t)$$

*if  $g = k_1 k_2 a_t k_2'$  ( $k_1 \in K_1; k_2, k_2' \in K_2; t \in \mathbf{R}$ ).*

*Proof.* For  $F \in \mathcal{D}(G; \tau_l)$ ,  $F(a_t)$  is a scalar multiple of the identity, since it commutes with  $\tau_l(k_1)$  for all  $k_1 \in K_1$ .  $\square$

2.3. REMARK. Formula (2.13) and Lemma 2.2 remain true for all continuous functions  $f$  such that  $f = f^0$ ,  $f * d_l \chi_l = f$ .