

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 45 (1999)  
**Heft:** 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** GENERALIZED FØLNER CONDITION AND THE NORMS OF  
RANDOM WALK OPERATORS ON GROUPS  
**Autor:** UK, Andrzej  
**Kapitel:** 3.1 Existence of eigenfunctions  
**DOI:** <https://doi.org/10.5169/seals-64454>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

**Download PDF:** 23.12.2024

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

This implies that

$$\frac{1}{\#S} |A|_{f^2} = \sum_{\gamma \in A} f^2(\gamma),$$

$$\frac{1}{\lambda(\#S)^2} |\partial A|_{f^2} \leq \sum_{\gamma \in \partial A} f^2(\gamma) \leq \lambda |\partial A|_{f^2}.$$

By Theorem 2, the first condition implies the second one.  $\square$

REMARK. The proof of Theorem 3 can easily be generalized to the case where  $P$  is a convolution operator with a finitely supported probability measure.

### 3. REMARKS

We will now make some comments about Theorems 2 and 3. We will state some theorems about the existence of eigenfunctions for the Markov operator and discuss whether one can take in the generalized Følner condition the eigenfunctions to be in  $L^2(X, \mu)$ .

For simplicity we will suppose that  $X$  is a connected, locally finite graph (i.e. the degree of each vertex is finite) and we consider the *simple random walk* going with equal probability from one vertex to any of its neighbors. We associate with this random walk the simple random walk operator  $P$  defined by

$$Pf(v) = \frac{1}{N(v)} \sum_{w \sim v} f(w) \quad \text{for } f \in l^2(X, N)$$

where  $N(v)$  is the degree of vertex  $v$  in  $X$  (i.e. the number of edges adjacent to  $v$ ), where  $w \sim v$  means that  $w$  and  $v$  are connected by an edge and where  $l^2(X, N)$  is the space of real-valued functions  $f$  on the vertices of  $X$  such that  $\sum_{x \in X} f^2(x)N(x)$  is finite.

#### 3.1 EXISTENCE OF EIGENFUNCTIONS

THEOREM 4 ([20]). *Let  $X$  be an infinite, locally finite graph and let  $P$  be the simple random walk operator on  $l^2(X, N)$ . For any  $\lambda \geq \|P\|$  there exists a positive eigenfunction  $f$  of  $P$  with eigenvalue  $\lambda$ , i.e.*

$$Pf(x) = \lambda f(x) \quad \text{and } f(x) > 0 \text{ for } x \in X.$$

*For  $\lambda < \|P\|$  there are no positive eigenfunctions of  $P$  with eigenvalue  $\lambda$ .*

*Proof.* There are several proofs of this theorem. In [20] one can find the proof where the analogue of Perron-Frobenius theory is developed and in [11] the truncation method is used.  $\square$

### 3.2 EIGENFUNCTIONS IN $l^2$

One can ask whether the positive eigenfunctions of the random walk operator are in  $l^2(X, N)$ . The answer is no in the case when  $X$  is the Cayley graph of an infinite group  $\Gamma$  (see Theorem 5). But in the general case there are examples of eigenfunctions which are in  $l^2(X, N)$  (see Proposition 2).

#### 3.2.1 THE CASE OF GROUPS

**THEOREM 5.** *Let  $f$  be a positive eigenfunction of the simple random walk operator  $P$  on the group  $\Gamma$  generated by a finite symmetric set  $S$ , i.e.  $Pf = \lambda f$ . If  $\Gamma$  is infinite then*

$$\sum_{\gamma \in \Gamma} f^2(\gamma) = +\infty.$$

*Proof.* Suppose the contrary, i.e. that there is a positive eigenfunction  $f$  of the operator  $P$  for which the  $l^2$  norm is finite:

$$\begin{aligned} Pf_0 &= \lambda f_0, \\ \sum_{\gamma \in \Gamma} f_0^2(\gamma) &< +\infty. \end{aligned}$$

The second condition implies that  $f_0$  is not constant and so there are  $\gamma_0, \gamma_1 \in \Gamma$  such that

$$f_0(\gamma_0) < f_0(\gamma_1).$$

Let us define the function  $f_1$  as a translation of  $f_0$  by  $\gamma_0\gamma_1^{-1}$ , i.e.

$$f_1(\gamma) = f_0(\gamma_0\gamma_1^{-1}\gamma).$$

The function  $f_1$ , being the translation of  $f_0$ , is an eigenfunction of  $P$ , i.e.

$$Pf_1 = \lambda f_1.$$

So the function  $\tilde{f}$  defined as follows:

$$\tilde{f}(\gamma) = \max\{f_0(\gamma), f_1(\gamma)\},$$

satisfies

$$P\tilde{f} \geq \lambda\tilde{f}.$$