

Zeitschrift: L'Enseignement Mathématique
Band: 45 (1999)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: GENERALIZED FØLNER CONDITION AND THE NORMS OF
RANDOM WALK OPERATORS ON GROUPS

Bibliographie

Autor: UK, Andrzej
DOI: <https://doi.org/10.5169/seals-64454>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 15.10.2024

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

$$(16) \quad a_{n+1} = |S_{n+1}|g^2(n+1) \leq (k-1)|S_n|g^2(n+1) = \left(1 + \frac{k-2}{(k-2)n+k}\right)^2 a_n.$$

We have to show that

$$(17) \quad \frac{\sum_{v \in S_{n+1}} f^2(v)}{\sum_{v \in B_n} f^2(v)} = \frac{a_{n+1}}{a_1 + \cdots + a_n} \xrightarrow{n \rightarrow \infty} 0.$$

It is a standard exercise to show that (16) implies (17). \square

Let f_n be the sequence of functions which are restrictions of f to the vertices that are at a distance not greater than n :

$$f_n = f|_{B_n}.$$

By Lemma 6 and Lemma 7 it follows that

$$\limsup_{n \rightarrow +\infty} \frac{\|Pf_n\|_{l^2(X,N)}}{\|f_n\|_{l^2(X,N)}} \geq \frac{2\sqrt{k-1}}{k},$$

which proves Theorem 8. \square

Some examples of upper bounds on the norm of the simple random walk operator on graphs and their comparison with the lower bound from Theorem 8 can be found in [22].

REFERENCES

- [1] AOMOTO, K. and Y. KATO. Green functions and spectra on free products of cyclic groups. *Ann. Inst. Fourier (Grenoble)* 38, 1 (1988), 59–85.
- [2] BOUGEROL, PH. and L. ÉLIE. Existence of positive harmonic functions on groups and on covering manifolds. *Ann. Inst. H. Poincaré Probab. Statist.* 31, No. 1 (1995), 59–80.
- [3] CARTWRIGHT, D. I. and P. M. SOARDI. Random walks on free products, quotients and amalgams. *Nagoya Math. J.* 102 (1986), 163–180.
- [4] FØLNER, E. On groups with full Banach mean value. *Math. Scand.* 3 (1955), 243–254.
- [5] GOODMAN, F. M., P. DE LA HARPE and V. F. R. JONES. Coxeter graphs and towers of algebras. *Math. Sci. Res. Inst. Publ.* 14. Springer-Verlag (1989).
- [6] GROMOV, M. Groups of polynomial growth and expanding maps. *Inst. Hautes Études Sci. Publ. Math.* 53 (1981), 53–78.
- [7] KAIMANOVICH, V. Dirichlet norms, capacities and generalized isoperimetric inequalities for Markov operators. *Potential Anal.*, 1 (1992), 61–82.

- [8] KESTEN, H. Full Banach mean values on countable groups. *Math. Scand.* 7 (1959), 146–156.
- [9] ——— Symmetric random walks on groups. *Trans. Amer. Math. Soc.* 92 (1959), 336–354.
- [10] MILNOR, J. A note on curvature and fundamental group. *J. Differential Geom.* 2 (1968), 1–7.
- [11] MOHAR, B. and W. WOESS. A survey on spectra of infinite graphs. *Bull. London Math. Soc.* 21 (1989), 209–234.
- [12] POPA, S. On Connes' joint distribution trick. *L'Enseignement Math.* (2) 44 (1998), no. 1–2, 57–70.
- [13] SALOFF-COSTE, L. Marches aléatoires sur les groupes discrets. Cours de DEA, Toulouse 1995.
- [14] SALOFF-COSTE, L. and W. WOESS. Transition operators, groups, norms, and spectral radii. *Pacific J. Math.* 180, No. 2 (1997), 333–367.
- [15] SALOFF-COSTE, L. and W. WOESS. Computing norms of group-invariant transition operators, *Combin. Probab. Comput.* 5 (1996), 161–178.
- [16] SAWYER, S. Isotropic random walks in a tree. *Z. Wahrsch. Verw. Gebiete* 42, No. 4 (1978), 279–292.
- [17] SOARDI, P.M. The resolvent for simple random walks on the free product of two discrete groups. *Math. Z.* 192 (1986), 109–116.
- [18] ŠVARC, A.S. Volume invariants of coverings. *Dokl. Akad. Nauk* 105 (1955), 32–34.
- [19] VAROPOULOS, N.TH., L. SALOFF-COSTE and T. COULHON. *Analysis and Geometry on Groups*. Cambridge University Press, 1992.
- [20] VERE-JONES, D. Ergodic properties of nonnegative matrices. *Pacific J. Math.* 22, No. 2 (1967), 361–386.
- [21] WOESS, W. Nearest neighbour random walks on free products of discrete groups. *Boll. Un. Mat. Ital. B(6)* 5 (1986), 961–982.
- [22] ŽUK, A. On the norms of random walks on planar graphs. *Ann. Inst. Fourier (Grenoble)* 47, 5 (1997), 1463–1490.

(Reçu le 15 juin 1998; version révisée reçue le 29 avril 1999)

Andrzej Żuk

CNRS, École Normale Supérieure de Lyon
Unité de Mathématiques Pures et Appliquées
46, allée d'Italie
F-69364 Lyon Cedex 07
France
e-mail: azuk@umpa.ens-lyon.fr