

Zeitschrift: L'Enseignement Mathématique
Band: 45 (1999)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: LOCAL-GLOBAL PRINCIPLE FOR NORMS FROM CYCLIC EXTENSIONS OF $\mathbb{Q}(t)$ (A DIRECT, CONSTRUCTIVE AND QUANTITATIVE APPROACH)
Kapitel: 4. An example for the non-cyclic case
Autor: ZANNIER, Umberto
DOI: <https://doi.org/10.5169/seals-64456>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 15.10.2024

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

$$N(r, b_1, \dots, b_d) = f(r)\mu$$

where $\mu \in \mathbf{Q}_v$ is very close to 1; in fact $f(r)$ is near to $f(a_v)$, which is nonzero. By the previous remarks, μ^{-1} is in the image of $N(r, x_1, \dots, x_d)$ on \mathbf{Q}_v^d , hence the same must be true for $f(r)$, by the basic multiplicative identity for N . In particular $f(r)$ will be a norm from $\mathbf{Q}_v(P_r)$ to \mathbf{Q}_v .

Let now S consist of the elements of \mathbf{Q} which are not poles or zeros of f , which satisfy $[\mathbf{Q}(P_s) : \mathbf{Q}] = d$ and which are sufficiently close (in the mentioned sense) to a_v , for each $v \in \Sigma$. We have proved that $f(s)$ is a norm from $\mathbf{Q}_v(P_s)$, for all $s \in S$ and for all places v . By Hasse's theorem, $f(s)$ is a norm from $\mathbf{Q}(P_s)$, so $S \subset N_f$. On the other hand $S \cap \mathbf{Z}$ contains the complement of a thin set in an arithmetic progression, whence N_f cannot satisfy the conclusion of the Theorem (or of Corollary 1), as required. \square

4. AN EXAMPLE FOR THE NON-CYCLIC CASE

We show that assuming that L/K is cyclic is essential in the Theorem (as in the number-field case, as shown in [CF, Ex. 5]).

To describe a counterexample, define $L = \mathbf{Q}(t, \sqrt{4t+3}, \sqrt{4t+7})$, $f(t) = t^2$. We proceed to show that $\mathbf{N} \subset N_f$. We have to show that for all large integers n , n^2 is a norm from $L(n) := \mathbf{Q}(\sqrt{4n+3}, \sqrt{4n+7})$. By [CF, Ex. 5.1 and 5.2, p.360] it is sufficient to show that the local degree $[L(n)_w : \mathbf{Q}_p]$ is 4 for some prime p . Observe that the Jacobi symbol $\left(\frac{4n+3}{4n+7}\right) = \left(\frac{-1}{4n+7}\right) = -1$. Hence there exists some prime p dividing $4n+7$ with an odd multiplicity and such that $\left(\frac{4n+3}{p}\right) = -1$. Then p ramifies in $L(n)$ and the residual degree is 2, proving the claim. Observe that the first conclusion of Corollary 1 does not hold for N_f .

On the other hand, t^2 is not a norm from L to K . Otherwise by [CF, Ex. 5.1] we could write t as the product of three norms from the three quadratic subfields of L . In other words we could write nontrivially

$$q^2(t)t = (a_1^2(t) - (4t+3)b_1^2(t))(a_2^2(t) - (4t+7)b_2^2(t))(a_3^2(t) - (4t+3)(4t+7)b_3^2(t)),$$

where $q, a_i, b_j \in \mathbf{Q}[t]$. We may suppose that a_i and b_i are coprime for each i , otherwise we can divide out a common factor. Now, putting $t = 0$ we get a contradiction.