

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 45 (1999)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** FREE GROUP ACTING ON  $Z^2$  WITHOUT FIXED POINTS  
**Autor:** Kenzi, Satô  
**Kapitel:** Proof of the main result  
**DOI:** <https://doi.org/10.5169/seals-64445>

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Siehe Rechtliche Hinweise.

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. Voir Informations légales.

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. See Legal notice.

**Download PDF:** 06.01.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

Though we cannot solve the above problem, the purpose of this note is to show that, if one replaces  $\mathbf{R}^2$  by  $\mathbf{Z}^2$ , the new problem has a positive solution. In fact, we will prove the following result, which shows somewhat more, namely that the action on  $\mathbf{Z}^2$  may be an action without non-trivial fixed points, rather than only locally commutative. We denote by  $\text{SA}_2(\mathbf{Z})$  the group of all transformations  $\vec{x} \mapsto A\vec{x} + \vec{a}$  of  $\mathbf{Z}^2$ , with  $A \in \text{SL}_2(\mathbf{Z})$  and  $\vec{a} \in \mathbf{Z}^2$ .

**THEOREM.** *The group  $\text{SA}_2(\mathbf{Z})$  has a free subgroup  $F_2$  of rank 2 which acts on  $\mathbf{Z}^2$  without non-trivial fixed points, namely the subgroup generated by*

$$\zeta: \begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} 7 & 3 \\ 9 & 4 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} 1 \\ -1 \end{pmatrix}$$

and

$$\eta: \begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} 94 & 39 \\ 147 & 61 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} 3 \\ 2 \end{pmatrix}.$$

The theorem implies the existence of a partition of  $\mathbf{Z}^2$  into three pieces  $P$ ,  $Q$  and  $R$  such that the six pieces  $P$ ,  $Q$ ,  $R$ ,  $P \sqcup Q$ ,  $Q \sqcup R$ ,  $R \sqcup P$  are pairwise  $F_2$ -congruent, without the axiom of choice ([S0], and Corollary 4.12 in [W]).

As observed in the discussion which follows Proposition 7.1 in [W], it is known that the above theorem does not carry over to  $\mathbf{R}^2$ ; more precisely, it is known that a subgroup of  $\text{SA}_2(\mathbf{R})$  which acts on  $\mathbf{R}^2$  without non-trivial fixed points is soluble, and consequently does not contain non-commutative free subgroups.

### PROOF OF THE MAIN RESULT

Recall that a matrix in  $\text{SL}_2(\mathbf{Z})$  is *hyperbolic* if the absolute value of its trace is strictly larger than 2, or equivalently if it has an eigenvalue of absolute value strictly larger than 1.

**LEMMA 0.** *The subgroup of  $\text{SL}_2(\mathbf{Z})$  generated by*

$$\begin{pmatrix} 7 & 3 \\ 9 & 4 \end{pmatrix} \text{ and } \begin{pmatrix} 94 & 39 \\ 147 & 61 \end{pmatrix}$$

*is free of rank 2 and all its elements distinct from the identity are hyperbolic.*

*Proof.* It is well-known that the subgroup of  $\mathrm{SL}_2(\mathbf{Z})$  generated by

$$\begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 5 & 2 \\ 2 & 1 \end{pmatrix}$$

is free of rank 2, and that all its elements distinct from the identity are hyperbolic. (See Appendix B in [K], [Ma], [MW], [N], or the proof of Theorem 6.8 in [W].) The lemma follows, because

$$\begin{pmatrix} 7 & 3 \\ 9 & 4 \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 5 & 2 \\ 2 & 1 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 94 & 39 \\ 147 & 61 \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}^2 \begin{pmatrix} 5 & 2 \\ 2 & 1 \end{pmatrix}^2$$

(see for example Exercice 12 of Section 1.4 in [MKS]).  $\square$

The following is elementary linear algebra.

LEMMA 1. *For  $A \in \mathrm{SL}_2(\mathbf{R})$  with  $\mathrm{tr} A \neq 2$  and for  $\vec{a} \in \mathbf{R}^2$ , the affine transformation*

$$\begin{cases} \mathbf{R}^2 & \rightarrow \mathbf{R}^2 \\ \vec{x} & \mapsto A\vec{x} + \vec{a} \end{cases}$$

*has a unique fixed point.*

Our preparations are complete.

*Proof of the main theorem.* The two transformations  $\zeta$  and  $\eta$  of our main result generate a group which is free of rank 2, by Lemma 0. As both these transformations fix the point

$$\begin{pmatrix} 2/3 \\ -5/3 \end{pmatrix} \in \mathbf{R}^2,$$

each element of the group they generate fix the same point. As this point is not in  $\mathbf{Z}^2$ , the theorem follows by Lemma 1.  $\square$

REMARK. Let  $\alpha, \beta \in \mathrm{SA}_2(\mathbf{Z})$  be as in the introduction. Then we can prove the main theorem by using the group generated by  $\alpha\beta^{-1}\alpha\beta^{-2}\alpha$  and  $\beta\alpha^{-1}\beta\alpha^{-2}\beta$ , because the transformations

$$\alpha\beta^{-1}\alpha\beta^{-2}\alpha : \begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} 13 & 22 \\ 10 & 17 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} -17 \\ -13 \end{pmatrix}$$

and

$$\beta\alpha^{-1}\beta\alpha^{-2}\beta : \begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} 17 & 10 \\ 22 & 13 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} -13 \\ -17 \end{pmatrix}$$

have a common fixed point  $\begin{pmatrix} 1/2 \\ 1/2 \end{pmatrix}$  in  $\mathbf{R}^2$  and the subgroup of  $\mathrm{SL}_2(\mathbf{Z})$  generated by

$$\begin{pmatrix} 13 & 22 \\ 10 & 17 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 17 & 10 \\ 22 & 13 \end{pmatrix}$$

is free of rank 2 and all its elements distinct from the identity are hyperbolic. See [K] and the following calculations:

$$\begin{pmatrix} 13 & 22 \\ 10 & 17 \end{pmatrix} = tu(tu^{-1})^3(tu)^2tu^{-1}tu,$$

$$\begin{pmatrix} 17 & 10 \\ 22 & 13 \end{pmatrix} = tu^{-1}(tu)^3(tu^{-1})^2tutu^{-1},$$

where

$$t = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \quad \text{and} \quad u = \begin{pmatrix} 0 & -1 \\ 1 & 1 \end{pmatrix}$$

with  $\langle t, u \rangle / \{\pm 1\} \cong \mathbf{Z}_2 * \mathbf{Z}_3$ .

The referee suggested to the author that the following could be shown (without the axiom of choice).

**COROLLARY.** *There exists a subset  $E_1$  of  $\mathbf{Z}^2$  such that, for every finite subset  $F$  of  $\mathbf{Z}^2$ , the symmetric difference of  $E_1$  and  $F$  is congruent to  $E_1$  relative to the group  $\mathrm{SA}_2(\mathbf{Z})$ .*

*Proof.* This is a consequence of our main result and of Theorem 2 in [My] ( $S = \mathbf{Z}^2$ ,  $G = \langle \zeta, \eta \rangle$ ,  $M = \{\zeta\eta, \zeta^2\eta^2, \zeta^3\eta^3, \dots\}$ ,  $\mathbf{F} = \{F \subseteq \mathbf{Z}^2 \mid F \text{ is finite}\}$ ).  $\square$

## REFERENCES

- [BT] BANACH, S. and A. TARSKI. Sur la décomposition des ensembles de points en parties respectivement congruentes. *Fund. Math.* 6 (1924), 244–277.
- [B] BOREL, A. On free subgroups of semi-simple groups. *L'Enseignement Math.* 29 (1983), 151–164.
- [D] DEKKER, Th. J. Decompositions of sets and spaces I. *Indag. Math.* 18 (1956), 581–589.
- [DS] DELIGNE, P. and D. SULLIVAN. Division algebras and the Hausdorff-Banach-Tarski paradox. *L'Enseignement Math.* 29 (1983), 145–150.
- [K] KUROSH, A. G. *The Theory of Groups*, vol. 2. Chelsea Publishing Company, New York, 1956.