

7. Corollaries of the isomorphism conjecture

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **46 (2000)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **13.09.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

REMARK 3. For G discrete the reduced C^* -algebra of G , denoted C^*G , comes equipped with a trace. An element in C^*G is a formal sum $\sum_{g \in G} \lambda_g g$ where $\lambda_g \in \mathbf{C}$. The trace of such an element is λ_1 where 1 is the identity element of G . This trace then induces a map

$$\text{tr}: K_0 C^*G \rightarrow \mathbf{R}.$$

Let Z be a proper G -manifold and let D be a G -invariant elliptic operator on Z . If ξ is the symbol of D then (Z, ξ) is a K -cocycle for (\cdot, G) and the Chern character defined above assigns to (Z, ξ)

$$\text{ch}(Z, \xi) \in H_*(BG; \mathbf{C}).$$

Let $\epsilon: BG \rightarrow \cdot$ be the map of BG to a point. Identify $H_*(\cdot, \mathbf{C}) = \mathbf{C}$ and consider

$$\epsilon_* \text{ch}(Z, \xi) \in \mathbf{C}.$$

The K -theory index of the elliptic operator D is an element of $K_0 C^*G$

$$\text{Index}(D) \in K_0 C^*G.$$

We then have the following formula for $\text{tr}[\text{Index}(D)]$:

$$\text{tr}[\text{Index}(D)] = \epsilon_* \text{ch}(Z, \xi).$$

For the special case when the action of G on Z is free this formula was obtained by M.F. Atiyah [3].

7. COROLLARIES OF THE ISOMORPHISM CONJECTURE

The conjecture stated in §2 above asserts that

$$\mu: K^*(X, G) \rightarrow K_*[C_0(X) \rtimes G]$$

is an isomorphism. Suppose that G is a discrete group and X is a point. The conjecture then asserts that $\mu: K^*(\cdot, G) \rightarrow K_* C^*G$ is an isomorphism where C^*G is the reduced C^* -algebra of G . Throughout this section G will be a discrete group and we shall consider some corollaries of the conjecture that $\mu: K^0(\cdot, G) \rightarrow K_0 C^*G$ is an isomorphism. “*Proof*” will mean “*Proof modulo the conjecture*”.

COROLLARY 1. *If G is torsion free then $\text{tr}: K_0 C^*G \rightarrow \mathbf{R}$ maps $K_0 C^*G$ onto the integers \mathbf{Z} .*

“*Proof*”. Let (Z, ξ) be a K -cocycle for (\cdot, G) . Let D be a G -invariant elliptic operator on Z whose symbol is ξ . By the definition of $\mu: K^0(\cdot, G) \rightarrow K_0 C^*G$ given in §2 above

$$\mu(Z, \xi) = \text{Index}(D).$$

If G is torsion free then the action of G on Z must be free. Hence Atiyah’s result applies [3] and $\text{tr}[\text{Index}(D)]$ must be an integer. Thus the surjectivity of $\mu: K^0(\cdot, G) \rightarrow K_0 C^*G$ implies that $\text{tr}: K_0 C^*G \rightarrow \mathbf{R}$ takes on only integer values. \square

COROLLARY 2. *If G is torsion free then there are no non-trivial projections in C^*G .*

“*Proof*”. A non-trivial projections in C^*G would give an element $\alpha \in K_0 C^*G$ with $0 < \text{tr}(\alpha) < 1$. \square

REMARK 3. For G torsion-free abelian, Corollary 2 can be proved by applying Pontrjagin duality. At the other extreme, Pimsner and Voiculescu [27] have proved that Corollary 2 is valid for a finitely generated free group.

In the statement of Corollary 2 it is essential that C^*G be the reduced C^* -algebra of G . Corollary 2 is not valid if one uses the maximal C^* -algebra C_{\max}^*G .

A classical conjecture [24] in the theory of group rings is that the group ring of a torsion-free group has no (non-trivial) divisors of zero. J. Cohen has observed that Corollaries 1 and 2 may be relevant to this zero-divisor conjecture.

If G has torsion then we conjecture that $\text{tr}: K_0 C^*G \rightarrow \mathbf{R}$ maps $K_0 C^*G$ onto the additive subgroup of \mathbf{Q} generated by all rational numbers of the form $\frac{1}{n}$, where n is the order of a finite subgroup of G . This would follow from the conjectured surjectivity of $K^0(\cdot, G) \rightarrow K_0 C^*G$ plus the unproved assertion that $\text{tr}[\text{Index}(D)]$ can only take on such values, where D is any G -invariant elliptic operator on a proper G -manifold.

COROLLARY 4. *The Novikov conjecture on homotopy invariance of higher signatures [11].*

“*Proof*”. Let M be a closed oriented C^∞ -manifold, $G = \pi_1(M)$ and let $f: M \rightarrow BG$ be the classifying map of the universal covering space of M .

The Novikov conjecture is that

$$\langle \mathbf{L}(M) \cup f^*(a), [M] \rangle$$

is an invariant of oriented homotopy type, where $\mathbf{L}(M)$ is the total \mathbf{L} class of TM and a is any element in $H^*(BG; \mathbf{Q})$.

Kasparov [19] and Miscenko-Fomenko [21] [22] define a map

$$K_0(BG) \rightarrow K_0 C^*G$$

and prove that the Novikov conjecture is implied by its rational injectivity. This enabled them to prove the Novikov conjecture for any discrete subgroup of a linear Lie group. The relation with our conjecture is clear from the following commutative diagram

$$\begin{array}{ccc} K_0(BG) & \longrightarrow & K_0 C^*G \\ & \searrow & \swarrow \\ & K^0(\cdot, G) & \end{array}$$

and the Proposition of §6 above. (In this factorization, the topological definition of K -homology given in [9] is being used.) \square

COROLLARY 5. *(Stable) Riemannian geometry conjectures of Gromov-Lawson-Rosenberg [30].*

For the same reason our conjecture implies the stable¹⁾ form of the Riemannian geometry conjectures of Gromov-Lawson-Rosenberg [30] on topological obstructions to the existence of metrics of positive scalar curvature.

8. TWISTING BY A 2-COCYCLE

This section is motivated by the papers [16], [26], [29], on the range of the trace for the C^* -algebra of the projective regular representation of a discrete group.

All of §2 adapts to the projective situation where together with the G -manifold X one is given a 2-cocycle $\gamma \in Z^2(X \rtimes G, S^1)$. For simplicity we

¹⁾ Paul Baum comments: It is important to emphasize "stable" because Thomas Schick has shown that the original unstable Gromov-Lawson-Rosenberg conjecture is false. On the other hand, Stephan Stolz (with contributions from J. Rosenberg and others) has proved that the real form of Baum-Connes implies the stable Gromov-Lawson-Rosenberg conjecture. Also, Max Karoubi and I have proved that the usual (i.e. complex K -theory) form of Baum-Connes implies the real form of Baum-Connes.