

3.1 $L_p(s, \lambda)$ FOR $\tau \in \bar{Q}_p$, $|\tau|_p \leq 1$

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **46 (2000)**

Heft 3-4: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **13.09.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

3. THE p -ADIC L -FUNCTION $L_p(s, t; \chi)$

In the following, we apply Theorem 2.7 to the sequence $\{b_n(\tau)\}_{n=0}^\infty$, where $b_n(\tau) = B_{n, \chi_n}(q\tau) - \chi_n(p)p^{n-1}B_{n, \chi_n}(p^{-1}q\tau)$, for $\tau \in \overline{\mathbf{Q}}_p$, $|\tau|_p \leq 1$, to show that there exists a power series $A_\chi(s, \tau) \in K_\tau[[s]]$, $K_\tau = \mathbf{Q}_p(\chi, \tau)$, which converges on $\{s \in \mathbf{C}_p : |s|_p < |p|_p^{1/(p-1)}|q|_p^{-1}\}$. From this we can prove the existence of a p -adic function, $L_p(s, \tau; \chi)$, that interpolates the values $L_p(1 - n, \tau; \chi) = -\frac{1}{n}b_n(\tau)$ for $n \in \mathbf{Z}$, $n \geq 1$, and converges in $\{s \in \mathbf{C}_p : |s - 1|_p < |p|_p^{1/(p-1)}|q|_p^{-1}\}$, except $s \neq 1$ if $\chi = 1$. After this we will show that there exists $L_p(s, \tau; \chi)$ for each $\tau \in \mathbf{C}_p$, $|\tau|_p \leq 1$, satisfying

$$L_p(1 - n, \tau; \chi) = -\frac{1}{n}b_n(\tau),$$

and converging in the domain above.

3.1 $L_p(s, \tau; \chi)$ FOR $\tau \in \overline{\mathbf{Q}}_p$, $|\tau|_p \leq 1$

Let p be prime, and let χ be a Dirichlet character with conductor f_χ . Let $\tau \in \overline{\mathbf{Q}}_p$, $|\tau|_p \leq 1$, and let $K_\tau = \mathbf{Q}_p(\chi, \tau)$, the field generated over \mathbf{Q}_p by adjoining τ and the values $\chi(a)$, $a \in \mathbf{Z}$. Since τ and each of the $\chi(a)$ are in $\overline{\mathbf{Q}}_p$, we see that K_τ is a finite extension of \mathbf{Q}_p in $\overline{\mathbf{Q}}_p$. For each $\tau \in \overline{\mathbf{Q}}_p$, $|\tau|_p \leq 1$, we shall derive our L -function $L_p(s, \tau; \chi)$ in a manner similar to that given for the derivation of $L_p(s; \chi)$ found in Chapter 3 of [13].

For $\tau \in \overline{\mathbf{Q}}_p$, $|\tau|_p \leq 1$, define the sequences $\{b_n(\tau)\}_{n=0}^\infty$ and $\{c_n(\tau)\}_{n=0}^\infty$ in K_τ according to

$$b_n(\tau) = B_{n, \chi_n}(q\tau) - \chi_n(p)p^{n-1}B_{n, \chi_n}(p^{-1}q\tau),$$

and

$$c_n(\tau) = \sum_{m=0}^n \binom{n}{m} (-1)^{n-m} b_m(\tau).$$

In order to derive our L -function $L_p(s, \tau; \chi)$, we will prove a particular bound on the magnitude of $c_n(\tau)$, but to do so, we shall need the following:

LEMMA 3.1. *Let $m, r \in \mathbf{Z}$, with $m \geq 0$ and $r \geq 1$. Then*

$$\sum_{a=0}^{p^r-1} a^m \equiv 0 \pmod{p^{r-1}},$$

where we take $0^0 = 1$ in the case of $a = 0$ and $m = 0$.

Proof. This is obvious for $m = 0$, so assume that $m \geq 1$. We shall prove this result for the remaining values of m by induction on r .

Since any sum of elements of \mathbf{Z} must also be in \mathbf{Z} , the lemma is true for $r = 1$. Now assume that the lemma holds for some $r \in \mathbf{Z}$, $r \geq 1$. By rewriting the sum

$$\sum_{a=0}^{p^{r+1}-1} a^m = \sum_{v=0}^{p-1} \sum_{u=0}^{p^r-1} (u + p^r v)^m,$$

and reducing this modulo p^r , we obtain

$$\begin{aligned} \sum_{a=0}^{p^{r+1}-1} a^m &\equiv \sum_{v=0}^{p-1} \sum_{u=0}^{p^r-1} u^m \pmod{p^r} \\ &\equiv p \sum_{u=0}^{p^r-1} u^m \pmod{p^r}. \end{aligned}$$

By our induction hypothesis we must then have

$$\sum_{a=0}^{p^{r+1}-1} a^m \equiv 0 \pmod{p^r},$$

and the lemma follows. \square

LEMMA 3.2. Let $\tau \in \mathbf{C}_p$, $|\tau|_p \leq 1$, and let $n \in \mathbf{Z}$, $n \geq 0$. For all $h \in \mathbf{Z}$, $h \geq 1$,

$$\frac{1}{q^h f_\chi} \sum_{\substack{a=1 \\ (a,p)=1}}^{q^h f_\chi} \chi(a) (\langle a + q\tau \rangle - 1)^n \equiv 0 \pmod{f_\chi^{-1} p^{-1} q^{n-1} \mathfrak{o}}.$$

Proof. This is obvious for $n = 0$ since writing

$$\sum_{\substack{a=1 \\ (a,p)=1}}^{q^h f_\chi} \chi(a) = \sum_{a=1}^{q^h f_\chi} \chi(a) - \sum_{a=1}^{p^{-1} q^h f_\chi} \chi(pa)$$

allows us to derive

$$\sum_{\substack{a=1 \\ (a,p)=1}}^{q^h f_\chi} \chi(a) = \begin{cases} p^{-1} q^h (p-1), & \text{if } \chi = 1 \\ 0, & \text{if } \chi \neq 1. \end{cases}$$

So let us assume that $n \geq 1$.

Let $h = 1$. Then $\langle a + q\tau \rangle \equiv 1 \pmod{q\mathfrak{o}}$ for all $a \in \mathbf{Z}$ such that $(a, p) = 1$ implies that

$$(\langle a + q\tau \rangle - 1)^n \equiv 0 \pmod{q^n\mathfrak{o}},$$

and the lemma holds for this case.

Now assume that $h \geq 1$. We can rewrite our sum as follows:

$$\sum_{\substack{a=1 \\ (a,p)=1}}^{q^h f_\chi} \chi(a) (\langle a + q\tau \rangle - 1)^n = \sum_{v=0}^{q^{h-1}-1} \sum_{\substack{u=1 \\ (u+vqf_\chi, p)=1}}^{qf_\chi} \chi(u + vqf_\chi) (\langle u + vqf_\chi + q\tau \rangle - 1)^n.$$

Since $|\tau|_p \leq 1$, we can write

$$\begin{aligned} \langle u + vqf_\chi + q\tau \rangle &= (u + vqf_\chi + q\tau) \omega^{-1} (u + vqf_\chi + q\tau) \\ &= (u + q\tau) \omega^{-1} (u + q\tau) + vqf_\chi \omega^{-1} (u + q\tau) \\ &= \langle u + q\tau \rangle + vqf_\chi \omega^{-1} (u). \end{aligned}$$

Thus

$$\begin{aligned} \sum_{\substack{a=1 \\ (a,p)=1}}^{q^h f_\chi} \chi(a) (\langle a + q\tau \rangle - 1)^n &= \sum_{\substack{u=1 \\ (u,p)=1}}^{qf_\chi} \chi(u) \sum_{v=0}^{q^{h-1}-1} (\langle u + q\tau \rangle - 1 + vqf_\chi \omega^{-1} (u))^n. \end{aligned}$$

By expanding, the inner sum on the right can be written

$$\begin{aligned} \sum_{v=0}^{q^{h-1}-1} (\langle u + q\tau \rangle - 1 + vqf_\chi \omega^{-1} (u))^n &= \sum_{k=0}^n \binom{n}{k} (\langle u + q\tau \rangle - 1)^{n-k} q^k f_\chi^k \omega^{-k} (u) \sum_{v=0}^{q^{h-1}-1} v^k. \end{aligned}$$

Since $(u, p) = 1$, we obtain the equivalence

$$q^k (\langle u + q\tau \rangle - 1)^{n-k} \equiv 0 \pmod{q^n\mathfrak{o}}$$

for each k , $0 \leq k \leq n$. Furthermore, by Lemma 3.1

$$\sum_{v=0}^{q^{h-1}-1} v^k \equiv 0 \pmod{p^{-1}q^{h-1}}$$

for each such k . Therefore

$$\sum_{v=0}^{q^{h-1}-1} (\langle u + q\tau \rangle - 1 + vqf_\chi \omega^{-1}(u))^n \equiv 0 \pmod{p^{-1}q^{n+h-1}\mathfrak{o}}.$$

This implies that

$$\sum_{\substack{a=1 \\ (a,p)=1}}^{q^h f_\chi} \chi(a) (\langle a + q\tau \rangle - 1)^n \equiv 0 \pmod{p^{-1}q^{n+h-1}\mathfrak{o}},$$

yielding the result. \square

We now derive our bound on the magnitude of $c_n(\tau)$.

PROPOSITION 3.3. *For all $\tau \in \mathbf{C}_p$, $|\tau|_p \leq 1$, and for $n \in \mathbf{Z}$, $n \geq 0$, we have $|c_n(\tau)|_p \leq |pqf_\chi|_p^{-1} |q|_p^n$.*

Proof. This follows in a manner similar to that given for the proof of the bound $|c_n(0)|_p \leq |q^2 f_\chi|_p^{-1} |q|_p^n$ found in [13] (Lemma 4 of Chapter 3). However, in this case we use Lemma 2.3 and the properties of χ and ω to derive

$$b_n(\tau) = \lim_{h \rightarrow \infty} \frac{1}{q^h f_\chi} \sum_{\substack{a=1 \\ (a,p)=1}}^{q^h f_\chi} \chi(a) \langle a + q\tau \rangle^n$$

for each $n \geq 0$, and thus

$$c_n(\tau) = \lim_{h \rightarrow \infty} \frac{1}{q^h f_\chi} \sum_{\substack{a=1 \\ (a,p)=1}}^{q^h f_\chi} \chi(a) (\langle a + q\tau \rangle - 1)^n$$

for each such n . From Lemma 3.2 we obtain

$$c_n(\tau) \equiv 0 \pmod{f_\chi^{-1} p^{-1} q^{n-1} \mathfrak{o}},$$

and thus the result. \square

For our immediate concern we only need this proposition to hold for all $\tau \in \overline{\mathbf{Q}}_p$ such that $|\tau|_p \leq 1$. However, later on we shall need it in the form in which we have it.

We are now ready to begin the construction of our L -function.

THEOREM 3.4. For each $\tau \in \overline{\mathbf{Q}}_p$, with $|\tau|_p \leq 1$, there exists a power series $A_\chi(s, \tau)$ in $K_\tau[[s]]$ such that the power series converges on $\{s \in \mathbf{C}_p : |s|_p < |p|_p^{1/(p-1)}|q|_p^{-1}\}$, and for each $n \in \mathbf{Z}$, $n \geq 0$, $A_\chi(n, \tau)$ satisfies

$$A_\chi(n, \tau) = B_{n, \chi_n}(q\tau) - \chi_n(p)p^{n-1}B_{n, \chi_n}(p^{-1}q\tau).$$

Proof. By Proposition 3.3, $|c_n(\tau)|_p \leq C|q|_p^n$ for all $n \geq 0$, where $C = |pqf_\chi|_p^{-1}$. Therefore we can apply Theorem 2.7 to the sequences $\{b_n(\tau)\}_{n=0}^\infty$ and $\{c_n(\tau)\}_{n=0}^\infty$ in $K_\tau = \mathbf{Q}_p(\chi, \tau)$, and for $\rho = |q|_p < |p|_p^{1/(p-1)}$, yielding this result. \square

Let us denote $\mathfrak{D} = \{s \in \mathbf{C}_p : |s - 1|_p < |p|_p^{1/(p-1)}|q|_p^{-1}\}$.

THEOREM 3.5. For each $\tau \in \overline{\mathbf{Q}}_p$, with $|\tau|_p \leq 1$, there exists a unique p -adic, meromorphic function $L_p(s, \tau; \chi)$ that can be expressed in the form

$$L_p(s, \tau; \chi) = \frac{a_{-1}(\tau)}{s - 1} + \sum_{n=0}^\infty a_n(\tau)(s - 1)^n,$$

where the power series converges in the domain \mathfrak{D} , having coefficients $a_n(\tau) \in \mathbf{Q}_p(\chi, \tau)$, with

$$a_{-1}(\tau) = \begin{cases} 1 - \frac{1}{p}, & \text{if } \chi = 1 \\ 0, & \text{if } \chi \neq 1. \end{cases}$$

Furthermore, for each $n \in \mathbf{Z}$, $n \geq 1$,

$$L_p(1 - n, \tau; \chi) = -\frac{1}{n} (B_{n, \chi_n}(q\tau) - \chi_n(p)p^{n-1}B_{n, \chi_n}(p^{-1}q\tau)).$$

Proof. Let

$$(13) \quad L_p(s, \tau; \chi) = \frac{1}{s - 1} A_\chi(1 - s, \tau)$$

with the $A_\chi(s, \tau)$ as in Theorem 3.4. Then from the properties of $A_\chi(s, \tau)$, the power series must converge in the given domain, and for $n \in \mathbf{Z}$, $n \geq 1$,

$$L_p(1 - n, \tau; \chi) = -\frac{1}{n} A_\chi(n, \tau) = -\frac{1}{n} (B_{n, \chi_n}(q\tau) - \chi_n(p)p^{n-1}B_{n, \chi_n}(p^{-1}q\tau)).$$

Note that

$$\begin{aligned} a_{-1}(\tau) &= A_\chi(0, \tau) = B_{0, \chi}(q\tau) - \chi(p)p^{-1}B_{0, \chi}(p^{-1}q\tau) \\ &= (1 - \chi(p)p^{-1})B_{0, \chi}, \end{aligned}$$

and thus

$$a_{-1}(\tau) = \begin{cases} 1 - \frac{1}{p}, & \text{if } \chi = 1 \\ 0, & \text{if } \chi \neq 1. \end{cases}$$

The uniqueness of $L_p(s, \tau; \chi)$ follows from Lemma 2.5. \square

At this point we have not completed our goal of showing that the p -adic function $L_p(s, \tau; \chi)$ exists for each $\tau \in \mathbf{C}_p$, $|\tau|_p \leq 1$. In order to prove this, we will need to study the coefficients, $a_n(\tau)$, of the power series expansion of $L_p(s, \tau; \chi)$ for each $\tau \in \overline{\mathbf{Q}}_p$, $|\tau|_p \leq 1$. From the results of this we will show that the function $L_p(s, \tau; \chi)$ exists for each $\tau \in \mathbf{C}_p$, $|\tau|_p \leq 1$, and for any sequence $\{\tau_i\}_{i=0}^{\infty}$ in $\overline{\mathbf{Q}}_p$, with $|\tau_i|_p \leq 1$, converging to τ , the values $L_p(1-n, \tau_i; \chi)$ converge to $L_p(1-n, \tau; \chi)$ for each $n \in \mathbf{Z}$, $n \geq 1$.

3.2 $L_p(s, \tau; \chi)$ FOR $\tau \in \mathbf{C}_p$, $|\tau|_p \leq 1$

Our previous work has been for $\tau \in \overline{\mathbf{Q}}_p$, $|\tau|_p \leq 1$. To extend this result to all $\tau \in \mathbf{C}_p$, $|\tau|_p \leq 1$, we need to find a way to express $a_n(\tau)$ so that it can be defined for these values of τ .

For $k \in \mathbf{Z}$, $k \geq 0$, the Stirling numbers of the first kind, $s(n, k)$, are defined by the generating function

$$(14) \quad \sum_{n=0}^{\infty} s(n, k) \frac{t^n}{n!} = \frac{1}{k!} (\log(1+t))^k.$$

Since the power series expansion of $\log(1+t)$ lacks a constant term, we must have $s(n, k) = 0$ whenever $0 \leq n < k$. We also have $s(n, n) = 1$ for all $n \geq 0$. The $s(n, k)$ are integers, where $n, k \in \mathbf{Z}$, $n \geq 0$, $k \geq 0$, and they satisfy the relation

$$(15) \quad \binom{x}{n} = \frac{1}{n!} \sum_{k=0}^n s(n, k) x^k.$$

For additional information on Stirling numbers of the first kind we refer the reader to [6], pp. 214–217.

LEMMA 3.6. *Let $\tau \in \overline{\mathbf{Q}}_p$, $|\tau|_p \leq 1$. For $n \in \mathbf{Z}$, $n \geq -1$,*

$$a_n(\tau) = (-1)^{n+1} \sum_{m=n+1}^{\infty} \frac{1}{m!} s(m, n+1) c_m(\tau).$$

Proof. From Corollary 2.8 we can write