

Objektyp: **ReferenceList**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **46 (2000)**

Heft 3-4: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **12.07.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

3. In order to make use of π_1 in locating other (not locally minimal) closed geodesics without non-degeneracy condition, one has to extend [Gr] to the non simply connected situation. When V is homeomorphic to $V_0 \times S^1$ and V_0 is simply connected, we can apply [Gr] directly and get $N_m(V) \geq C \log(m)$ for some $C > 0$. (Probably this is true when $H_1(V)$ is infinite or at least when $\pi_1(V) = \mathbf{Z}$.) The last estimate can be sharpened and we show this here for the simplest example when V_0 is the sphere S^3 and the proof is obvious [Gr].

Let V be homeomorphic to $S^3 \times S^1$. Then there exist closed geodesics $g_j^i \subset V$ (not necessarily simple) such that

1. Each g_j^i , $i, j = 1, 2, \dots$, represents $\gamma^i \in \pi_1(V)$ where γ is a generator in $\pi_1(V)$.
2. For each i the geodesic g_1^i is the shortest in its homotopy class.

Denote by $|g_j^i|$ the length of g_j^i .

3. $|g_1^{i+k}| + C \geq |g_1^i| + |g_1^k| \geq |g_1^{i+k}|$, where $C \geq 0$ and $i, k = 1, 2, \dots$
4. $|g_j^i| + C \geq |g_{j+1}^i| \geq |g_j^i|$ for some $C > 0$ and $i, j = 1, 2, \dots$
5. $||g_j^i| - |g_j^k|| \leq C|i - k|$ for some $C > 0$ and $i, j, k = 1, 2, \dots$
6. $|g_j^i| \geq \frac{i}{C}$ for some $C > 0$ and $i, j = 1, 2, \dots$

COROLLARY. *If V is as above, then $\limsup_{m \rightarrow \infty} \frac{N_m(V)}{m^2} \geq \text{const} > 0$.*

All our estimates give a rather poor approximation to the (unknown) reality. Probably, in most cases N_m grows exponentially. That is so, of course, for “ C^0 -generic” manifolds (“ C^0 -generic” is used for C^0 -generic manifolds having uncountably many closed geodesics).

REFERENCES

- [Fr] FRANKS, J. Anosov diffeomorphisms. In: *Global Analysis, Berkeley, 1968*. Proc. Symp. Pure Math. 14, Amer. Math. Soc. 1970, 61–93.
- [G-M] GROMOLL, D. and W. MEYER. Periodic geodesics on compact manifolds. *J. Differential Geom.* 3 (1969), 493–510.
- [Gr] GROMOV, M. Homotopical effects of dilatation. *J. Differential Geom.* 13 (1978), 303–310.
- [KI] KLINGENBERG, W. *Lectures on Closed Geodesics*. Third ed. Mathematisches Institut der Universität Bonn, 1977.

- [Ma] MANNING, A. Topological entropy and the first homology group. *Springer Lecture Notes in Math.* 468 (1974), 185–191.
- [Nie] NIELSEN, J. Über topologische Abbildungen geschlossener Flächen. *Abh. Math. Sem. Univ. Hamburg* 3 (1924), 246–260.
- [Pu] PUGH, C. On the entropy conjecture. *Springer Lecture Notes in Math.* 468 (1974), 257–262.
- [Sh] SHUB, M. Endomorphisms of compact manifolds. *Amer. J. Math.* 91 (1969), 185–199.
- [Sh-S] SHUB, M. and D. SULLIVAN. A remark on the Lefschetz fixed point formula. *Topology* 13 (1974), 189–191.

WHY THE APPENDICES WERE NOT WRITTEN :

AUTHOR'S APOLOGIES TO THE READERS

APPENDIX 2. The stable homeomorphism suggests a geometric link between the homotopy and topological invariance of Pontryagin classes, at least for manifolds with negative curvature but I did not manage to forge this to my satisfaction till 1996 (see [Gro₂]); also see [Fa-Jo] for a deeper analysis.

APPENDIX 3. One can define a notion of hyperbolicity for an automorphism α of an arbitrary finitely generated group Γ , such that (Γ, α) functorially defines a Bowen-Franks hyperbolic system (see [Gro₁]). Unfortunately, this class of (Γ, α) is rather limited, e.g. is not closed under free products and does not include hyperbolic automorphisms of surface groups. I still do not know what the right setting is.

APPENDIX 4. An obvious example of semi-hyperbolicity is provided by non-strictly expanding endomorphisms, where the geometric picture is rather clear. However, I still do not see a functorial description, in the spirit of the symbolic dynamics, of more general semi-hyperbolic systems, not even for the geodesic (or Weil chamber) flows on locally symmetric spaces (compare [B-G-S] and [Br-Ha]).

APPENDIX 5. The section on entropy was inspired by Manning's paper [Ma₁], but I was unaware of the prior paper by Dinaburg (see [Din]) that essentially contained the entropy estimate for geodesic flows (also discussed in [Ma₂]). On the other hand, estimating the entropy of an endomorphism (or an automorphism) f in terms of $f_*: \pi_1 \rightarrow \pi_1$ appears now much less clear than it seemed to me back in 1976. It is not hard to bound the entropy from below