

4. Existence of almost complex structures

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **46 (2000)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **12.07.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

It follows that

$$\mathfrak{o}_0(X_m \# X_n \# S^4 \times S^4, J_m + J_n + J'_0) = 3(m(m+1) + n(n+1))/7$$

(and $\mathfrak{o}_2 = 0$). With the given constraints on m and n , this can only be zero for $m = n = 0$ (even if we allow for the orientation of the summands to be changed).

For the allowed choices of m and n , this connected sum is homotopy equivalent to $\mathbf{HP}^2 \# \mathbf{HP}^2 \# S^4 \times S^4$. The fact that the homotopy equivalence $X_m, X_n \simeq X_0$ induces a homotopy equivalence of the connected sums is a simple consequence of the Whitehead theorem, since we are dealing with simply-connected manifolds. This concludes the proof of Proposition 6. \square

4. EXISTENCE OF ALMOST COMPLEX STRUCTURES

In this section we prove Theorem 4(b). We already know that condition (b)(i) is necessary. We now show that condition (b)(ii) is necessary.

Given an almost complex structure J on M , we have

$$2\chi(M) - 2c_1(J)c_3(J) + c_2(J)^2 - p_2(M) = 0$$

by (1). In the sequel we suppress M and J . In case II_0 , c_1 is a torsion class, so this simplifies to

$$2\chi + c_2^2 - p_2 = 0.$$

Squaring the relation $p_1 = c_1^2 - 2c_2$, and again observing that c_1 is a torsion class, we get $p_1^2 = 4c_2^2$. Multiplying the equation above by 4 and substituting p_1^2 for $4c_2^2$ yields condition (b)(ii).

In fact, this argument also shows that (b)(ii) is a sufficient condition. By (a) we have a stable a.c.s. \tilde{J} on M and thus a corresponding J_0 as in Section 3. If condition (b)(ii) holds, then reversing the argument just given we find

$$4\mathfrak{o}_0(M, J_0) = 2\chi(M) + c_2(J_0)^2 - p_2(M) = 0.$$

Since J_0 is induced by \tilde{J} , the stable part \mathfrak{o}_2 of the obstruction vanishes as well, so J_0 extends to an almost complex structure on M .

Next we prove that condition (b)(i) is sufficient for the existence of an a.c.s. We begin with a preparatory lemma.

LEMMA 10. *Let M be an 8-manifold with $b_2 > 0$ which satisfies the assumptions of Theorem 4(a). Then there is a family \tilde{J}_k , $k \in \mathbf{Z}$, of stable almost complex structures on M such that*

$$c_1(\tilde{J}_k) c_3(\tilde{J}_k) = c_1(\tilde{J}_0) c_3(\tilde{J}_0) + 4k$$

and

$$c_2(\tilde{J}_k) = c_2(\tilde{J}_0) \text{ modulo 2-torsion,}$$

so in particular $c_2(\tilde{J}_k)^2 = c_2(\tilde{J}_0)^2$.

Proof. Case I: By Theorem 3(a) we can find a stable a.c.s. \tilde{J}_0 such that the free part x of $c_1(\tilde{J}_0)$ is indivisible. By Poincaré duality there is an element $x' \in H^6(M; \mathbf{Z})$ such that $c_1(\tilde{J}_0)x' = xx' = 1$. Then the pair $(c_1(\tilde{J}_0), c_3(\tilde{J}_0) + 4kx')$, $k \in \mathbf{Z}$, still satisfies the assumptions of Theorem 3(a), so there are stable almost complex structures \tilde{J}_k , $k \in \mathbf{Z}$, with

$$c_1(\tilde{J}_k) = c_1(\tilde{J}_0) \text{ and } c_3(\tilde{J}_k) = c_3(\tilde{J}_0) + 4kx'.$$

This is the desired family, since the relation $p_1 = c_1^2 - 2c_2$ shows that c_2 is determined modulo 2-torsion by c_1 and p_1 .

Case II₊: Let \tilde{J}_0 be a stable a.c.s. such that $c_1(\tilde{J}_0)$ is torsion. Let x be an indivisible element of $H^2(M; \mathbf{Z})$ and x' a dual element of $H^6(M; \mathbf{Z})$, i.e. $xx' = 1$ and $yx' = 0$ for all y in a complement of $\mathbf{Z}x \subset H^2(M; \mathbf{Z})$ (of course x' depends on the choice of this complement). Then by Theorem 3(a) there exists a stable a.c.s. \tilde{J}_k with

$$c_1(\tilde{J}_k) = c_1(\tilde{J}_0) + 2x \text{ and } c_3(\tilde{J}_k) = c_3(\tilde{J}_0) + 2kx'.$$

This family has the desired properties. \square

Now, assuming that condition (b)(i) is satisfied, we choose a stable a.c.s. \tilde{J} on M as in the proof of the preceding lemma. Hence we get an a.c.s. J_1 on $M - D^8$ with

$$\sigma(M, J_1) = (a_1, 0) \in \mathbf{Z} \oplus \mathbf{Z}_2.$$

If a_1 is even, then by formulae (1) and (2) for σ and the preceding lemma, we can find a different a.c.s. J'_1 on $M - D^8$ with $\sigma(M, J'_1) = (0, 0)$, so that J'_1 extends to an a.c.s. on M .

We complete the proof of Theorem 4 by showing that a_1 has to be even. Since

$$\sigma(M \# \mathbf{H}P^2 \# \mathbf{H}P^2, J_1 + J_0 + J_0) = (a_1 - 2, 0),$$

we can find an 8-manifold M_2 and an a.c.s. J_2 on $M_2 - D^8$ with

$$\mathfrak{o}(M_2, J_2) = (-a_2, 0)$$

with $a_2 > 0$ of the same parity as a_1 , and

$$\chi(M_2) - \tau(M_2) = \chi(M) - \tau(M) \equiv 0 \pmod{4}.$$

Then

$$\mathfrak{o}(M_2 \# a_2 S^4 \times S^4, J_2 + a_2 J'_0) = (0, 0).$$

So $M_2 \# a_2 S^4 \times S^4$ admits an a.c.s. Now compute

$$\begin{aligned} (\chi - \tau)(M_2 \# a_2 S^4 \times S^4) &= \chi(M_2) + 2a_2 - \tau(M_2) \\ &\equiv 2a_2 \pmod{4}. \end{aligned}$$

By the necessity of condition (b) (i) we conclude $a_1 \equiv a_2 \equiv 0 \pmod{2}$. \square

REFERENCES

- [1] AUDIN, M. Exemples de variétés presque complexes. *L'Enseignement Math.* (2) 37 (1991), 175–190.
- [2] DESSAI, A. Some remarks on almost and stable almost complex manifolds. *Math. Nachr.* 192 (1998), 159–172.
- [3] EELLS, J. JR. and N. H. KUIPER. Manifolds which are like projective planes. *Inst. Hautes Études Sci. Publ. Math.* 14 (1962), 5–46.
- [4] GEIGES, H. Contact structures on 1-connected 5-manifolds. *Mathematika* 38 (1991), 303–311.
- [5] —— Applications of contact surgery. *Topology* 36 (1997), 1193–1220.
- [6] HEAPS, T. Almost complex structures on eight- and ten-dimensional manifolds. *Topology* 9 (1970), 111–119.
- [7] HILLER, H. Almost complex structures on four-dimensional complete intersections. *Bull. Austral. Math. Soc.* 29 (1984), 143–152.
- [8] HIRZEBRUCH, F. Über die quaternionalen projektiven Räume. *Sitzungsber. Bayer. Akad. Wiss. Math.-Naturwiss. Kl.* 27 (1953), 301–312.
- [9] —— Komplexe Mannigfaltigkeiten. *Proc. International Congress of Mathematicians* 1958, Cambridge University Press (1960), 119–136.
- [10] —— *Collected Papers*. Springer, Berlin, 1987.
- [11] HIRZEBRUCH, F. and H. HOPF. Felder von Flächenelementen in 4-dimensionalen Mannigfaltigkeiten. *Math. Ann.* 136 (1958), 156–172.
- [12] KAHN, P. J. Obstructions to the existence of almost X-structures. *Illinois J. Math.* 13 (1969), 336–357.
- [13] MASSEY, W. S. Obstructions to the existence of almost complex structures. *Bull. Amer. Math. Soc. (N.S.)* 67 (1961), 559–564.
- [14] MÜLLER, S. *Zur Topologie einfacher 8-Mannigfaltigkeiten*. Inaugural-Dissertation, Universität Zürich, 1998.
- [15] STEENROD, N. *The Topology of Fibre Bundles*. Princeton University Press, 1951.