

2.1 DISCRETE 4-VERTEX THEOREM

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **47 (2001)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **12.07.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

2. THEOREMS ON PLANE POLYGONS

In this section we formulate our results for plane polygonal curves. The proofs will be given in Section 4.1.

2.1 DISCRETE 4-VERTEX THEOREM

The *osculating circle* of a smooth plane curve at a point is the circle (or straight line) that has 3rd order of contact with the curve at the given point. One may say that the osculating circle goes through 3 infinitely close points; at a vertex the osculating circle passes through 4 infinitely close points. Moreover, a generic curve crosses the osculating circle at a generic point and stays on one side of it at a vertex. This well-known fact motivates the following definition.

Let P be a plane convex n -gon; throughout this section we assume that $n \geq 4$. Denote the consecutive vertices by V_1, \dots, V_n ; the subscripts are understood cyclically, that is, $V_{n+1} = V_1$, etc.

DEFINITION 2.1. A triple of vertices (V_i, V_{i+1}, V_{i+2}) is said to be *extremal*¹⁾ if V_{i-1} and V_{i+3} lie on the same side of the circle through V_i, V_{i+1}, V_{i+2} (this does not exclude the case where V_{i-1} or V_{i+3} belongs to the circle).




FIGURE 1

The next result follows from a somewhat more general theorem due to O. Musin and V. Sedykh [12] (see also [13]).

¹⁾ We have a terminological difficulty here: as we are dealing with polygons, we cannot use the term "vertex" in the same sense as in the smooth case; hence the term "extremal".

THEOREM 2.2. *Every plane convex polygon P has at least 4 extremal triples of vertices.*

EXAMPLE 2.3. If P is a quadrilateral then the theorem holds tautologically since the $(i - 1)^{\text{st}}$ vertex coincides with the $(i + 3)^{\text{rd}}$ for every i .

REMARK 2.4. An alternative approach to discretization of the 4-vertex theorem consists in inscribing circles in consecutive triples of sides of a polygon (the centre of such a circle is the intersection point of the bisectors of consecutive angles of the polygon). Then a triple of sides (l_i, l_{i+1}, l_{i+2}) is said to be *extremal* if the lines l_{i-1}, l_{i+3} either both intersect the corresponding circle or both fail to intersect it. With this definition an analogue of Theorem 2.2 holds true [19, 16], and this, in the limit, also provides the smooth 4-vertex theorem.

Both formulations, concerning circumscribed or inscribed circles, make sense on the sphere. Moreover, they are equivalent via projective duality.

2.2 DISCRETE THEOREM ON 6 AFFINE VERTICES

Five generic points in the plane determine a conic. Considering the plane as an affine part of the projective plane, the complement of the conic has two connected components. Let P be a plane convex n -gon; throughout this section we assume that $n \geq 6$. As in the previous section, we introduce the following definition.

DEFINITION 2.5. Five consecutive vertices V_i, \dots, V_{i+4} are said to be *extremal* if V_{i-1} and V_{i+5} lie on the same side of the conic through these 5 points (this does not exclude the case where V_{i-1} or V_{i+5} belongs to the conic).

If P is replaced by a smooth convex curve, and V_i, \dots, V_{i+4} are infinitely close points, we recover the definition of an affine vertex. Hence the following theorem is a discrete version of the smooth theorem on 6 affine vertices.

THEOREM 2.6. *Every plane convex polygon P has at least 6 extremal quintuples of vertices.*

EXAMPLE 2.7. If P is a hexagon then the theorem holds tautologically for the same reason as in Example 2.3.