

2. CONJUGATION AND PARTIAL CONJUGATION

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **47 (2001)**

Heft 3-4: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **10.08.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

well as the work of Mellor and Thurston, of course) shows the existence of non-trivial finite type link concordance invariants.

To extend the applicability of our general philosophy slightly, we find that the operation on the vector $\{\mu(rst)\}$ induced by reversing the orientation of each component of a string link is to change it by a negative sign followed by a translation whose translation vector's coordinates are quadratic polynomials in l_{ij} . If the dimension of the subspace generated by this vector together with the translation vectors of conjugations and partial conjugations is still less than $\binom{k}{3}$ for generic values of the linking numbers, and this is the case indeed, we can construct a non-trivial link-homotopy invariant polynomial which is changed by a sign when the orientation of each component of a link is reversed. We say that such a link invariant detects the invertibility for links. Recall that the reversion of the orientation of every component of a link does not change the quantum invariant associated with an irreducible representation of a semi-simple Lie algebra (see, for example, [8]). Thus our invariant is of finite type but is not determined by quantum invariants. The existence of a finite type knot invariant which detects the invertibility for knots is a major problem in the theory of finite type invariants (see, for example, [8] and [4]). We believe that finite type knot invariants can not detect the invertibility for knots.

It remains unclear whether we can have a complete set of link-homotopy invariant polynomials which determines uniquely link-homotopy classes of links. See [5] for an earlier attempt on this problem²). This problem could probably be translated to the problem of understanding the sublattice generated by the translation vectors of conjugations and partial conjugations. A better understanding of this sublattice might also be useful in answering the following question. If we let $\deg(l_{ij}) = 1$ and $\deg(\mu(rst)) = 2$, the link-homotopy invariant polynomial for $k = 6$ we construct in Section 3, which detects the invertibility for links, is a linear combination of 113,700 monomials of degree 22, homogeneous in both l_{ij} and $\mu(rst)$ and linear in $\mu(rst)$. Is there a shorter link-homotopy invariant polynomial detecting the invertibility for links?

2. CONJUGATION AND PARTIAL CONJUGATION

We first recall the classification of ordered, oriented links up to link-homotopy given in [3]. We will follow the notations of [3].

²) See [6] for another approach to the similar problem for surgery equivalence of links. Notice that both approaches attempted to reduce the indeterminacies of the $\bar{\mu}$ -invariants.

Let $\mathcal{H}(k)$ be the group of link-homotopy classes of ordered, oriented string links with k components. The components of a string link will be ordered by $1, 2, \dots, k$. Recall that a string link is a concordance of k marked points inside of the 2-disk D^2 to itself in $D^2 \times [0, 1]$, such that it has no closed component. Two string links are link-homotopic if they are homotopic in such a way that at any moment of the homotopy, different components remain disjoint (but they are allowed to have self-intersections). Two string links can be put together to form a new string link and this gives rise to a group structure on the set of all link-homotopy classes of string links. This is the group $\mathcal{H}(k)$.

A pure braid is by definition a string link of the same number of components. So we have a natural map from the pure braid group $P(k)$ of k components to $\mathcal{H}(k)$. It is shown in [3] that this natural map $P(k) \rightarrow \mathcal{H}(k)$ is onto.

Deletion of the i^{th} component of the string link gives rise to a group homomorphism $d_i: \mathcal{H}(k) \rightarrow \mathcal{H}(k-1)$. If $F(k)$ denotes the free group of rank k generated by x_1, x_2, \dots, x_k , the reduced free group $RF(k)$ is the quotient of $F(k)$ by adding relations $[x_i, x_i^g] = 1$ for all i and all $g \in F(k)$.

LEMMA 2.1. *There is a split short exact sequence of groups*

$$(1) \quad 1 \longrightarrow RF(k-1) \longrightarrow \mathcal{H}(k) \xrightarrow{d_i} \mathcal{H}(k-1) \longrightarrow 1$$

where $RF(k-1)$ is the reduced free group generated by $x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_k$.

Notice that the split exact sequence (1) depends on the deleted component so that there are k such split exact sequences altogether. A split exact sequence determines a semi-direct product decomposition

$$\mathcal{H}(k) = \mathcal{H}(k-1) \rtimes RF(k-1).$$

Conjugation in the group $\mathcal{H}(k)$ is defined as usual: A conjugation of $\sigma \in \mathcal{H}(k)$ by $\beta \in \mathcal{H}(k)$ is the element $\beta\sigma\beta^{-1} \in \mathcal{H}(k)$. A *partial conjugation* of $\sigma \in \mathcal{H}(k)$ is an element of the form $\theta h g h^{-1}$, where we write $\sigma = \theta g$ according to a decomposition $\mathcal{H}(k) = \mathcal{H}(k-1) \rtimes RF(k-1)$, for $\theta \in \mathcal{H}(k-1)$ and $g \in RF(k-1)$, and for an arbitrary $h \in RF(k-1)$.

To form the closure of a string link $\sigma \in \mathcal{H}(k)$, we may think of it as a pure braid in $P(k)$ and its closure will be the usual braid closure. The closure of $\sigma \in \mathcal{H}(k)$ is an ordered, oriented link of k components. It is not hard to see that every link-homotopy class of ordered, oriented links with k components can be realized as the closure of an element in $\mathcal{H}(k)$, and thus the closure of a pure braid in $P(k)$. One of the main results of [3] is the following classification theorem.

THEOREM 2.2. *Let $\sigma, \sigma' \in \mathcal{H}(k)$. Then the closures of σ and σ' are link-homotopic as ordered, oriented links if and only if there is a sequence $\sigma = \sigma_0, \sigma_1, \dots, \sigma_n = \sigma'$ of elements of $\mathcal{H}(k)$ such that σ_{j+1} is either a conjugation or a partial conjugation of σ_j .*

For a group G , we will denote by G_n the n^{th} term of the lower central series of G , i.e. $G_1 = G$ and $G_{n+1} = [G_n, G]$, the normal subgroup of G generated by elements of the form $[g, h] = ghg^{-1}h^{-1}$ for all $g \in G_n$ and $h \in G$. A group G is nilpotent of class n if $G_{n+1} = 1$ but $G_n \neq 1$. We summarize some known facts about the group structures of $\mathcal{H}(k)$ in the following lemma.

LEMMA 2.3. 1) $\mathcal{H}(k)$ is torsion free and nilpotent of class $k - 1$.

2) Corresponding to a decomposition $\mathcal{H}(k) = \mathcal{H}(k - 1) \rtimes RF(k - 1)$, we have

$$\mathcal{H}(k)_n = \mathcal{H}(k - 1)_n \rtimes RF(k - 1)_n.$$

3) $\mathcal{H}(k)_{n-1}/\mathcal{H}(k)_n$ is a free abelian group of rank $(n - 2)! \binom{k}{n}$.

For $\sigma \in \mathcal{H}(k)$, its image in $\mathcal{H}(k)/\mathcal{H}(k)_3$ can be described by $\binom{k}{2} + \binom{k}{3}$ integers. These integers are linking numbers l_{ij} , for $1 \leq i < j \leq k$, and Milnor's triple linking numbers $\mu(rst)$, for $1 \leq r < s < t \leq k$. We want to have them defined precisely and understand how they change when σ is changed by a conjugation or a partial conjugation.




FIGURE 1
The pure braid τ_{rs}

We will denote by $\tau_{rs} = \tau_{sr}$, for $1 \leq r < s \leq k$, the pure braid depicted in Figure 1. Let $\sigma \in \mathcal{H}(k)/\mathcal{H}(k)_3$. For $1 \leq r < s < t \leq k$, after deleting

all components other than the r, s, t -th components, σ can be written in the following normal form

$$(2) \quad \sigma = \tau_{rs}^\alpha \tau_{rt}^\beta \tau_{st}^\gamma [\tau_{rt}, \tau_{st}]^\delta,$$

where $\alpha = l_{rs}$, $\beta = l_{rt}$, $\gamma = l_{st}$. By definition, we have $\delta = \mu(rst)$ for $\sigma \in \mathcal{H}(k)$.

LEMMA 2.4. *In $\mathcal{H}(k)/\mathcal{H}(k)_3$, if r', s', t' is a permutation of r, s, t and ϵ is the sign of the permutation, then*

$$[\tau_{r't'}, \tau_{s't'}] = [\tau_{rt}, \tau_{st}]^\epsilon.$$

Furthermore, we have

$$[\tau_{rt}^\eta, \tau_{st}] = [\tau_{rt}, \tau_{st}]^\eta.$$

This lemma is useful in the following calculation and its proof is straightforward.

To understand how $\mu(rst)$ changes under the conjugation, we only need to calculate the conjugation of $\sigma \in \mathcal{H}(k)/\mathcal{H}(k)_3$ under the normal form (2) by $\tau_{rs}, \tau_{rt}, \tau_{st}$. This calculation is straightforward:

$$\begin{aligned} \tau_{rs}\sigma\tau_{rs}^{-1} &= \tau_{rs}\tau_{rs}^\alpha\tau_{rt}^\beta\tau_{st}^\gamma[\tau_{rt}, \tau_{st}]^\delta\tau_{rs}^{-1} \\ &= \tau_{rs}^\alpha\tau_{rt}^\beta\tau_{st}^\gamma[\tau_{rs}, \tau_{rt}]^\beta[\tau_{rs}, \tau_{st}]^\gamma[\tau_{rt}, \tau_{st}]^\delta \\ &= \tau_{rs}^\alpha\tau_{rt}^\beta\tau_{st}^\gamma[\tau_{rt}, \tau_{st}]^{\delta+\beta-\gamma}; \\ \tau_{rt}\sigma\tau_{rt}^{-1} &= \tau_{rt}\tau_{rs}^\alpha\tau_{rt}^\beta\tau_{st}^\gamma[\tau_{rt}, \tau_{st}]^\delta\tau_{rt}^{-1} \\ &= \tau_{rs}^\alpha\tau_{rt}^\beta\tau_{st}^\gamma[\tau_{rt}, \tau_{rs}]^\alpha[\tau_{rt}, \tau_{st}]^\gamma[\tau_{rt}, \tau_{st}]^\delta \\ &= \tau_{rs}^\alpha\tau_{rt}^\beta\tau_{st}^\gamma[\tau_{rt}, \tau_{st}]^{\delta-\alpha+\gamma}; \\ \tau_{st}\sigma\tau_{st}^{-1} &= \tau_{st}\tau_{rs}^\alpha\tau_{rt}^\beta\tau_{st}^\gamma[\tau_{rt}, \tau_{st}]^\delta\tau_{st}^{-1} \\ &= \tau_{rs}^\alpha\tau_{rt}^\beta\tau_{st}^\gamma[\tau_{st}, \tau_{rs}]^\alpha[\tau_{st}, \tau_{rt}]^\beta[\tau_{rt}, \tau_{st}]^\delta \\ &= \tau_{rs}^\alpha\tau_{rt}^\beta\tau_{st}^\gamma[\tau_{rt}, \tau_{st}]^{\delta+\alpha-\beta}. \end{aligned}$$

We summarize the calculation into the following lemma.

LEMMA 2.5. *The change of $\mu(rst)$ under a conjugation is given by*

$$\text{Conjugation by } \tau_{rs}: \quad \mu(rst) \rightarrow \mu(rst) + l_{rt} - l_{st};$$

$$\text{Conjugation by } \tau_{rt}: \quad \mu(rst) \rightarrow \mu(rst) - l_{rs} + l_{st};$$

$$\text{Conjugation by } \tau_{st}: \quad \mu(rst) \rightarrow \mu(rst) + l_{rs} - l_{rt}.$$

Furthermore, $\mu(rst)$ will not change under a conjugation by τ_{ij} where $\{i, j\}$ and $\{r, s, t\}$ have at most one element in common.

The calculation of partial conjugations is slightly more complicated. We will start with partial conjugations by τ_{rt} and τ_{st} . These two operations are denoted by \mathbf{t}^r and \mathbf{t}^s , respectively. For $\sigma \in \mathcal{H}(k)/\mathcal{H}(k)_3$ under the normal form (2), we have:

$$\begin{aligned}\sigma &\xrightarrow{\mathbf{t}^r} \tau_{rs}^\alpha \tau_{rt} \tau_{rt}^\beta \tau_{st}^\gamma [\tau_{rt}, \tau_{st}]^\delta \tau_{rt}^{-1} \\ &= \tau_{rs} \alpha \tau_{rt}^\beta \tau_{st}^\gamma [\tau_{rt}, \tau_{st}]^{\delta+\gamma}; \\ \sigma &\xrightarrow{\mathbf{t}^s} \tau_{rs}^\alpha \tau_{st} \tau_{rt}^\beta \tau_{st}^\gamma [\tau_{rt}, \tau_{st}]^\delta \tau_{st}^{-1} \\ &= \tau_{rs} \alpha \tau_{rt}^\beta \tau_{st}^\gamma [\tau_{rt}, \tau_{st}]^{\delta-\beta}.\end{aligned}$$

To calculate partial conjugations by τ_{rs} and τ_{ts} , which are denoted by \mathbf{s}^r and \mathbf{s}^t , respectively, we need to rewrite σ as follows:

$$\sigma = \tau_{rs}^\alpha \tau_{rt}^\beta \tau_{st}^\gamma [\tau_{rt}, \tau_{st}]^\delta = \tau_{rt}^\beta \tau_{rs}^\alpha \tau_{ts}^\gamma [\tau_{rs}, \tau_{ts}]^{-\delta-\alpha\beta}.$$

Then, we have:

$$\begin{aligned}\sigma &\xrightarrow{\mathbf{s}^r} \tau_{rt}^\beta \tau_{rs} \tau_{rs}^\alpha \tau_{ts}^\gamma [\tau_{rs}, \tau_{ts}]^{-\delta-\alpha\beta} \tau_{rs}^{-1} \\ &= \tau_{rt}^\beta \tau_{rs}^\alpha \tau_{ts}^\gamma [\tau_{rs}, \tau_{ts}]^{-\delta-\alpha\beta+\gamma} \\ &= \tau_{rs}^\alpha \tau_{rt}^\beta \tau_{st}^\gamma [\tau_{rt}, \tau_{st}]^{\delta-\gamma}; \\ \sigma &\xrightarrow{\mathbf{s}^t} \tau_{rt}^\beta \tau_{ts} \tau_{rs}^\alpha \tau_{ts}^\gamma [\tau_{rs}, \tau_{ts}]^{-\delta-\alpha\beta} \tau_{ts}^{-1} \\ &= \tau_{rt}^\beta \tau_{rs}^\alpha \tau_{ts}^\gamma [\tau_{rs}, \tau_{ts}]^{-\delta-\alpha\beta-\alpha} \\ &= \tau_{rs}^\alpha \tau_{rt}^\beta \tau_{st}^\gamma [\tau_{rt}, \tau_{st}]^{\delta+\gamma}.\end{aligned}$$

Similarly, to calculate partial conjugations \mathbf{r}^s and \mathbf{r}^t , we first rewrite σ :

$$\sigma = \tau_{rs}^\alpha \tau_{rt}^\beta \tau_{st}^\gamma [\tau_{rt}, \tau_{st}]^\delta = \tau_{st}^\gamma \tau_{sr}^\alpha \tau_{tr}^\beta [\tau_{sr}, \tau_{tr}]^{\delta-\alpha\gamma+\beta\gamma}.$$

Then, we have

$$\begin{aligned}\sigma &\xrightarrow{\mathbf{r}^s} \tau_{st}^\gamma \tau_{sr} \tau_{sr}^\alpha \tau_{tr}^\beta [\tau_{sr}, \tau_{tr}]^{\delta-\alpha\gamma+\beta\gamma} \tau_{sr}^{-1} \\ &= \tau_{st}^\gamma \tau_{sr}^\alpha \tau_{tr}^\beta [\tau_{sr}, \tau_{tr}]^{\delta-\alpha\gamma+\beta\gamma+\beta} \\ &= \tau_{rs}^\alpha \tau_{rt}^\beta \tau_{st}^\gamma [\tau_{rt}, \tau_{st}]^{\delta+\beta}; \\ \sigma &\xrightarrow{\mathbf{r}^t} \tau_{st}^\gamma \tau_{tr} \tau_{sr}^\alpha \tau_{tr}^\beta [\tau_{sr}, \tau_{tr}]^{\delta-\alpha\gamma+\beta\gamma} \tau_{tr}^{-1} \\ &= \tau_{st}^\gamma \tau_{sr}^\alpha \tau_{tr}^\beta [\tau_{sr}, \tau_{tr}]^{\delta-\alpha\gamma+\beta\gamma-\alpha} \\ &= \tau_{rs}^\alpha \tau_{rt}^\beta \tau_{st}^\gamma [\tau_{rt}, \tau_{st}]^{\delta-\alpha}.\end{aligned}$$

We summarize the previous calculation into the following lemma.

LEMMA 2.6. *The change of $\mu(rst)$ under a partial conjugation is given by*

$$\begin{aligned} \mathbf{t}^r &: \mu(rst) \rightarrow \mu(rst) + l_{st}; \\ \mathbf{t}^s &: \mu(rst) \rightarrow \mu(rst) - l_{rt}; \\ \mathbf{s}^r &: \mu(rst) \rightarrow \mu(rst) - l_{st}; \\ \mathbf{s}^t &: \mu(rst) \rightarrow \mu(rst) + l_{rs}; \\ \mathbf{r}^s &: \mu(rst) \rightarrow \mu(rst) + l_{rt}; \\ \mathbf{r}^t &: \mu(rst) \rightarrow \mu(rst) - l_{rs}. \end{aligned}$$

Furthermore, a partial conjugation by \mathbf{i}^j will not change $\mu(rst)$ if $\{i, j\}$ and $\{r, s, t\}$ have at most one element in common.

For a given string link $\sigma \in \mathcal{H}(k)$, we will think of the whole collection $\{\mu(rst); 1 \leq r < s < t \leq k\}$ as an element in $\mathbf{Z}^{\binom{k}{3}}$. Then the conjugations and partial conjugations act on $\mathbf{Z}^{\binom{k}{3}}$ by translations. We will abuse the notation by using the same symbol to denote both a translation operation and the corresponding translation vector. Thus, a translation operation $T: V \rightarrow V$ on a vector space V is given by $T(v) = v + T$, for all $v \in V$ and a fixed $T \in V$. If T_1 and T_2 are two translations, we have

$$(T_1 \cdot T_2)(v) = v + T_1 + T_2, \quad \text{for all } v \in V.$$

The following two theorems follow directly from Lemmas 2.5 and 2.6.

THEOREM 2.7. *The translation operation on $\mathbf{Z}^{\binom{k}{3}}$ given by the conjugation of τ_{ij} is the same as the composition of the translation operations given by the partial conjugations \mathbf{i}^j and \mathbf{j}^i , i.e. it is equal to $\mathbf{i}^j + \mathbf{j}^i$.*

THEOREM 2.8. *The translation operations \mathbf{i}^j satisfy the following relations:*

$$\sum_{j \neq i} \mathbf{j}^i = 0, \quad \sum_{j \neq i} l_{ij} \mathbf{i}^j = 0$$

for all $i = 1, 2, \dots, k$.

String links are oriented in the sense that each component is given an orientation from the bottom to the top. See Figure 1. Reversing the orientation on each component of a string link defines a bijection

$$\sigma \mapsto \bar{\sigma}: \mathcal{H}(k) \rightarrow \mathcal{H}(k).$$

This bijection is an anti-homomorphism: $\overline{\sigma_1\sigma_2} = \overline{\sigma_2}\overline{\sigma_1}$. This bijection induces an operation on $\mathbf{Z}^{(k)}_{(3)}$.

THEOREM 2.9. *The operation on $\mathbf{Z}^{(k)}_{(3)}$ induced by reversing the orientation of each component of a string link is to change each $\mu(rst)$ to $-\mu(rst)$ followed by the translation operation*

$$\mu(rst) \longrightarrow \mu(rst) - l_{rs}l_{rt} + l_{rs}l_{st} - l_{rt}l_{st}.$$

Proof. Consider the normal form (2) of $\sigma \in \mathcal{H}(k)/\mathcal{H}(k)_3$ in the r, s, t -th components. The normal form for $\overline{\sigma}$ is obtained as follows:

$$\begin{aligned} \overline{\sigma} &= [\tau_{rt}, \tau_{st}]^{-\delta} \tau_{st}^{\gamma} \tau_{rt}^{\beta} \tau_{rs}^{\alpha} \\ &= \tau_{rs}^{\alpha} \tau_{rt}^{\beta} \tau_{st}^{\gamma} [\tau_{rt}, \tau_{st}]^{-\delta - \alpha\beta + \alpha\gamma - \beta\gamma}. \end{aligned}$$

Thus the operation on $\mathbf{Z}^{(k)}_{(3)}$ induced by $\sigma \mapsto \overline{\sigma}$ is given by

$$\mu(rst) \longrightarrow -\mu(rst) - l_{rs}l_{rt} + l_{rs}l_{st} - l_{rt}l_{st}. \quad \square$$

3. CONSTRUCTION OF THE INVARIANT

By Theorems 2.2 and 2.7, we shall look for polynomials in l_{ij} and $\mu(rst)$ invariant under the translation operations on $\{\mu(rst)\} \in \mathbf{Z}^{(k)}_{(3)}$ induced by partial conjugations. There are $k(k-1)$ partial conjugations altogether and their induced translations subject to $2k$ linear equations given in Theorem 2.8. If these equations are linearly independent for generic values of $\{l_{ij}\}$, the sublattice of $\mathbf{Z}^{(k)}_{(3)}$ generated by the translation vectors of the partial conjugations will be of dimension no larger than $k(k-1) - 2k = k^2 - 3k$.

LEMMA 3.1. *For $k > 3$, the $2k$ equations in Theorem 2.8 are linearly independent for generic values of $\{l_{ij}\}$.*

Proof. We write the two sets of equations in Theorem 2.8 as follows:

$$\begin{aligned} \mathbf{1}^i + \mathbf{2}^i + \cdots + \mathbf{j}^i + \cdots + \mathbf{k}^i &= 0, \quad j \neq i; \\ l_{i1}\mathbf{i}^1 + l_{i2}\mathbf{i}^2 + \cdots + l_{ij}\mathbf{i}^j + \cdots + l_{ik}\mathbf{i}^k &= 0, \quad j \neq i, \end{aligned}$$

for each $i = 1, 2, \dots, k$.

For generic values of $\{l_{ij}\}$, using the first $k-1$ equations from the first set of k equations, we can solve for $\mathbf{k}^1, \mathbf{k}^2, \dots, \mathbf{k}^{k-1}$. Similarly, we can solve