

6.2 The Euler class of a group action on the circle

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **47 (2001)**

Heft 3-4: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **14.09.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Now consider the group $\tilde{\Gamma}_g$ defined by the presentation

$$\tilde{\Gamma}_g = \langle z, a_1, b_1, \dots, a_g, b_g \mid a_1 b_1 a_1^{-1} b_1^{-1} \dots a_g b_g a_g^{-1} b_g^{-1} = z, \quad za_i = a_i z, \quad zb_i = b_i z \rangle.$$

The central subgroup A generated by z turns out to be infinite cyclic so that $\tilde{\Gamma}_g$ defines a central extension of Γ_g by \mathbf{Z} , hence an Euler class in $H^2(\Gamma_g, \mathbf{Z})$. It is a fact that $H^2(\Gamma_g, \mathbf{Z})$ is isomorphic with \mathbf{Z} and that the element that we have just constructed is a generator of this cohomology group. We shall not prove this here but we note that this is related to the fact that a closed oriented surface of genus $g \geq 1$ has a contractible universal cover and that the cohomology of Γ_g can therefore be identified with the cohomology of the compact oriented surface of genus g (see [11] for more details).

6.2 THE EULER CLASS OF A GROUP ACTION ON THE CIRCLE

We have already met a central extension related to groups of homeomorphisms

$$0 \longrightarrow \mathbf{Z} \longrightarrow \widetilde{\text{Homeo}}_+(\mathbf{S}^1) \xrightarrow{p} \text{Homeo}_+(\mathbf{S}^1) \longrightarrow 1.$$

The cohomology group $H^2(\text{Homeo}_+(\mathbf{S}^1), \mathbf{Z})$ has been computed. It is isomorphic to \mathbf{Z} and a generator is the Euler class of this central extension [50].

Consider now a homomorphism ϕ from some group Γ to $\text{Homeo}_+(\mathbf{S}^1)$. Then, we can pull back the previous extension by ϕ . In other words, we consider the set of $(\gamma, \tilde{f}) \in \Gamma \times \widetilde{\text{Homeo}}_+(\mathbf{S}^1)$ such that $\phi(\gamma) = p(\tilde{f})$. This is a group $\tilde{\Gamma}$ equipped with a canonical projection onto Γ whose kernel is isomorphic to \mathbf{Z} , *i.e.* $\tilde{\Gamma}$ is a central extension of Γ by \mathbf{Z} . In case ϕ is injective, $\tilde{\Gamma}$ is just the pre-image of $\phi(\Gamma)$ under p , which is the group of lifts of $\phi(\Gamma)$. The Euler class of this central extension of Γ is called *the Euler class of the homomorphism ϕ* and denoted by $eu(\phi) \in H^2(\Gamma, \mathbf{Z})$. It is obviously a dynamical invariant in the sense that two conjugate homomorphisms ϕ_1 and ϕ_2 have the same Euler class in $H^2(\Gamma, \mathbf{Z})$. Note that it follows from the definition that $eu(\phi)$ is zero if and only if the homomorphism ϕ lifts to a homomorphism $\tilde{\phi}: \Gamma \rightarrow \widetilde{\text{Homeo}}_+(\mathbf{S}^1)$ such that $\phi = p \circ \tilde{\phi}$.

A few examples are in order. In the case of a single homeomorphism, *i.e.* when $\Gamma = \mathbf{Z}$, we saw that $H^2(\mathbf{Z}, \mathbf{Z}) = 0$. Hence the Euler class vanishes and our new invariant is very poor indeed: in particular, it does not detect the rotation number. A similar phenomenon occurs when Γ is free.

If Γ_g is the fundamental group of a closed oriented surface of genus $g \geq 1$, we know that $H^2(\Gamma_g, \mathbf{Z})$ is isomorphic to \mathbf{Z} so that the Euler class

$eu(\phi)$ in this case is an integer. In [51], Milnor gives an algorithm to compute this number. With the same notation as above, for each $1 \leq i \leq g$, choose lifts \tilde{a}_i and \tilde{b}_i of $\phi(a_i)$ and $\phi(b_i)$. Now compute the product of commutators $\tilde{a}_1 \tilde{b}_1 \tilde{a}_1^{-1} \tilde{b}_1^{-1} \dots \tilde{a}_g \tilde{b}_g \tilde{a}_g^{-1} \tilde{b}_g^{-1}$. Since this homeomorphism is a lift of the identity, it is an integral translation. This amplitude of this translation does not depend on the choices made and is the Euler number $eu(\phi)$.

As an explicit example, also computed by Milnor, recall that any closed orientable surface of genus $g > 1$ can be endowed with a riemannian metric of constant negative curvature. Recall also that the Poincaré upper half space \mathcal{H} can be equipped with a metric of curvature -1 whose group of orientation preserving isometries is precisely $\text{PSL}(2, \mathbf{R})$. Moreover, any complete simply connected riemannian surface of curvature -1 is isometric to \mathcal{H} . Hence there are embeddings ϕ of the fundamental group Γ_g of a closed oriented surface of genus $g > 1$ in $\text{PSL}(2, \mathbf{R})$ such that the corresponding action of Γ_g on \mathcal{H} is free, proper and cocompact. Since we know that $\text{PSL}(2, \mathbf{R})$ is a subgroup of $\text{Homeo}_+(\mathbf{S}^1)$, we can compute the corresponding Euler number $eu(\phi)$. The result of the computation is $2g - 2$. Note that each element of $\phi(\Gamma_g)$ is hyperbolic since the action is free and cocompact so that the rotation number of every element of $\phi(\Gamma_g)$ is 0. So we are in a situation in which the topological invariant $eu(\phi)$ is not 0 but the rotation number invariants are trivial; a situation different from the case where $\Gamma = \mathbf{Z}$.

6.3 BOUNDED COHOMOLOGY AND THE MILNOR-WOOD INEQUALITY

It was observed very early that the Euler class of a homomorphism $\phi: \Gamma \rightarrow \text{Homeo}_+(\mathbf{S}^1)$ cannot be arbitrary. Milnor and Wood proved the following [51, 71].

THEOREM 6.1 (Milnor-Wood). *Let Γ_g be the fundamental group of a closed oriented surface of genus $g \geq 1$ and $\phi: \Gamma_g \rightarrow \text{Homeo}_+(\mathbf{S}^1)$ be any homomorphism. Then the Euler number satisfies $|eu(\phi)| \leq 2g - 2$.*

Proof. We shall not give a complete proof since this result will follow from later considerations but we prove a weaker version. Keeping the previous notation, we know that $eu(\phi)$ is the translation number of the homeomorphism $\tilde{a}_1 \tilde{b}_1 \tilde{a}_1^{-1} \tilde{b}_1^{-1} \dots \tilde{a}_g \tilde{b}_g \tilde{a}_g^{-1} \tilde{b}_g^{-1}$. We also know that the translation number function τ is a quasi-homomorphism, i.e. there is some inequality of the form $|\tau(\tilde{f}_1 \tilde{f}_2) - \tau(\tilde{f}_1) - \tau(\tilde{f}_2)| \leq D$ for some D . We also know that $\tau(\tilde{f}^{-1}) = -\tau(\tilde{f})$. So, if we evaluate τ on this element, we get a bound of the form $|eu(\phi)| \leq (4g - 1)D$. This is not quite the bound given in the