

5. FURTHER PROPERTIES OF THE CONES

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **47 (2001)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **14.09.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

LEMMA 4.2. For $m \geq 2k + 1$, the group $\pi_m(BU(k))$ is finite.

Proof. We fix $m \geq 3$. The fibration $BU(k-1) \rightarrow BU(k)$, with fibre S^{2k-1} , yields the following long exact sequence in homotopy:

$$\dots \rightarrow \pi_m(S^{2k-1}) \rightarrow \pi_m(BU(k-1)) \rightarrow \pi_m(BU(k)) \rightarrow \pi_{m-1}(S^{2k-1}) \rightarrow \dots$$

By Serre [Serre], $\pi_j(S^{2k-1})$ is finite for $j \neq 2k-1$, and we can conclude by induction over k (with $k \geq 1$ and $2k+1 \leq m$), since when $k=1$, one has $\pi_m(BU(1)) = \pi_{m-1}(U(1)) = 0$ for $m \geq 3$. \square

From this, we now infer that the image of $(i_k)_*$ is zero for $k < n$. This implies that $\text{g-dim}(lx) = n$ when $l \neq 0$, and concludes the second proof.

REMARK 4.3.

i) Since we were motivated by Elliott's classification of unital C^* -algebras of type AF by means of their K -theory, their positive cone and the K -theory class [1] of the unit (see [Bla1]), it is important to single out the fact that the positive cone of S^{2n} and that of S^{2m} are non-isomorphic as monoids if n is different from m . (There is no need here to distinguish the K -theory class 1 of the trivial one-dimensional bundle.) Let us provide a short proof of this claim. For $n \geq 1$, let M_n denote the positive cone of S^{2n} (identified as above with a sub-monoid of \mathbf{Z}^2 , in order to designate its elements). The abelian monoid M_n has a minimal set A_n of generators, in other words a generating set (as a monoid) that is contained in any other generating set, namely

$$A_n = \{(0, 1)\} \cup \{(k, n) \mid k \in \mathbf{Z} \setminus \{0\}\}.$$

Now, consider the function $\sigma: A_n \rightarrow \{2, 3, \dots\}$ defined, for $x \in A_n$, by

$$\sigma(x) := \min \{l \geq 2 \mid lx \text{ decomposes as a sum of elements of } A_n \setminus \{x\}\}.$$

Clearly, such an l exists for any $x \in A_n$ and $\sigma(A_n) = \{2, 2n\}$. Since A_n and σ are isomorphism invariants of M_n , this proves our claim.

ii) For odd-dimensional spheres the positive cone is "trivial"; in other words, $K(S^{2n-1}) = \mathbf{Z}$ and $K_+(S^{2n-1}) = \mathbf{N}$.

5. FURTHER PROPERTIES OF THE CONES

We now investigate naturality properties and behaviour under products of the positive cone, the γ -cone and the c -cone.

The following result is obvious.

PROPOSITION 5.1. *Let $f: X \rightarrow Y$ be a map between connected finite CW-complexes. Let $f^*: K(Y) \rightarrow K(X)$ be the λ -homomorphism induced by f . Then, for any $y \in \tilde{K}(Y)$, one has*

$$\begin{aligned} \text{g-dim}(f^*(y)) &\leq \text{g-dim}(y) \\ \gamma\text{-dim}(f^*(y)) &\leq \gamma\text{-dim}(y) \\ \text{c-dim}(f^*(y)) &\leq \text{c-dim}(y), \end{aligned}$$

and in particular,

$$\begin{aligned} f^*(K_+(Y)) &\subseteq K_+(X) \\ f^*(K_\gamma(Y)) &\subseteq K_\gamma(X) \\ f^*(K_c(Y)) &\subseteq K_c(X). \end{aligned}$$

Furthermore, if f^* is an isomorphism, then

$$f^*(K_\gamma(Y)) = K_\gamma(X).$$

For the next corollary we need a new definition.

DEFINITION 5.2. Let X and Y be two connected finite CW-complexes. A map $f: X \rightarrow Y$ is called a K^0 -equivalence (or K -equivalence for short) if there exists a map $g: Y \rightarrow X$ such that, on the level of the K^0 -groups,

$$f^* \circ g^* = \text{Id}_{K^0(X)} \quad \text{and} \quad g^* \circ f^* = \text{Id}_{K^0(Y)}.$$

Note that a K -equivalence is *not* necessarily a homotopy equivalence: there are homotopically non-trivial (i.e. non-contractible) finite CW-complexes X for which $\tilde{K}(X) = 0 = \tilde{K}(pt)$; see example iii) below.

PROPOSITION 5.3. *If $f: X \rightarrow Y$ is a K -equivalence, then f induces the following isomorphisms of semigroups:*

$$K_+(Y) \xrightarrow{f^*} K_+(X) \quad \text{and} \quad K_\gamma(Y) \xrightarrow{f^*} K_\gamma(X).$$

Proof. Applying Proposition 5.1 twice, we get (in the notations of Definition 5.2)

$$K_+(X) = f^* \circ g^*(K_+(X)) \subseteq f^*(K_+(Y)) \subseteq K_+(X).$$

This establishes the first isomorphism, whereas the second one is obvious. \square

The following result is more technical to state.

COROLLARY 5.4. *Let X and Y be two connected finite CW-complexes. Assume that $K^1(X) = 0$ and that $\tilde{K}^0(Y) = 0$. Then the projection $p: X \times Y \rightarrow X$ induces isomorphisms*

$$K_+(X) \stackrel{p^*}{\cong} K_+(X \times Y) \quad \text{and} \quad K_\gamma(X) \stackrel{p^*}{\cong} K_\gamma(X \times Y).$$

Proof. Invoking the Künneth theorem for K -theory, our hypotheses imply that $p^*: K^0(X) \rightarrow K^0(X \times Y)$ is an isomorphism with inverse i^* , where i is the inclusion of X in $X \times Y$. Consequently, p^* is a K -equivalence. \square

The following is a useful result.

PROPOSITION 5.5. *Let X and Y be connected finite CW-complexes. Assume that the positive cone and the γ -cone of Y coincide, and let $f: X \rightarrow Y$ be a map inducing an isomorphism $f^*: K(Y) \rightarrow K(X)$. Then f induces an isomorphism of positive cones, and the γ -cone of X coincides with the positive cone:*

$$K_+(Y) \stackrel{f^*}{\cong} K_+(X) = K_\gamma(X).$$

Proof. By Proposition 5.1 we have $f^*(K_+(Y)) = f^*(K_\gamma(Y)) = K_\gamma(X)$ and $f^*(K_+(Y)) \subseteq K_+(X)$, hence $K_\gamma(X) \subseteq K_+(X)$. We conclude with iii) of Proposition 3.2. \square

EXAMPLES.

i) Let X be a connected finite CW-complex of dimension ≤ 3 . Since for suitable CW-decompositions, one has $BU(1)^{[3]} = BU^{[3]}$ and since $BU(1) = CP^\infty = K(\mathbf{Z}, 2)$, any $x \in \tilde{K}(X) = [X, BU]$ lifts to a class in $[X, BU(1)]$, giving an isomorphism $\tilde{K}(X) \cong H^2(X; \mathbf{Z})$ mapping x to $c_1(x)$. It follows that the positive cone coincides with the c -cone and is given by

$$K_+(X) = \mathbf{N} \times \{0\} \cup \mathbf{N}^* \times \tilde{K}(X) \subset \mathbf{Z} \times \tilde{K}(X).$$

ii) Example i) applies to a closed oriented surface Σ_g of genus g . Since it is torsion-free, its positive cone coincides with its c -cone and with its γ -cone. Moreover, let $f: \Sigma_g \rightarrow S^2$ be a map of degree 1 (it exists, since both the 2-sphere and Σ_g are quotients of the square $[0, 1]^2$). Then f not only induces an isomorphism in K -theory, but also an isomorphism of positive cones, as follows from Proposition 5.1.

iii) Let X and Y be the Moore spaces $M(\mathbf{Z}/3, 2q+11) = S^{2q+11} \cup_3 e^{2q+12}$ and $M(\mathbf{Z}/3, 2q-1) = S^{2q-1} \cup_3 e^{2q}$ respectively. In [Adams], Adams shows that for q large enough, there exists a map $A: X = \Sigma^{12}Y \longrightarrow Y$ such that the induced map $A^*: \tilde{K}(Y) \longrightarrow \tilde{K}(X)$ is an isomorphism (take $p = m = 3$, $f = 1$ and $r = 6$ in Theorem 1.7 and in Lemmas 12.4 and 12.5 of [Adams]). Therefore, A is a K -isomorphism between simply connected finite CW-complexes, but it is *not* a homotopy equivalence. The mapping cone C_A is a non-contractible finite CW-complex with $\tilde{K}(C_A) = 0$. (It is non-contractible because its homology is non-trivial.)

iv) In [GrMo], pp. 203-206, a CW-complex $X = (S^1 \vee S^2) \cup e^3$ is defined, with the property that the inclusion $i: S^1 = X^{[1]} \hookrightarrow X$ of the 1-skeleton induces an isomorphism in integral homology (and on the level on fundamental groups); however, i is *not* a homotopy equivalence since $\pi_2(X) \neq 0$. Consequently, by the universal coefficient theorem (see Corollary V.7.2 in [Bred]), i induces an isomorphism in integral cohomology, and, by a direct application of the Atiyah-Hirzebruch spectral sequence, also in K -theory. In particular, i is a K -equivalence, but *not* an equivalence. (As C_A in the preceding example, the quotient space $X/X^{[1]}$ has vanishing \tilde{K} , however it is the closed 3-ball and is therefore contractible.)

Let us finally mention that in [Matt], the positive cone, the c -cone and the γ -cone are also studied from the rational point of view, and rational K -theory is considered.

6. THE CONES OF THE PRODUCTS $S^n \times S^{2m-1}$

In this section, we will compute the cones for the products $S^{2n} \times S^{2m-1}$ and $S^{2n-1} \times S^{2m-1}$.

We begin with $S^{2n} \times S^{2m-1}$. Since $\tilde{K}(S^{2m-1}) = 0$ and $K^1(S^{2n}) = 0$, the answer immediately follows from Proposition 5.5.

THEOREM 6.1. *The projection $p: S^{2n} \times S^{2m-1} \longrightarrow S^{2n}$ induces an isomorphism of positive cones, and, for $S^{2n} \times S^{2m-1}$, the γ -cone and the c -cone coincide with the positive cone:*

$$K_+(S^{2n}) \stackrel{p^*}{\cong} K_+(S^{2n} \times S^{2m-1}) = K_\gamma(S^{2n} \times S^{2m-1}).$$