

4.2 Proof of Theorem 2

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **48 (2002)**

Heft 3-4: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **10.08.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

4. NILPOTENT LIE ALGEBRAS WITH INFINITELY MANY
NON-ISOMORPHIC RATIONAL FORMS

In this section we propose a construction which can provide a series of nilpotent Lie algebras with infinitely many isomorphism classes of rational forms.

4.1 BASIC LEMMA

Let

$$\mathfrak{h} = \bigoplus_{i=1}^c \mathfrak{h}_i = \mathfrak{h}(\mathbf{Q})$$

be a graded Lie algebra over \mathbf{Q} generated by \mathfrak{h}_1 . Let \mathbf{K} be a number field, $\dim_{\mathbf{Q}} \mathbf{K} = d$, of type (s, t) , that is, there are s real and $2t$ complex embeddings of \mathbf{K} in \mathbf{C} ($d = s + 2t$) whence there exists an isomorphism of \mathbf{R} -algebras

$$\mathbf{K} \otimes_{\mathbf{Q}} \mathbf{R} \cong \bigoplus_{k=1}^s \mathbf{R} \oplus \bigoplus_{l=1}^t \mathbf{C}.$$

More generally one can take a finite-dimensional commutative associative algebra \mathbf{A} over \mathbf{Q} instead of \mathbf{K} . We consider the Lie algebra $\mathfrak{h}(\mathbf{K}) = \mathfrak{h} \otimes_{\mathbf{Q}} \mathbf{K}$ as a Lie algebra over \mathbf{Q} . This algebra has two important properties. Firstly,

$$\mathfrak{h}(\mathbf{K}) \otimes_{\mathbf{Q}} \mathbf{R} \cong (\mathfrak{h} \otimes_{\mathbf{Q}} \mathbf{K}) \otimes_{\mathbf{Q}} \mathbf{R} \cong \mathfrak{h} \otimes_{\mathbf{Q}} (\mathbf{K} \otimes_{\mathbf{Q}} \mathbf{R}) \cong \bigoplus_{k=1}^s \mathfrak{h}(\mathbf{R}) \oplus \bigoplus_{l=1}^t \mathfrak{h}(\mathbf{C}),$$

i.e., $\mathfrak{h}(\mathbf{K})$ is a \mathbf{Q} -form of the last Lie algebra for any number field \mathbf{K} of type (s, t) . Secondly, there is an embedding $R: \mathbf{K}^* \rightarrow \text{Aut}_{\mathbf{Q}}(\mathfrak{h}(\mathbf{K}))$ of the multiplicative group \mathbf{K}^* such that $R(k)(h_i \otimes k_1) = h_i \otimes \tilde{k}k^i$ where $h_i \in \mathfrak{h}_i$ is homogenous of degree i . The following lemma is straightforward.

LEMMA 4.1. *Let $\mathbf{K} \neq \mathbf{K}'$ be two distinct number fields of the same type. If there is no injection of \mathbf{K}^* into $\text{Aut}_{\mathbf{Q}}(\mathfrak{h}(\mathbf{K}'))$ then two \mathbf{Q} -forms $\mathfrak{h}(\mathbf{K})$ and $\mathfrak{h}(\mathbf{K}')$ are not isomorphic.*

4.2 PROOF OF THEOREM 2

We start with the class of nilpotence $c = 2$. Let $\mathbf{K} = \mathbf{Q}(\sqrt{m})$ and $\mathbf{K}' = \mathbf{Q}(\sqrt{n})$, where $m \neq n$ are two positive (resp. negative) square-free integers. Consider the automorphism $A = R(\sqrt{m})$ of $\mathfrak{h}(\mathbf{K}) = \mathfrak{f}_2(p, \mathbf{K})$. One immediately checks that

- 1) A^2 acts on $\mathfrak{h}(\mathbf{K})/[\mathfrak{h}(\mathbf{K}), \mathfrak{h}(\mathbf{K})]$ as $m \cdot Id$;
- 2) the restriction

$$A|_{[\mathfrak{h}(\mathbf{K}), \mathfrak{h}(\mathbf{K})]} = m \cdot Id.$$

By Lemma 4.1 we must prove that there is no such automorphism for $\mathfrak{h}(\mathbf{K}') = \mathfrak{h}(\mathbf{Q}(\sqrt{n}))$. We choose the following basis of $\mathfrak{h}(\mathbf{K}')$ over \mathbf{Q} :

$$X_i = x_i \otimes 1, \quad Y_i = x_i \otimes \sqrt{n}, \quad C_{ij} = c_{ij} \otimes 1, \quad Z_{ij} = c_{ij} \otimes \sqrt{n},$$

$x_1, \dots, x_p, c_{ij} = [x_i, x_j]$ being the standard basis of $\mathfrak{f}_2(p, \mathbf{Q})$.

Suppose that there exists an automorphism A' with two above properties. First of all, let us show that $[X_i, A'(X_i)] = 0$. On the one hand,

$$A'[X_i, A'(X_i)] = [A'(X_i), mX_i] = -m[X_i, A'(X_i)].$$

On the other hand,

$$A'[X_i, A'(X_i)] = m[X_i, A'(X_i)].$$

Since the centralizer of X_i is generated modulo the centre by X_i, Y_i it follows that

$$A'(X_i) = p_i X_i + q_i Y_i + \varepsilon = x_i \otimes (p_i + q_i \sqrt{n}) + \varepsilon, \quad q_i \neq 0.$$

Here ε stands for a central element which plays no role below.

Consider now $[X_i, A'(X_j)] = c_{ij} \otimes (p_j + q_j \sqrt{n})$. On the one hand,

$$A'[X_i, A'(X_j)] = [A'(X_i), mX_j] = c_{ij} \otimes m(p_i + q_i \sqrt{n}).$$

On the other hand,

$$A'[X_i, A'(X_i)] = m[X_i, A'(X_j)] = c_{ij} \otimes m(p_j + q_j \sqrt{n}),$$

whence

$$p_i + q_i \sqrt{n} = p_j + q_j \sqrt{n} = p + q \sqrt{n} \notin \mathbf{Q} \quad \forall i, j.$$

Finally, we apply A' to $[A'(X_i), A'(X_j)] = c_{ij} \otimes (p + q \sqrt{n})^2$. On the one hand,

$$A'[A'(X_i), A'(X_j)] = [mX_i, mX_j] = c_{ij} \otimes m^2.$$

On the other hand,

$$A'[A'(X_i), A'(X_i)] = m[A'(X_i), A'(X_j)] = c_{ij} \otimes m(p + q \sqrt{n})^2.$$

It follows that $m = (p + q \sqrt{n})^2$. We have obtained a contradiction since $q \neq 0$. Thus, there are infinitely many non-isomorphic rational forms of $\mathfrak{f}_2(p, \mathbf{R}) \oplus \mathfrak{f}_2(p, \mathbf{R})$ and of $\mathfrak{f}_2(p, \mathbf{C})$.

More generally let $\mathfrak{g} = \mathfrak{f}_c(p, \mathbf{R})$ be a free nilpotent Lie algebra of class $c \geq 3$ on p generators. Then $\mathfrak{g} \oplus \mathfrak{g}$ and $\mathfrak{g} \otimes_{\mathbf{R}} \mathbf{C} = \mathfrak{f}_c(p, \mathbf{C})$ (as a Lie algebra over \mathbf{R}) also have infinitely many non-isomorphic rational forms. Consider the automorphism A as above and note that it respects the descending central series. Any isomorphism between $\mathfrak{f}_c(p, \mathbf{K})$ and $\mathfrak{f}_c(p, \mathbf{K}')$ must respect it, too. Then we can take the free nilpotent quotients of class 2 of both algebras and obtain a contradiction just like in the first part of the proof. \square

Thus, the case of a free nilpotent Lie algebra $f_c(p, \mathbf{C})$ (as a Lie algebra over \mathbf{R}) on p generators differs from the case 2.2.

REMARK. All rational forms of $f_2(2, \mathbf{C}) = \mathfrak{hei}_3(\mathbf{C})$ and $f_2(2, \mathbf{R}) \oplus f_2(2, \mathbf{R}) = \mathfrak{hei}_3(\mathbf{R}) \oplus \mathfrak{hei}_3(\mathbf{R})$ are listed in Theorem 3.

COROLLARY 4.2. *There are infinitely many non-commensurable (in any sense) lattices in the Lie groups of type $F_c(p, \mathbf{R}) \times F_c(p, \mathbf{R})$ where $F_c(p, \mathbf{R})$ is the free nilpotent Lie group on p free generators.*

4.3 CLASSIFICATION OF RATIONAL FORMS FOR SOME 6-DIMENSIONAL LIE ALGEBRAS

Let m be a rational number and $A_m = \mathbf{Q}[x]/(x^2 - m)$. A_m is a 2-dimensional commutative algebra over \mathbf{Q} which depends only on m modulo square factors. Thus there are four types of A_m :

- 1) if $m = 1$ then $A_m \cong \mathbf{Q} \oplus \mathbf{Q}$;
- 2) if $m > 1$ is a positive square-free integer then $A_m \cong \mathbf{Q}(\sqrt{m})$ is a real quadratic field over \mathbf{Q} ;
- 3) if $m = 0$ then A_0 is the algebra of dual numbers over \mathbf{Q} ;
- 4) if m is a negative square-free integer then $A_m \cong \mathbf{Q}(\sqrt{m})$ is an imaginary quadratic field over \mathbf{Q} .

Let $\mathfrak{hei}_3(A_m)$ be a Heisenberg algebra over A_m considered over \mathbf{Q} . Then $\mathfrak{hei}_3(A_m)$ is a rational form of either $\mathfrak{hei}_3(\mathbf{R}) \oplus \mathfrak{hei}_3(\mathbf{R})$, or $\mathfrak{hei}_3(\mathbf{R}[x]/(x^2))$, or $\mathfrak{hei}_3(\mathbf{C})$. More precisely,

THEOREM 3. *Let \mathfrak{h} be a 6-dimensional nilpotent Lie algebra of class 2 over \mathbf{Q} . Suppose that $[\mathfrak{h}, \mathfrak{h}]$ coincides with the 2-dimensional centre of \mathfrak{h} . Then $\mathfrak{h} \cong \mathfrak{hei}_3(A_m)$ for some $m \in \mathbf{Q}$ as above.*

Moreover,

- 1) $\mathfrak{h} \otimes_{\mathbf{Q}} \mathbf{R} \cong \mathfrak{hei}_3(\mathbf{R}) \oplus \mathfrak{hei}_3(\mathbf{R}) = \mathfrak{g}_+$ iff $m > 0$,
- 2) $\mathfrak{h} \otimes_{\mathbf{Q}} \mathbf{R} \cong \mathfrak{hei}_3(\mathbf{R}[x]/(x^2)) = \mathfrak{g}_0$ iff $m = 0$,
- 3) $\mathfrak{h} \otimes_{\mathbf{Q}} \mathbf{R} \cong \mathfrak{hei}_3(\mathbf{C}) = \mathfrak{g}_-$ iff $m < 0$,

and up to isomorphism there are no more rational forms for \mathfrak{g}_- , \mathfrak{g}_0 , \mathfrak{g}_+ . The Lie algebras $\mathfrak{hei}_3(A_m)$ and $\mathfrak{hei}_3(A_n)$ are isomorphic over \mathbf{Q} if and only if A_m and A_n are isomorphic.