

# 1. Introduction

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **48 (2002)**

Heft 3-4: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **14.09.2024**

## **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

ON THE CLASSIFICATION OF CERTAIN PIECEWISE LINEAR  
AND DIFFERENTIABLE MANIFOLDS IN DIMENSION EIGHT  
AND AUTOMORPHISMS OF  $\#_{i=1}^b(S^2 \times S^5)$

by Alexander SCHMITT

ABSTRACT. In this paper, we will be concerned with the explicit classification of closed, oriented, simply-connected spin manifolds in dimension eight with vanishing cohomology in the odd dimensions. The study of such manifolds was begun by Stefan Müller. In order to understand the structure of these manifolds, we will analyze their minimal handle presentations and describe explicitly to what extent these handle presentations are determined by the cohomology ring and the characteristic classes. It turns out that the cohomology ring and the characteristic classes do not suffice to reconstruct a manifold of the above type completely. In fact, the group  $\text{Aut}_0(\#_{i=1}^b(S^2 \times S^5)) / \text{Aut}_0(\#_{i=1}^b(S^2 \times D^6))$  of automorphisms of  $\#_{i=1}^b(S^2 \times S^5)$  which induce the identity on cohomology modulo those which extend to  $\#_{i=1}^b(S^2 \times D^6)$  acts on the set of oriented homeomorphism classes of manifolds with fixed cohomology ring and characteristic classes, and we will be also concerned with describing this group and some facts about the above action.

1. INTRODUCTION

The classification of topological manifolds up to homeomorphism is an extremely interesting and important problem. Let us restrict our attention to the case of closed (i.e., compact without boundary) and oriented simply connected manifolds. As a general classification scheme, surgery theory [1] solves this problem for manifolds within a given homotopy type, e.g., that of a sphere. Another approach to the classification “up to finite indeterminacy”, using rational homotopy theory, is due to Sullivan [34]. Nevertheless, there are only a few explicit results which characterize the oriented homeomorphism type of a manifold in terms of easily computable invariants. They usually require many simplifying assumptions, such as high connectedness [36]. In this paper, we will

consider *even cohomology manifolds* (or *E-manifolds*, for short) in dimension eight, by which we mean closed, oriented, simply connected, piecewise linear or smooth manifolds all of whose odd-dimensional homology groups with integer coefficients vanish. The universal coefficient theorem implies that all homology groups of an E-manifold are without torsion. Moreover, since  $H^3(X, \mathbf{Z}_2) = 0$  for an E-manifold, two E-manifolds of dimension at least 6 are homeomorphic (as topological manifolds) if and only if they are piecewise linearly homeomorphic [16].

Though the class of E-manifolds is fairly restricted, it still contains many interesting examples from various areas of mathematics, such as the piecewise linear manifolds underlying the toric manifolds from Algebraic Geometry [4] or the polygon spaces [12], to mention a few. So far, E-manifolds have been classified up to dimension 6. Of course, in dimension 2 there is only  $S^2$ , in dimension 4, there is the famous classification result of Freedman various interesting aspects of which are discussed in [15], and finally in dimension 6, the classification was achieved by Wall [37] and Jupp [14]. Various applications of the latter result to Algebraic Geometry are surveyed in [26]. Finally, we refer to [2], [3], and [29] for the determination of projective algebraic structures on certain 6- and 8-dimensional E-manifolds.

ACKNOWLEDGEMENTS. My thanks go to J.-C. Hausmann for his interest in this work and for pointing out to me that an E-manifold of dimension eight is actually not determined by its classical invariants.

## 2. STATEMENT OF THE RESULT

We now discuss the main result of this note, namely the classification of E-manifolds of dimension 8 with vanishing second Stiefel-Whitney class in the form of an exact sequence of pointed sets. This result was motivated by the work [24]. In order to state it in a more elegant form, we will work with based manifolds. By a *based piecewise linear (smooth) E-manifold*, we mean a triple  $(X, \underline{x}, \underline{y})$ , consisting of a piecewise linear (smooth) E-manifold  $X$  and bases  $\underline{x} = (x_1, \dots, x_{b_2(M)})$  for  $H^2(X, \mathbf{Z})$  and  $\underline{y} = (y_1, \dots, y_{b_4(M)})$  for  $H^4(X, \mathbf{Z})$ . Recall that by definition E-manifolds are oriented, so that the above data specify a basis for  $H^*(X, \mathbf{Z})$ , such that the bases for  $H^i(X, \mathbf{Z})$  and  $H^{8-i}(X, \mathbf{Z})$  are dual to each other with respect to the cup product. An *isomorphism between piecewise linear (smooth) based E-manifolds*  $(X, \underline{x}, \underline{y})$  and  $(X', \underline{x}', \underline{y}')$  is an orientation preserving piecewise linear (smooth) isomorphism  $f: X \rightarrow X'$  with  $f^*(\underline{x}') = \underline{x}$