

**Zeitschrift:** L'Enseignement Mathématique  
**Band:** 48 (2002)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** TRIPLE RATIO ON THE UNITARY STIEFEL MANIFOLD  
**Kapitel:** 4. The triple ratio on S  
**Autor:** Clerc, Jean-Louis  
**DOI:** <https://doi.org/10.5169/seals-66067>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

**Download PDF:** 16.10.2024

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

4. THE TRIPLE RATIO ON  $S$ 

We return to the notation introduced in Sections 1 and 2.

For  $z_1, z_2, z_3 \in \text{Mat}(p \times q, \mathbf{C})$  define, whenever it makes sense, the element  $T(z_1, z_2, z_3) \in \text{GL}(q, \mathbf{C})$  by the following formula

$$(24) \quad \begin{aligned} T(z_1, z_2, z_3) &= k(z_1, z_2) k(z_3, z_2)^{-1} k(z_3, z_1) \\ &= (\mathbf{1}_q - z_2^* z_1)^{-1} (\mathbf{1}_q - z_2^* z_3) (\mathbf{1}_q - z_1^* z_3)^{-1}. \end{aligned}$$

It satisfies the following transformation law

$$(25) \quad T(g(z_1), g(z_2), g(z_3)) = j(g, z_1) T(z_1, z_2, z_3) j(g, z_1)^*$$

for  $g \in G$ . In particular, we see that  $T(\sigma_1, \sigma_2, \sigma_3)$  is well defined on  $S_{\top}^3$  and that the  $\text{GL}(q, \mathbf{C})$ -orbit of  $T(\sigma_1, \sigma_2, \sigma_3)$  is constant along any  $G$ -orbit in  $S_{\top}^3$ .

LEMMA 4.1. Let  $\sigma = \begin{pmatrix} \sigma_p \\ \sigma' \end{pmatrix} \in S$ , tranverse to  $ie$  and  $-ie$ . Then

$$(26) \quad T(ie, -ie, \sigma) = \frac{1}{2i} (i\mathbf{1}_q + \sigma_q) (\mathbf{1}_q + i\sigma_q)^{-1}.$$

*Proof.* This is an easy computation.

PROPOSITION 4.2. Let  $(\sigma_1, \sigma_2, \sigma_3) \in S_{\top}^3$ . Then

$$2i T(\sigma_1, \sigma_2, \sigma_3) \in \tilde{T}_q^{(r)}.$$

*Proof.* Let us first assume  $\sigma_1 = ie, \sigma_2 = -ie, \sigma_3 = \sigma$ . Except for the factor  $\frac{1}{2i}$ , a comparison with (9) shows that  $T(ie, -ie, \sigma)$  is the first term of the Cayley transform of  $\sigma$ . More precisely, let  $c(\sigma) = \xi = \begin{pmatrix} \xi_q \\ \xi' \end{pmatrix}$ . Then we may rewrite (26) as

$$T(ie, -ie, \sigma) = \frac{1}{2i} \xi_q.$$

Now  $\xi$  belongs to  ${}^cS$ , and hence  $\frac{1}{2i}(\xi_q - \xi_q^*) = \xi'^* \xi'$ . But  $\text{rank}(\xi') \leq r$ , so  $\text{rank}(\xi'^* \xi') \leq r$  and hence  $\xi_q$  belongs to  $\tilde{T}_q^{(r)}$ . Now the transformation law (25) for the triple ratio implies that for any  $(\sigma_1, \sigma_2, \sigma_3) \in S_{\top}^3$ ,  $2i T(\sigma_1, \sigma_2, \sigma_3)$  belongs to  $\tilde{T}_q^{(r)}$ .  $\square$

**THEOREM 4.3.** *Let  $(\sigma_1, \sigma_2, \sigma_3)$  and  $(\tau_1, \tau_2, \tau_3)$  belong to  $S^3_\top$ . They belong to the same  $G$ -orbit if and only if  $T(\sigma_1, \sigma_2, \sigma_3)$  and  $T(\tau_1, \tau_2, \tau_3)$  belong to the same  $GL(q, \mathbf{C})$ -orbit.*

*Proof.* One way is obvious from the transformation law (25) for the triple ratio. For the converse, we assume (as we may) that  $\sigma_1 = \tau_1 = ie$  and  $\sigma_2 = \tau_2 = -ie$ , and set for simplicity  $\sigma = \sigma_3$  and  $\tau = \tau_3$ . Then the assumption implies that  $(i\mathbf{1}_q + \sigma_q)(\mathbf{1}_q - i\sigma_q)^{-1}$  and  $(i\mathbf{1}_q + \tau_q)(\mathbf{1}_q - i\tau_q)^{-1}$  are in the same  $GL(q, \mathbf{C})$ -orbit. By Lemma 2.3,  $c(\sigma)$  and  $c(\tau)$  are in the same  ${}^cL$ -orbit. So  $\sigma$  and  $\tau$  are in the same  $L$ -orbit.  $\square$

Now to give a description of the invariant in terms of Theorem 3.13, we need to define the analog of the function  $\arg \det$ . For  $z_1 \in D$  and  $z_2 \in \bar{D}$ , the function  $k(z_1, z_2) = (\mathbf{1}_q - z_2^* z_1)^{-1}$  is well defined and belongs to  $GL(q, \mathbf{C})$ . So we can extend the definition of  $T$  to the set

$$\tilde{D}_\top = \{(z_1, z_2, z_3) \mid z_i \in D \cup S, 1 \leq i \leq 3, z_1 \top' z_2, z_2 \top' z_3, z_3 \top' z_1\},$$

where by definition  $z \top' w$  is satisfied if  $z$  or  $w$  belongs to  $D$ , and reduces to the condition  $z \top w$  if both  $z$  and  $w$  belong to  $S$ . As  $\tilde{D}_\top$  is stable by  $(z_1, z_2, z_3) \mapsto (tz_1, tz_2, tz_3)$  for  $0 \leq t \leq 1$ , this is a simply connected set. For  $z_1 \in D$ ,  $\det T(z_1, z_1, z_1)$  is a positive real number. So there is a well defined continuous determination of the argument of  $\det(T(z_1, z_2, z_3))$  on  $\tilde{D}_\top$  such that it takes the value 0 whenever  $z_1 = z_2 = z_3 \in D$ . Denote this determination by  $\arg \det T(z_1, z_2, z_3)$ . It is clearly invariant under the  $G$ -action, and so it defines an invariant for the  $G$ -orbits.

On the other hand, let

$$S(z_1, z_2, z_3) = T(z_1, z_2, z_3)^{*^{-1}} T(z_1, z_2, z_3)$$

be the angular matrix associated to  $T(z_1, z_2, z_3)$ .

**THEOREM 4.4.** *Let  $(\sigma_1, \sigma_2, \sigma_3)$  and  $(\tau_1, \tau_2, \tau_3)$  belong to  $S^3_\top$ . They belong to the same  $G$ -orbit if and only if  $S(\sigma_1, \sigma_2, \sigma_3)$  and  $S(\tau_1, \tau_2, \tau_3)$  are conjugate under  $GL(q, \mathbf{C})$  and  $\arg \det T(\sigma_1, \sigma_2, \sigma_3) = \arg \det T(\tau_1, \tau_2, \tau_3)$ .*

*Proof.* This is a direct consequence of Theorem 4.3 and Theorem 3.13.

**REMARK 1.** Let us consider the case where  $q = 1$ . The Stiefel manifold is the unit sphere  $S^{2p-1}$  in  $\mathbf{C}^p$ . The transversality condition  $\sigma \top \tau$  just means  $\sigma \neq \tau$ , as is easily seen from the Cauchy-Schwarz inequality. The triple ratio

is the complex number

$$T(\sigma_1, \sigma_2, \sigma_3) = (1 - \sigma_2^* \sigma_1)^{-1} (1 - \sigma_2^* \sigma_3) (1 - \sigma_1^* \sigma_3)^{-1}.$$

The group  $GL(q, \mathbf{C}) \simeq \mathbf{C}^*$  acts on the upper halfplane by  $(\lambda, z) \mapsto |\lambda|^2 z$  and so the orbits are described by the argument of the complex number  $z$ . So the characteristic invariant in this case is just

$$\arg \left( (1 - \sigma_2^* \sigma_1)^{-1} (1 - \sigma_2^* \sigma_3) (1 - \sigma_1^* \sigma_3)^{-1} \right).$$

It is equivalent to the invariant  $\theta$  considered in [KR]. This invariant, almost in our terms, was known to E. Cartan (see [Ca]).

REMARK 2. Let us consider the case where  $p = q$ . Then the Stiefel manifold is  $U(q)$ , and the content of Proposition 4.2 is that for  $(\sigma_1, \sigma_2, \sigma_3) \in S_{\top}^3$

$$T(\sigma_1, \sigma_2, \sigma_3) = (1 - \sigma_2^* \sigma_1)^{-1} (1 - \sigma_2^* \sigma_3) (1 - \sigma_1^* \sigma_3)^{-1}$$

is an invertible skew-Hermitian matrix. The orbits of  $GL(q, \mathbf{C})$  in its action on nondegenerate Hermitian forms are characterized by the signature. So the characteristic invariant as described in Theorem 4.3 in this case reduces to  $\text{sgn } iT(\sigma_1, \sigma_2, \sigma_3)$ . As concerns Theorem 4.4, notice that the invariant  $S$  is trivial (equal to  $-\mathbf{1}_q$ ), so one is only concerned with the invariant  $\arg \det T$ . The bounded domain  $D$  is of tube type and the description of the invariant through the function  $\arg \det$  coincides with the approach of this problem in [CØ], where the invariant was introduced under the name of *generalized Maslov index*.

## REFERENCES

- [Ca] CARTAN, E. Sur le groupe de la géométrie hypersphérique. *Comment. Math. Helv.* 4 (1932), 158–171.
- [CØ] CLERC, J.-L. and B. ØRSTED. The Maslov index revisited. *Transform. Groups* 6 (2001), 303–320.
- [FK] FARAUT, J. and A. KORÁNYI. *Analysis on Symmetric Cones*. Oxford Mathematical Monographs, Clarendon Press, Oxford, 1994.
- [Fal] FARAUT, J. et al. *Analysis and Geometry on Complex Homogeneous Domains*. Progress in Mathematics 185. Birkhäuser Verlag, Boston, 2000.
- [H] HUA, L.-K. Geometries of matrices, I. Generalizations of von Staudt's theorem. *Trans. Amer. Math. Soc.* 57 (1945), 441–481.
- [KR] KORÁNYI, A. and H. M. REIMANN. The complex cross ratio on the Heisenberg group. *L'Enseign. Math.* (2) 33 (1987), 291–300.