

## 5.3 The special unitary group

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **49 (2003)**

Heft 3-4: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **12.07.2024**

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

### **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

$$(\Psi_i - \Psi_j)(\exp \nu) = (\nu - \mu_i) - (\nu - \mu_j) = \mu_j - \mu_i.$$

By equivariance, it follows that  $\Psi_i - \Psi_j$  takes values in the adjoint orbit through  $\mu_j - \mu_i$ . The difference  $\varpi_i - \varpi_j$  vanishes on the maximal torus  $T$ , and is therefore determined by its contractions with generating vector fields. Since  $\Psi_i - \Psi_j$  is a moment map for  $\varpi_i - \varpi_j$ , it follows that  $\varpi_i - \varpi_j$  equals the pull-back of the symplectic form on  $G \cdot (\mu_j - \mu_i)$ .

### 5.3 THE SPECIAL UNITARY GROUP

For the special unitary group  $G = \mathrm{SU}(d+1)$ , the construction of the basic gerbe simplifies due to the fact that in this case all vertices  $\mu_j$  of the alcove are contained in the weight lattice. In fact the gerbe is presented as a Chatterjee-Hitchin gerbe for the cover  $\mathcal{V} = \{V_i, i = 0, \dots, d\}$ .

For each weight  $\mu \in \Lambda^* \subset \mathfrak{t} \subset \mathfrak{g}$ , let  $G_\mu$  be its stabilizer for the adjoint action and let  $\mathbf{C}_\mu$  the 1-dimensional  $G_\mu$ -representation with infinitesimal character  $\mu$ . Let the line bundle  $L_\mu = G \times_{G_\mu} \mathbf{C}_\mu$  equipped with the unique left-invariant connection  $\nabla$ . Then  $L_\mu$  is a  $G$ -equivariant pre-quantum line bundle for the orbit  $\mathcal{O} = G \cdot \mu$ . That is,

$$\frac{i}{2\pi} \mathrm{curv}_G(\nabla) = (\omega_{\mathcal{O}})_G := \omega_{\mathcal{O}} - \Phi_{\mathcal{O}}$$

where  $\omega_{\mathcal{O}}$  is the symplectic form and  $\Phi_{\mathcal{O}}: \mathcal{O} \hookrightarrow \mathfrak{g}^*$  is the moment map given as inclusion.

In particular, in the case of  $\mathrm{SU}(d+1)$  all orbits  $\mathcal{O}_{ij} = G \cdot (\mu_j - \mu_i)$  carry  $G$ -equivariant pre-quantum line bundles. Recall the fibrations  $p_{ij}: V_{ij} \rightarrow \mathcal{O}_{ij}$  defined by  $\Psi_i - \Psi_j$ , and let

$$L_{ij} = p_{ij}^*(L_{\mu_j - \mu_i}),$$

equipped with the pull-back connection. For any triple intersection  $V_{ijk} = G \times_{G_{ijk}} S_{ijk}$ , the tensor product  $(\delta L)_{ijk} = L_{jk} L_{ik}^{-1} L_{ij}$  is the pull-back of the line bundle over  $G/G_{ijk}$ , defined by the zero weight

$$(\mu_k - \mu_j) - (\mu_k - \mu_i) + (\mu_j - \mu_i) = 0$$

of  $G_{ijk}$ . It is hence canonically trivial, with  $(\delta \nabla)_{ijk}$  the trivial connection. The trivializing section  $t_{ijk} = 1$  satisfies  $\delta t = 1$  and  $(\delta \nabla)t = 0$ . Take  $(B_j)_G = (\varpi_j)_G$ . Then

$$(B_j)_G - (B_i)_G = (\varpi_j)_G - (\varpi_i)_G = -p_{ij}^*(\omega_{\mathcal{O}_{ij}})_G = \frac{1}{2\pi i} \mathrm{curv}_G(\nabla^{L_{ij}}).$$

Thus  $\mathcal{G} = (\mathcal{V}, L, t)$  is a equivariant gerbe with connection  $(\nabla, B)$ . Since

$$d_G(B_j)_G = d_G(\varpi_j)_G = \eta_G|_{V_j},$$

this is the basic gerbe for  $SU(d + 1)$ . The transition line bundles  $L_{ij}$  may be expressed in terms of eigenspace line bundles, leading to the description of the basic gerbe from the introduction.

REMARK 5.3. This description of the basic gerbe over the special unitary group was found independently by Gawędzki-Reis [13], who also discuss the much more difficult case of quotients of  $SU(d + 1)$  by subgroups of the center.

A similar construction works for the group  $C_d = Sp(d)$ , the only case besides  $A_d = SU(d + 1)$  for which the vertices of the alcove are in the weight lattice. The following table lists, for all simply connected compact simple groups, the smallest integer  $k_0 > 0$  such that  $k_0\mathfrak{A}$  is a weight lattice polytope<sup>4</sup>). The construction for  $SU(d + 1)$  generalizes to describe the  $k_0$ -th power of the basic gerbe in all cases.

(5.2)

$G$	$A_d$	$B_d$	$C_d$	$D_d$	$E_6$	$E_7$	$E_8$	$F_4$	$G_2$
$k_0$	1	2	1	2	6	12	60	6	2

#### 5.4 THE BASIC GERBE FOR GENERAL SIMPLE, SIMPLY CONNECTED $G$

The extra difficulty for the groups with  $k_0 > 1$  comes from the fact that the pull-back maps  $H_G^3(G, \mathbf{Z}) \rightarrow H_G^3(C_j, \mathbf{Z}) \cong H_G^3(V_j, \mathbf{Z})$  may be a non-zero torsion class, in general. In this case the restriction of the basic gerbe to  $V_j$  will be non-trivial. Our strategy for the general case is to first construct equivariant bundle gerbes over  $V_j$ , and then glue the local data as explained in Section 4.

The centralizers  $G_g$  of elements  $g \in G$  are always connected [11, Corollary (3.15)] but need not be simply-connected. The conjugacy classes  $C_j = q^{-1}(\mu_j)$  corresponding to the vertices of the alcove are exactly the conjugacy classes of elements for which the centralizer is semi-simple. Since

$$H_G^3(C_j, \mathbf{Z}) = H_G^3(G/G_j, \mathbf{Z}) = H_{G_j}^3(\text{pt}, \mathbf{Z}),$$

we see that the torsion problem described above is related to a possibly non-trivial central extension of the centralizers  $G_j$  of  $\exp(\mu_j)$  by the circle  $U(1)$ .

<sup>4</sup>) This information is extracted from the tables in Bourbaki [5]. Letting  $w_1, \dots, w_d$  be the fundamental weights, one determines  $k_0$  as the least common multiple of the numbers  $\alpha_{max} \cdot w_j$ , using the basic inner product defined by  $\alpha_{max} \cdot \alpha_{max} = 2$ . The number  $k_0$  is equal to the smallest Dynkin index of a representation  $G \rightarrow SU(n)$ , see [28, p. 128] where the same table appears in a different context.

PROPOSITION 5.4. Any vertex  $\mu_j$  of the alcove  $\mathfrak{A}$  is in the dual of the co-root lattice for the corresponding centralizer  $G_j$ . It hence defines a homomorphism  $\varrho_j \in \text{Hom}(\pi_1(G_j), \text{U}(1))$ , or equivalently a central extension of  $G_j$  by  $\text{U}(1)$ .

*Proof.* Let  $\tilde{G}_j$  be the universal cover of  $G_j$ . A system of simple roots for  $\tilde{G}_j$  is given by the list of all  $\alpha_i$  ( $i = 0, \dots, d$ ) with  $j \neq i$ . The lattice  $\Lambda_j$  is spanned by the corresponding coroots  $\check{\alpha}_i$ . To show that  $\mu_j$  is in the dual of the co-root lattice, we have to verify that  $\langle \mu_j, \check{\alpha}_i \rangle \in \mathbf{Z}$  for  $i \neq j$ . For  $i \neq 0, j$  this is obvious since  $\mu_j(\check{\alpha}_i) = 0$ . For  $i = 0$ , we have  $\|\check{\alpha}_0\|^2 = 2$ , and therefore  $\check{\alpha}_0 = \alpha_0$  and  $\mu_j(\check{\alpha}_0) = \alpha_0(\mu_j) = -1$ .

Recall that for  $i \neq j$ ,  $G_{ij}$  is the centralizer of points  $\exp \mu$  with  $\mu = t\mu_j + (1-t)\mu_i$  for some  $0 < t < 1$ . Let  $\varrho_{ij} \in \text{Hom}(\pi_1(G_{ij}), \text{U}(1))$  be the quotient of  $\pi_1(G_{ij}) \rightarrow \pi_1(G_j) \xrightarrow{\varrho_j} \text{U}(1)$  by the homomorphism  $\pi_1(G_{ij}) \rightarrow \pi_1(G_i) \xrightarrow{\varrho_i} \text{U}(1)$ .

LEMMA 5.5. The difference  $\mu_j - \mu_i \in \mathfrak{g}_{ij}$  is fixed under  $G_{ij}$ , and  $\varrho_{ij} \in \text{Hom}(\pi_1(G_{ij}), \text{U}(1))$  is its image under the exact sequence (3.2) for  $K = G_{ij}$ .

*Proof.* Since  $G_{ij}$  fixes the curve  $g(t) = \exp(t\mu_j + (1-t)\mu_i) = \exp(\mu_i)\exp(t(\mu_j - \mu_i))$ , it stabilizes the Lie algebra element  $\mu_j - \mu_i$ . The second claim is immediate from the definition.

We are now in position to explain our construction of the basic gerbe in the general case. For all  $I \subset \{0, \dots, d\}$  let  $X_I \rightarrow V_I$  be the  $G$ -equivariant principal  $G_I$ -bundle,

$$X_I = G \times S_I \rightarrow V_I = G \times_{G_I} S_I.$$

$X_I$  is the pull-back of the  $G_I$ -bundle  $G \rightarrow G/G_I$ , and in particular carries a  $G$ -invariant connection  $\theta_I$  obtained by pull-back of the unique  $G$ -invariant connection on that bundle. For  $I \supset J$  there are natural  $G$ -equivariant inclusions  $f_I^J: X_I \rightarrow X_J$ , and these are compatible as in Section 4. The homomorphisms  $\varrho_j: \pi_1(G_j) \rightarrow \text{U}(1)$  define flat,  $G$ -equivariant bundle gerbes  $\mathcal{G}_j = (X_j, L_j, t_j)$  over  $V_j$ .

The quotient of the two gerbes on  $V_{ij}$ , obtained by pulling back  $\mathcal{G}_i, \mathcal{G}_j$  to  $X_{ij}$ , is just the gerbe defined by the homomorphism  $\varrho_{ij}: \pi_1(G_{ij}) \rightarrow \text{U}(1)$ . By Lemma 5.5 and Proposition 3.2(b), it follows that this quotient gerbe has

a distinguished, equivariant pseudo-line bundle  $(E_{ij}, s_{ij})$  (where  $E_{ij}$  is trivial), with connection  $\nabla^{E_{ij}}$  induced from the connection  $\theta_{ij}$ . From the definition of  $\theta_{ij}$ , it follows that the equivariant error 2-form for this connection is the pull-back of the equivariant symplectic form on the coadjoint orbit through  $\mu_j - \mu_i$ .

We now modify the bundle gerbe connection by adding the equivariant 2-form  $(\varpi_j)_G \in \Omega_G^2(V_j)$  to the gerbe connection. Proposition 5.2(d) shows that the equivariant error 2-form of  $\nabla^{E_{ij}}$  with respect to the new gerbe connection vanishes. The other conditions from the gluing construction in §4 are trivially satisfied. Since the equivariant 3-curvature for the new gerbe connection on  $\mathcal{G}_j$  is  $d_G(\varpi_j)_G = \eta_G|_{V_j}$ , we have constructed an equivariant bundle gerbe with connection, with equivariant curvature-form  $\eta_G$ .

REMARK 5.6. For  $G = \text{SU}(d + 1)$  this construction reduces to the construction in terms of transition line bundles: All  $L_i, t_i, E_{ij}, u_{ijk}$  are trivial in this case, hence the entire information on the gerbe resides in the functions  $s_{ij}: (X_{ij})^{[2]} \rightarrow \text{U}(1)$  defined by the differences  $\mu_j - \mu_i$ . The condition  $\delta s_{ij} = 1$  for these functions means that  $s_{ij}$  defines a line bundle  $L_{ij}$  over  $V_{ij}$ , as remarked at the beginning of Section 2.2. The condition  $s_{ij}s_{jk}s_{ki} = 1$  over  $X_{ijk}$  is the compatibility condition over triple intersections.

### 6. PRE-QUANTIZATION OF CONJUGACY CLASSES

It is a well-known fact from symplectic geometry that a coadjoint orbit  $\mathcal{O} = G \cdot \mu$  through  $\mu \in \mathfrak{t}_+^*$  has integral symplectic form, i.e. admits a pre-quantum line bundle, if and only if  $\mu$  is in the weight lattice  $\Lambda^*$ . The analogous question for conjugacy classes reads: For which  $\mu \in \mathfrak{A}$  and  $m \in \mathbf{N}$  does the pull-back of the  $m$ th power of the basic gerbe  $\mathcal{G}^m$  to the conjugacy class  $\mathcal{C} = G \cdot \exp(\mu)$  admit a pseudo-line bundle, with  $m\omega_{\mathcal{C}}$  as its error 2-form? For any positive integer  $m > 0$  let

$$\Lambda_m^* = \Lambda^* \cap m\mathfrak{A}$$

be the set of level  $m$  weights. As is well-known [26], the set  $\Lambda_m^*$  parametrizes the positive energy representations of the loop group  $LG$  at level  $m$ .

THEOREM 6.1. *The restriction of  $\mathcal{G}^m$  to a conjugacy class  $\mathcal{C}$  admits a pseudo-line bundle  $\mathcal{L}$  with connection, with error 2-form  $m\omega_{\mathcal{C}}$ , if and only if  $\mathcal{C} = G \cdot \exp(\mu/m)$  with  $\mu \in \Lambda_m^*$ . Moreover  $\mathcal{L}$  has an equivariant extension in this case, with  $m\omega_{\mathcal{C}}$  as its equivariant error 2-form.*