

Objektyp: **ReferenceList**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **53 (2007)**

Heft 1-2

PDF erstellt am: **12.07.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Ein Dienst der *ETH-Bibliothek*
ETH Zürich, Rämistrasse 101, 8092 Zürich, Schweiz, www.library.ethz.ch

<http://www.e-periodica.ch>

denote by $H^{p,q} \subset H^{p+q}(X, \mathbf{C})$ the subspace of de Rham cohomology classes of forms of type (p, q) ; we have $H^{q,p} = \overline{H^{p,q}}$. The fundamental result of Hodge theory is the Hodge decomposition

$$H^n(X, \mathbf{C}) = \bigoplus_{p+q=n} H^{p,q},$$

together with the canonical isomorphisms $H^{p,q} \xrightarrow{\sim} H^q(X, \Omega_X^p)$. In particular,

$$H^2(X, \mathbf{C}) = H^{2,0} \oplus H^{1,1} \oplus H^{0,2},$$

with $H^{2,0} \cong H^0(X, \Omega_X^2)$, embedded into $H^2(X, \mathbf{C})$ by associating to a holomorphic form its De Rham class.

To any hermitian metric g on X is associated a real 2-form ω of type $(1, 1)$, the *Kähler form*, defined by $\omega(V, W) = g(V, JW)$ for any real vector fields V, W ; the metric is Kähler if ω is closed. Then its class in $H^2(X, \mathbf{C})$ is called a Kähler class. The Kähler classes form an open cone in $H_{\mathbf{R}}^{1,1} = H^{1,1} \cap H^2(X, \mathbf{R})$.

Let L be a line bundle on X . The Chern class $c_1(L) \in H^2(X, \mathbf{C})$ is integral, that is comes from $H^2(X, \mathbf{Z})$, and belongs to $H^{1,1}$. Conversely, any integral class in $H^{1,1}$ is the Chern class of some line bundle on X (Lefschetz theorem).

If L is very ample, its Chern class is the pull-back by φ_L of the Chern class of $\mathcal{O}_{\mathbf{P}^1}(1)$, which is a Kähler class, and therefore $c_1(L)$ is a Kähler class. More generally, if L is ample, some multiple of $c_1(L)$ is a Kähler class, hence also $c_1(L)$. Conversely, the celebrated Kodaira embedding theorem asserts that *a line bundle whose Chern class is Kähler is ample*. As a corollary, we see that *any compact Kähler manifold X with $H^0(X, \Omega_X^2) = 0$ is projective*: we have $H^2(X, \mathbf{C}) = H^{1,1}$, hence the cone of Kähler classes is open in $H^2(X, \mathbf{R})$. Therefore it contains integral classes; by the above results such a class is the first Chern class of an ample line bundle, hence X is projective. More generally, the same argument shows that X is projective whenever the subspace $H^{1,1}$ of $H^2(X, \mathbf{C})$ is defined over \mathbf{Q} .

REFERENCES

- [A] ATTYAH, M. On analytic surfaces with double points. *Proc. Roy. Soc. London Ser. A* 247 (1958), 237–244.
- [Ba] BATYREV, V. Birational Calabi-Yau n -folds have equal Betti numbers. *New trends in algebraic geometry (Warwick, 1996)*, 1–11. London Math. Soc. Lecture Note Ser. 264, Cambridge Univ. Press, 1999.

- [B1] BEAUVILLE, A. Variétés kählériennes dont la première classe de Chern est nulle. *J. Differential Geom.* 18 (1983), 755–782.
- [B2] — Surfaces K3. Exp. 609 du Séminaire Bourbaki. *Astérisque* 105–106 (1983), 217–229.
- [B3] — Fano contact manifolds and nilpotent orbits. *Comment. Math. Helv.* 73 (1998), 566–583.
- [B-D] BEAUVILLE, A. et R. DONAGI. La variété des droites d’une hypersurface cubique de dimension 4. *C.R. Acad. Sc. Paris* 301, sér. I (1985), 703–706.
- [Be] BERGER, M. Sur les groupes d’holonomie homogène des variétés à connexion affine et des variétés riemanniennes. *Bull. Soc. Math. France* 83 (1955), 279–330.
- [Bs] BESSE, A. Einstein manifolds. *Ergebnisse der Math.* 10. Springer-Verlag, Berlin, 1999.
- [B-G] BROWN, R. and A. GRAY. Riemannian manifolds with holonomy group $\text{Spin}(9)$. In: *Differential Geometry, in Honor of K. Yano*, 41–59. Kinokuniya, Tokyo (1972).
- [B-Y] BOCHNER, S. and K. YANO. Curvature and Betti numbers. *Annals of Math. Studies* 32. Princeton University Press, 1953.
- [Bg1] BOGOMOLOV, F. Hamiltonian Kähler manifolds. *Soviet Math. Dokl.* 19 (1978), 1462–1465.
- [Bg2] — On the cohomology ring of a simple hyper-Kähler manifold (on the results of Verbitsky). *Geom. Funct. Anal.* 6 (1996), 612–618.
- [Bo] BOURGUIGNON, J.-P. Premières formes de Chern des variétés kählériennes compactes [d’après E. Calabi, T. Aubin et S.T. Yau]. Séminaire Bourbaki, Exp. 507, 1–21. Lecture Notes 710, Springer, Berlin, 1979.
- [C] CALABI, E. On Kähler manifolds with vanishing canonical class. *Algebraic Geometry and Topology (in honor of S. Lefschetz)*, 78–89. Princeton University Press, 1957.
- [De] DEBARRE, O. Un contre-exemple au théorème de Torelli pour les variétés symplectiques irréductibles. *C.R. Acad. Sci. Paris* 299, sér. I (1984), 681–684.
- [Dm] DEMAILLY, J.-P. On the Frobenius integrability of certain holomorphic p -forms. In: *Complex Geometry (Göttingen, 2000)*, 93–98. Springer, Berlin, 2002.
- [F] FOGARTY, J. Algebraic families on an algebraic surface. *Amer. J. Math.* 90 (1968), 511–521.
- [Fr] FRIEDMAN, R. On threefolds with trivial canonical bundle. *Complex Geometry and Lie theory*, 103–134. Proc. Sympos. Pure Math. 53, AMS, 1991.
- [F-N] FUJIKI, A. and S. NAKANO. Supplement to “On the inverse of monoidal transformation”. *Publ. Res. Inst. Math. Sci.* 7 (1971–72), 637–644.
- [H] HUYBRECHTS, D. Compact hyperkähler manifolds: basic results. *Invent. Math.* 135 (1999), 63–113; see also: Erratum, *Invent. Math.* 152 (2003), 209–212.
- [H-L] HUYBRECHTS, D. and M. LEHN. *The Geometry of Moduli Spaces of Sheaves*. Aspects of Math. E31. Vieweg, 1997.

- [H-S] HITCHIN, N. and J. SAWON. Curvature and characteristic numbers of hyper-Kähler manifolds. *Duke Math. J.* 106 (2001), 599–615.
- [J1] JOYCE, D. Compact Riemannian 7-manifolds with holonomy G_2 . *J. Differential Geom.* 43 (1996), 291–375.
- [J2] ——— Compact Riemannian 8-manifolds with holonomy $\text{Spin}(7)$. *Invent. Math.* 123 (1996), 507–552.
- [J3] ——— Compact manifolds with exceptional holonomy. *Doc. Math. J. DMV Extra Volume ICM II* (1998), 361–370.
- [K] KAPRANOV, M. Rozansky-Witten invariants via Atiyah classes. *Compositio Math.* 115 (1999), 71–113.
- [K-L-S] KALEDIN, D., M. LEHN and C. SORGER. Singular symplectic moduli spaces. *Invent. math.* 164 (2006), 591–614.
- [KPSW] KEBEKUS, S., T. PETERNELL, A. SOMMESE and J. WIŚNIEWSKI. Projective contact manifolds. *Invent. Math.* 142 (2000), 1–15.
- [K-N] KOBAYASHI, S. and K. NOMIZU. *Foundations of Differential Geometry*. Wiley-Interscience Publications, 1963.
- [Ko] KONTSEVICH, M. Homological algebra of mirror symmetry. *Proc. ICM Zürich 1994*, vol. 1, 120–139. Birkhäuser, Basel, 1995.
- [L] LE BRUN, C. Fano manifolds, contact structures, and quaternionic geometry. *Internat. J. Math.* 6 (1995), 419–437.
- [M] MUKAI, S. Symplectic structure of the moduli space of sheaves on an abelian or K3 surface. *Invent. Math.* 77 (1984), 101–116.
- [N] NAMIKAWA, Y. Counter-example to global Torelli problem for irreducible symplectic manifolds. *Math. Ann.* 324 (2002), 841–845.
- [Ni] NIEPER, M. Hirzebruch-Riemann-Roch formulae on irreducible symplectic Kähler manifolds. *J. Algebraic Geom.* 12 (2003), 715–739.
- [O] OLMOS, C. A geometric proof of the Berger holonomy theorem. *Ann. of Math. (2)* 161 (2005), 579–588.
- [OG1] O’GRADY, K. The weight-two Hodge structure of moduli spaces of sheaves on a K3 surface. *J. Algebraic Geom.* 6 (1997), 599–644.
- [OG2] ——— Desingularized moduli spaces of sheaves on a K3. *J. reine angew. Math.* 512 (1999), 49–117.
- [OG3] ——— A new six-dimensional irreducible symplectic variety. *J. Algebraic Geom.* 12 (2003), 435–505.
- [P] *Géométrie des surfaces K3 : modules et périodes* (séminaire Palaiseau 81–82). *Astérisque* 126 (1985).
- [R] DE RHAM, G. Sur la réductibilité d’un espace de Riemann. *Comment. Math. Helv.* 26 (1952), 328–344.
- [R-W] ROZANSKY, L. and E. WITTEN. Hyper-Kähler geometry and invariants of three-manifolds. *Selecta Math. (N.S.)* 3 (1997), 401–458.
- [S] SALAMON, S. Quaternionic Kähler manifolds. *Invent. Math.* 67 (1982), 143–171.
- [Si] SIMONS, J. On the transitivity of holonomy systems. *Ann. of Math. (2)* 76 (1962), 213–234.
- [S-Y-Z] STROMINGER, A., S. T. YAU and E. ZASLOW. Mirror symmetry is T -duality. *Nuclear Phys. B* 479 (1996), 243–259.

- [T] TIAN, G. Smoothness of the universal deformation space of compact Calabi-Yau manifolds and its Petersson-Weil metric. Math. aspects of string theory, 629–646. *Adv. Ser. Math. Phys. 1*. World Scientific, Singapore, 1987.
- [Va] VAROUCHAS, J. Stabilité de la classe des variétés kählériennes par certains morphismes propres. *Invent. Math.* 77 (1984), 117–127.
- [V1] VOISIN, C. *Symétrie miroir. Panoramas et Synthèses 2*. SMF, 1996.
- [V2] — On the homotopy types of compact Kähler and complex projective manifolds. *Invent. Math.* 157 (2004) 329–343.
- [W] WOLF, J. Complex homogeneous contact manifolds and quaternionic symmetric spaces. *J. Math. Mech.* 14 (1965), 1033–1047.
- [Y] YAU, S. T. On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation, I. *Comm. Pure Appl. Math.* 31 (1978), 339–411.

(Reçu le 17 janvier 2007)

Arnaud Beauville

Institut Universitaire de France et Laboratoire J.-A. Dieudonné
Université de Nice
Parc Valrose
F-06108 Nice Cedex 2
e-mail: beauville@math.unice.fr