

Blocking light in compact Riemannian manifolds

Autor(en): **Schmidt, Benjamin**

Objektyp: **Article**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **54 (2008)**

Heft 1-2

PDF erstellt am: **13.09.2024**

Persistenter Link: <https://doi.org/10.5169/seals-109927>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

58

BLOCKING LIGHT IN CLOSED RIEMANNIAN MANIFOLDS

by Benjamin SCHMIDT

To what extent does the collision of light determine the global geometry of space? In this note we'll discuss two conjectures, both of which assert that the focusing behavior of light in locally symmetric Riemannian manifolds are unique to these spaces. Throughout, (M, g) denotes a C^∞ -smooth, connected, and compact manifold without boundary equipped with a C^∞ -smooth Riemannian metric g . Geodesic segments are identified with their unit speed parametrization $\gamma: [0, L_\gamma] \rightarrow M$, where L_γ is the length of the segment γ .

DEFINITION 58.1 (Light). Let $X, Y \subset (M, g)$ be two nonempty subsets, and let $G_g(X, Y)$ denote the set of geodesic segments with initial point $\gamma(0) \in X$ and terminal point $\gamma(L_\gamma) \in Y$. The *light from X to Y* is the set

$$L_g(X, Y) = \{\gamma \in G_g(X, Y) \mid \text{interior}(\gamma) \cap (X \cup Y) = \emptyset\}.$$

DEFINITION 58.2 (Blocking Set). Let $X, Y \subset M$ be two nonempty subsets. A subset $B \subset M$ is a *blocking set* for $L_g(X, Y)$ provided that for every $\gamma \in L_g(X, Y)$,

$$\text{interior}(\gamma) \cap B \neq \emptyset.$$

We focus on closed Riemannian manifolds for which the light between pairs of points in M is blocked by a finite set of points. By a theorem of Serre ([5]), $G_g(x, y)$ is infinite when $x, y \in M$ are distinct points. However, $L_g(x, y) \subset G_g(x, y)$ may or may not be an infinite subset. This is the case, for example, in a round sphere where all of the infinitely many geodesics between a typical pair of points cover a single periodic geodesic.

DEFINITION 58.3 (Blocking Number). Let $x, y \in M$ be two (possibly not distinct) points in M . The *blocking number* $b_g(x, y)$ for $L_g(x, y)$ is defined as

$$b_g(x, y) = \inf\{n \in \mathbf{N} \cup \{\infty\} \mid L_g(x, y) \text{ is blocked by } n \text{ points}\}.$$

Our starting point is the following surprising theorem from [2]:

THEOREM 58.4 (Gutkin). *Let (M, g) be a closed flat Riemannian manifold. Then there exists an $n \in \mathbf{N}$ depending only on the dimension of M such that $b_g \leq n$ as a function on $M \times M$.*

We believe the following is true:

CONJECTURE 58.5. *Let (M, g) be a closed Riemannian manifold. If $b_g < \infty$ then g is a flat metric.*

This conjecture is true for Riemannian metrics of nonpositive sectional curvatures as shown independently by Burns–Gutkin and Lafont–Schmidt ([1], [4]). The focusing of light is also interesting in the context of the compact type locally symmetric spaces. In [3], Gutkin and Schroeder establish the following:

THEOREM 58.6 (Gutkin–Schroeder). *Let (M, g) be a closed locally symmetric space of compact type with \mathbf{R} -rank $k \geq 1$. Then $b_g(x, y) \leq 2^k$ for almost all $(x, y) \in M \times M$.*

We refer the reader to [3] for a more precise formulation and discussion of this result. Presently, we'll restrict attention to the compact rank one symmetric spaces or CROSSes. The CROSSes are classified and consist of the round spheres and the various projective spaces. The CROSSes all satisfy the following blocking property:

DEFINITION 58.7 (Cross Blocking). A closed Riemannian manifold (M, g) has property *CB* if

$$0 < d(x, y) < \text{Diam}(M, g) \implies b_g(x, y) \leq 2.$$

Round spheres additionally satisfy the following blocking property, a blocking interpretation of antipodal points:

DEFINITION 58.8 (Sphere Blocking). A closed Riemannian manifold (M, g) has property SB if $b_g(x, x) = 1$ for every $x \in M$.

We believe the following is true:

CONJECTURE 58.9. A simply connected closed Riemannian manifold (M, g) has property CB if and only if (M, g) is isometric to a simply connected compact rank one symmetric space. In particular, (M, g) has properties CB and SB if and only if (M, g) is isometric to a round sphere.

In [4], special cases of this conjecture are confirmed under various additional hypotheses.

REFERENCES

- [1] BURNS, K. and E. GUTKIN. Growth of the number of geodesics between points and insecurity for Riemannian manifolds. *Discrete Contin. Dyn. Syst.* 21 (2008), 403–413.
- [2] GUTKIN, E. Blocking of billiard orbits and security for polygons and flat surfaces. *Geom. Funct. Anal.* 15 (2005), 83–105.
- [3] GUTKIN, E. and V. SCHROEDER. Connecting geodesics and security of configurations in compact locally symmetric spaces. *Geom. Dedicata* 118 (2006), 185–208.
- [4] LAFONT, J.-F. and B. SCHMIDT. Blocking light in compact Riemannian manifolds. *Geom. Topol.* 11 (2007), 867–887.
- [5] SERRE, J.-P. Homologie singulière des espaces fibrés. *Ann. of Math.* (2) 54 (1951), 425–505.

B. Schmidt

University of Chicago
Department of Mathematics
5734 S. University Ave.
Chicago, IL 60637
USA
e-mail: schmidt@math.uchicago.edu