

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 54 (2008)
Heft: 1-2

Artikel: Finiteness properties of function-field-arithmetic groups
Autor: Wortman, Kevin
DOI: <https://doi.org/10.5169/seals-109934>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 13.01.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

65

FINITENESS PROPERTIES OF FUNCTION-FIELD-ARITHMETIC GROUPS

by Kevin WORTMAN

Let K be a global function field, and S a finite nonempty set of pairwise inequivalent valuations on K . We let $\mathcal{O}_S \leq K$ be the corresponding ring of S -integers, and we let \mathfrak{G} be a simple K -group. For any $v \in S$, we let K_v be the completion of K with respect to v .

The finiteness properties of function-field-arithmetic groups such as $\mathfrak{G}(\mathcal{O}_S)$ have been of interest for about 50 years. Some of that interest has fallen on which of these groups are of type FP_m for a given m . For example, for which m is it true that $\text{SL}_n(\mathbf{F}_q[t])$ is of type FP_m where $\mathbf{F}_q[t]$ denotes a polynomial ring with one variable and coefficients in a finite field with q elements.

We recall that a group Γ is of type FP_m if there exists a partial resolution

$$P_m \rightarrow P_{m-1} \rightarrow \cdots \rightarrow P_1 \rightarrow P_0 \rightarrow \mathbf{Z} \rightarrow 0$$

of the trivial $\mathbf{Z}\Gamma$ -module \mathbf{Z} by finitely generated projective $\mathbf{Z}\Gamma$ -modules.

All of the evidence thus far indicates the existence of a solution for the following

CONJECTURE 65.1. *With $\mathfrak{G}(\mathcal{O}_S)$ as above and with $k = \sum_{v \in S} \text{rank}_{K_v}(\mathfrak{G})$, the arithmetic group $\mathfrak{G}(\mathcal{O}_S)$ is of type FP_m if and only if either $\text{rank}_K(\mathfrak{G}) = 0$ or $k > m$.*

For example, according to the above conjecture $\text{SL}_n(\mathbf{F}_q[t])$ should be of type $\text{FP}_{(n-2)}$ but not of type $\text{FP}_{(n-1)}$. In fact this was shown independently by Abels and Abramenko for large values of q .

Plenty of other evidence exists to support the conjecture in general including a proof of the “only if” implication [2]. The existing evidence for the

conjecture can be found in papers of Abels, Abramenko, Behr, Bux–Wortman, Hurrelbrink, Keller, Kneser, Lubotzky, McHardy, Nagao, O’Meara, Rehmman–Soulé, Serre, Splitthoff, and Stuhler. For detailed references, see e.g. [1] and [2].

REFERENCES

- [1] BEHR, H. Arithmetic groups over function fields. I. A complete characterization of finitely generated and finitely presented arithmetic subgroups of reductive algebraic groups. *J. Reine Angew. Math.* 495 (1998), 79–118.
- [2] BUX, K.-U. and K. WORTMAN. Finiteness properties of arithmetic groups over function fields. *Invent. Math.* 167 (2007), 355–378.

K. Wortman

University of Utah
Department of Mathematics
155 South 1400 East
Salt Lake City, UT 84112-0090
USA
e-mail: wortman@math.utah.edu