

# Summary

Objekttyp: **Chapter**

Zeitschrift: **Veröffentlichungen des Geobotanischen Institutes der Eidg. Tech. Hochschule, Stiftung Rübel, in Zürich**

Band (Jahr): **67 (1980)**

PDF erstellt am: **03.07.2024**

## Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

## Summary

A survey made in 1970 on the distribution of the yew-tree (*Taxus baccata*) in Switzerland shows that this species has decreased considerably since 1904 when a similar survey was made, and that the next tree generation is practically non-existent. The yew-tree maintains a peculiar "in-between" position, i.e. between deciduous tree and evergreen conifer as well as between pioneer and climax tree. Owing to this position and the genetically controlled small height growth the species in question could grow in a large number of ecological niches of our forests as a secondary stand tree.

From a climatic point of view the potential range of the yew-tree covers half of Switzerland. The species can be found in all regions with oceanic to subcontinental climate and which are not subject to heavy winter frosts. Compared to beech with a similar distribution, yew-trees tolerate drier air and do not suffer from late frost. Their range therefore extends further into the continental zones.

At present, however, yew-trees occur in only half of their potential area. Site factors such as local climate, water properties, skeletal and nutrient content of the soil as well as the influence of man were investigated as potential limiting factors within the climatic zone. It was found that parent material of soils and configuration of ground were the only natural limiting factors: marly and unstabilized soils discourage beech. This, together with better light conditions on steep slopes promote the growth of yew-trees.

The 260 vegetation surveys reveal the wide phytosociological and accordingly ecological range of this species. Although the survey confirmed *Taxo-Fagetum* as the association for the area investigated, its range proved to be smaller than previously thought. Many stands containing yew-trees had to be attributed to other associations.

An ecogram has been made as a synthesis of the ecological and phytosociological results, integrating edaphic factors as well as competitive influence of beech and pine. The ecogram shows that the ecological niche of yew-trees is determined by extreme site factors which act physiologically as barriers (mainly hydrological factors) and by ecological conditions (mainly light competition).

However, all these ecological and botanical aspects do not fully explain why the yew-tree is missing from large parts of its potential range. This is in fact due to human influence. The yew-tree has been connected with human civilization for thousands of years. The investigation showed that already early civilizations have caused severe damage to this species. In addition, differences in local forest management systems over the last two centuries have further contributed to the existing pattern of distribution.

The present decline of yew-trees is based mainly on two factors: the conversion of coppice with standards into much darker high forests and the excessive game population density (also due to human influence). Nearly all young yew-trees suffer severely from browsing damage. In conclusion, we can say that the present yew-tree stands should be considered as relicts. Without active measures of protection and promotion in the near future this peculiar tree species will almost certainly disappear completely from Swiss forests with the death of the present tree generation - which would be yet another step towards the growing depletion of nature in our technical civilization.