

# On the asymptotic behaviour of atomic spectra near the continuum

Autor(en): **Hunziker, Walter**

Objektyp: **Article**

Zeitschrift: **Helvetica Physica Acta**

Band (Jahr): **48 (1975)**

Heft 1

PDF erstellt am: **10.08.2024**

Persistenter Link: <https://doi.org/10.5169/seals-114666>

## **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

# On the Asymptotic Behaviour of Atomic Spectra near the Continuum

by **Walter Hunziker**

Institut für theoretische Physik, ETH Hönggerberg, 8049 Zürich, Switzerland

(22. XI. 74)

*Abstract.* If  $\Sigma$  is the beginning of the continuous energy spectrum of an atom in any multiplet system, it is shown that this system contains groups of eigenvalues close to  $E_n = \Sigma - n^{-2}$ , compared with  $E_{n+1} - E_n$ , as  $n = 1, 2, \dots \rightarrow \infty$ .

## 1. Introduction

We consider the Hamiltonian

$$H = \sum_{i=1}^N \frac{p_i^2}{2m_i} + \sum_{i < k} \frac{e_i e_k}{|x_i - x_k|}$$

of a system of  $N$  charged particles, with centre of mass (CM) removed. The following is known about the spectrum  $\sigma(H)$  of  $H$  in any subspace of total angular momentum  $L$  or of given symmetry under permutations of identical particles:

- $\sigma(H)$  consists of a continuum  $[\Sigma, \infty]$  and, in the complement of this continuum, only of eigenvalues with finite multiplicities which can accumulate only at  $\Sigma$ .  $\Sigma$  is the lowest threshold for break-up of the system into independent parts [1, 2].
- If  $\Sigma$  is the threshold for break-up into two sub-systems  $C_1, C_2$  with total charges  $q_1, q_2$  such that  $q_1 q_2 < 0$ , then the number of eigenvalues below  $\Sigma$  is infinite. This is always the case for atoms [2].

Following a suggestion of SIMON [2], we want to study the asymptotic behaviour of the eigenvalues in case b). We introduce coordinates adapted to the decomposition  $(C_1, C_2)$  of the set of particles  $(1 \dots N)$ :

$x \in R^3$  = position of the CM of  $C_2$  with respect to the CM of  $C_1$ ,  
 $y_i \in R^{(3n_i-3)}$  = internal coordinates for the system  $C_i$ , which are taken as linear combinations of the cartesian coordinates  $x_1 \dots x_N$ .  $n_i$  stands for the number of particles in  $C_i$ .

If  $C_i$  has total mass  $M_i$ , we choose units such that  $\hbar = 1$ ,  $q_1 q_2 = -2$ ,  $M_1^{-1} + M_2^{-1} = 2$ . Then  $H$  takes the form

$$H = p^2 + V(x, y_1, y_2) + h_1 + h_2,$$

where  $p$  = momentum conjugate to  $x$ ,  $V$  = sum of all interactions linking  $C_1$  and  $C_2$  and  $h_i$  = Hamiltonian of  $C_i$  with CM removed. Now let

$$h_i \varphi_i = \varepsilon_i \varphi_i$$

for the ground state  $\varphi_i$  of  $C_i$ . For simplicity, we ignore possible degeneracies, and if  $C_i$  is a single particle we have, of course,  $h_i = 0$ ,  $\varepsilon_i = 0$ ,  $\varphi_i = 1$ . The operator

$$p^2 - 2|x|^{-1} + h_1 + h_2$$

has eigenvalues

$$E_n = \varepsilon_1 + \varepsilon_2 - n^{-2} = \Sigma - n^{-2}, \quad n = 1, 2, \dots$$

of multiplicity  $n^2$  with eigenfunctions

$$\psi_n = \eta_n(x) \varphi_1(y_1) \varphi_2(y_2),$$

where  $\eta_n$  satisfies

$$(p^2 - 2|x|^{-1}) \eta_n = -n^{-2} \eta_n.$$

We shall assume that  $\eta_n$ ,  $\varphi_1$ ,  $\varphi_2$ , are normalized to 1 in their respective  $L^2$ -spaces.

Since  $V(x, y_1, y_2) \sim -2|x|^{-1}$  for bounded  $y_i$  and  $|x| \rightarrow \infty$  one expects  $E_n$ ,  $\psi_n$  to become 'approximate' eigenvalues and eigenfunctions of  $H$  as  $n \rightarrow \infty$ . More precisely, the question is how rapidly  $\|(H - E_n)\psi_n\| \rightarrow 0$  as  $n \rightarrow \infty$ . This question can be answered now that *pointwise exponential bounds* are available for the bound state wave functions  $\varphi_i$  [3].

## 2. Basic Inequalities

For real  $E$  and  $a > 0$  let

$$N_a = \text{spectral subspace of } H \text{ where } (H - E)^2 \leq a^2. \quad (1)$$

Suppose that  $M$  is a subspace of  $D(H)$  such that for all  $\psi \in M$ ,  $\|\psi\| = 1$ ,

$$\|(H - E)\psi\| \leq \varepsilon. \quad (2)$$

If  $\psi'$  is the component of  $\psi$  orthogonal to  $N_a$  we then have  $\varepsilon \geq \|(H - E)\psi'\| > a \|\psi'\|$ , hence

$$\|\psi'\| = \text{dist}(\psi, N_a) < \varepsilon a^{-1}. \quad (3)$$

Setting  $a = \varepsilon$  it follows that the projection of  $M$  onto  $N_\varepsilon$  is injective, hence

$$\dim N_\varepsilon \geq \dim M. \quad (4)$$

If  $E + \varepsilon < \Sigma$ , we conclude that  $H$  possesses eigenvalues of total multiplicity  $\geq \dim M$  in the interval  $[E - \varepsilon, E + \varepsilon]$ .

In order to deal with symmetries in the case of identical particles, we consider a projection  $P$  (projecting onto a subspace of given symmetry under permutations)

commuting with  $H$ . Let  $M$  be as before and assume in addition to (2) that

$$\|P\psi\| \geq \lambda \tag{5}$$

for some  $\lambda > 0$  and all  $\psi \in M$ ,  $\|\psi\| = 1$ . Then  $\dim PM = \dim M$  and

$$\|(H - E)P\psi\| \leq \varepsilon \leq \varepsilon\lambda^{-1}\|P\psi\|.$$

Therefore, if  $E + \varepsilon\lambda^{-1} < \Sigma$ ,  $HP$  possesses eigenvalues of total multiplicity  $\geq \dim M$  within the bounds  $E \pm \varepsilon\lambda^{-1}$  and  $\|P\psi\|^{-1}P\psi$  has distance less than  $\varepsilon(a\lambda)^{-1}$  from  $PN_a$ .

### 3. Estimate of $\|(H - E_n)\psi_n\|$

$$\begin{aligned} \|(H - E_n)\psi_n\| &= \|(V + 2|x|^{-1})\psi_n\| \\ &\leq \sum_{\substack{i \in C_1 \\ k \in C_2}} |e_i e_k| \|U_{ik}\psi_n\| \end{aligned}$$

with

$$\begin{aligned} U_{ik} &= | |x - z_1 + z_2|^{-1} - |x|^{-1} | \\ &\leq |x|^{-1} |z_1 - z_2| |x - z_1 + z_2|^{-1}, \end{aligned}$$

where  $z_1(y_1)$ ,  $z_2(y_2)$  are the positions of particles  $i$ ,  $k$  relative to the CM of  $C_1$ ,  $C_2$ , respectively. Therefore

$$\|U_{ik}\psi_n\|^2 \leq \int dx |x|^{-2} f(x) |\eta_n(x)|^2$$

with

$$f(x) = \int dz_1 dz_2 \rho_1(z_1) \rho_2(z_2) |z_1 - z_2|^2 |x - z_1 + z_2|^{-2},$$

where  $\rho_i$  is the probability density for  $z_i$  in the state  $\varphi_i$ . From the exponential bounds [3] for  $\varphi_i(y_i)$  it follows that  $\rho_i(z) \leq \text{const} \cdot \exp(-\alpha|z|)$  for some  $\alpha > 0$ . Therefore,  $f(x) \leq \text{const} |x|^{-2}$  and

$$\|(H - E_n)\psi_n\| \leq \text{const} (\eta_n, |x|^{-4} \eta_n)^{1/2}, \tag{6}$$

with a constant independent of  $n$  and  $\eta_n$ . The matrix elements of  $|x|^{-4}$  in the conventional basis  $\eta_{nlm}(x) = R_{nl}(|x|) Y_{lm}(x/|x|)$  ( $l = 0, 1, \dots, n - 1$ ;  $m = l, \dots, -l$ ) are known [4]:

$$(\eta_{nlm}, |x|^{-4} \eta_{n'l'm'}) = \delta_{ll'} \delta_{mm'} [2n^5(l - \frac{1}{2})l(l + \frac{1}{2})(l + 1)(l + \frac{3}{2})]^{-1} [3n^2 - l(l + 1)] \tag{7}$$

for  $l > 0$ . We now restrict  $\eta_n$  to the subspace spanned by the basis vectors  $\eta_{nlm}$  with

$$l + 1 \geq n^{(2\alpha-3)/5} \quad (\alpha \leq 4) \tag{8}$$

the corresponding  $\psi_n$  then span a subspace  $M_n$  with

$$\dim M_n = n^2 - ([n^{(2\alpha-3)/5}] - 1)^2$$

where  $[a]$  = smallest integer  $\geq a$ . It follows from (6), (7) and (8) that for  $\alpha \leq 4$

$$\|(H - E_n)\psi_n\| \leq Cn^{-\alpha} \quad (9)$$

for all  $\psi_n \in M_n$ ,  $\|\psi_n\| = 1$ , with  $C$  not depending on  $n$ ,  $\psi_n$  or  $\alpha$ . From the inequalities of Section 2 we therefore obtain:

*Theorem:* There exists a constant  $C$  such that for  $3 < \alpha \leq 4$   $H$  possesses at least

$$n^2 - ([n^{(2\alpha-3)/5}] - 1)^2$$

eigenvalues (including multiplicities) in the intervals

$$I_n = [E_n - Cn^{-\alpha}, E_n + Cn^{-\alpha}],$$

if  $n$  is sufficiently large so that  $n^{-2} > Cn^{-\alpha}$ . For  $a_n = Cn^{-\beta}$ ,  $3 < \beta < \alpha$ , and all  $\psi_n \in M_n$ ,  $\|\psi_n\| = 1$ ,

$$\text{dist}(\psi_n, N_{a_n}) < n^{-(\alpha-\beta)},$$

where  $N_a$  and  $M_n$  are defined by (1) and (8).

*Remarks:* The conditions  $3 < \beta < \alpha$  merely serve to make the results significant. For example, the condition  $3 < \alpha$  implies that  $I_{n+1}$  is disjoint from  $I_n$  for  $n$  sufficiently large.

Is it necessary to increase  $l$  with  $n$ ? For  $|x| \ll n^2$  the radial Coulomb wave functions  $R_{nl}(|x|)$  behave like  $n^{-3/2}f_l(|x|)$  with  $f_l$  independent of  $n$  [5]. Therefore

$$\|(H - E_n)\psi_n\| = O(n^{-3/2})$$

for fixed  $l$  and  $n \rightarrow \infty$ , and this decrease is too slow since  $\Sigma - E_n = n^{-2}$ . But even the first-order perturbation  $(\psi_n, (V + 2|x|^{-1})\psi_n)$  of  $E_n$  is of the same order as  $E_{n+1} - E_n$ , namely  $O(n^{-3})$ . This shows the difficulty of proving a similar theorem in a subspace of fixed orbital angular momentum  $L$ .

*Inclusion of short-range forces:* The estimate (6) remains valid if  $V$  also contains two-body potentials  $W_{ik}(x_i - x_k)$  which are locally  $L^2$  and vanish at least like  $|x_i - x_k|^{-2}$  for  $|x_i - x_k| \rightarrow \infty$ .

#### 4. Identical Particles

If the system contains identical particles, the theorem remains true in any sector of given symmetry, with  $C$  replaced by  $C\lambda^{-1}$ , provided that (5) holds with  $\lambda$  independent of  $n$  for  $n$  sufficiently large.

To prove (5) one only has to show that the overlap-integrals vanish as  $n \rightarrow \infty$ , i.e. that

$$\lim_{n \rightarrow \infty} \sup_{\substack{\psi_n \in M_n \\ \|\psi_n\| = 1}} (\psi_n, \pi\psi_n) = 0, \quad (10)$$

where  $\pi\psi_n$  is obtained from  $\psi_n$  by a permutation  $\pi$  of identical particles which sends the partition  $(C_1, C_2)$  of  $(1 \dots N)$  into a *different* partition  $(C'_1, C'_2)$ . For a discussion of symmetries under permutations see [2].

The overlap-integral  $(\psi_n, \pi\psi_n)$  is of the form

$$I = \int dx dy_1 dy_2 \overline{\eta_n(x) \varphi_1(y_1) \varphi_2(y_2)} \eta_n(x') \varphi_1(y'_1) \varphi_2(y'_2),$$

where  $x', y'_1, y'_2$  are coordinates adapted to the decomposition  $(C'_1, C'_2)$ . We distinguish the regions

- (1)  $|y_1| > R$  or  $|y'_1| > R$
- (2)  $|y_2| > R$  or  $|y'_2| > R$
- (3)  $|y_i| \leq R$  and  $|y'_i| \leq R$  ( $i = 1, 2$ ),

and denote the contribution of region (i) to  $I$  by  $I_i$ . By the Schwarz inequality,

$$|I_1| \leq 2 \left( \int_{|y_1| > R} dy_1 |\varphi_1(y_1)|^2 \right)^{1/2} \rightarrow 0$$

for  $R \rightarrow \infty$ , and similarly for  $I_2$ . For region (3) we shall prove below that

$$|x| \leq aR \quad \text{and} \quad |x'| \leq aR \tag{11}$$

for some constant  $a$  depending only on the masses. Therefore

$$\begin{aligned} |I_3| &\leq \int_{|x| \leq aR} dx |\eta_n(x)|^2 \leq aR \int dx |x|^{-1} |\eta_n(x)|^2 \\ &= aRn^{-2} \end{aligned}$$

since  $(\eta_n, |x|^{-1} \eta_n) = n^{-2}$  [4]. By choosing first  $R$  and then  $n$  sufficiently large, we can make  $I_1 \dots I_3$  arbitrarily small. Therefore, (10) is satisfied.

To prove (11), finally, we remark that  $(C'_1, C'_2)$  is obtained from  $(C_1, C_2)$  by exchanging groups  $g_i \subset C_i$  of an equal number of identical particles. Let  $m$  be the total mass of  $g_i$ . Then  $m < M_1$  or  $m < M_2$  since otherwise  $C_1$  and  $C_2$  would be identical, which is impossible because  $q_1 q_2 < 0$ . Let  $z_i$  be the position of the CM of  $g_i$  with respect to the CM of  $C_i$ . Then the CM of  $C'_1, C'_2$  have the following coordinates relative to the CM of  $C_1$ :

$$C'_1: mM_1^{-1}(-z_1 + x + z_2)$$

$$C'_2: x + mM_2^{-1}(-z_2 - x + z_1).$$

Taking the difference we get

$$x' = x(1 - \alpha) + \alpha(z_1 - z_2)$$

with  $\alpha = m(M_1^{-1} + M_2^{-1})$ ,  $0 < \alpha < 2$ . Now  $|y_i| \leq R$  implies  $|z_i| \leq bR$  for some  $b$  depending on the choice of the internal coordinates  $y_i$ . Therefore

$$|x'| \leq |x| |1 - \alpha| + 2\alpha bR.$$

By the same argument,

$$\begin{aligned} |x| &\leq |x'| |1 - \alpha| + 2\alpha bR \\ &\leq |x|(1 - \alpha)^2 + 2\alpha |1 - \alpha| bR + 2\alpha bR, \end{aligned}$$

or, since  $(1 - \alpha)^2 < 1$ ,

$$|x| \leq [1 - (1 - \alpha)^2]^{-1} (|1 - \alpha| + 1) 2\alpha bR$$

and similarly for  $x'$ . This proves (11).

## References

- [1] W. HUNZIKER, *Helv. Phys. Acta* 39, 451 (1966).
- [2] B. SIMON, *Helv. Phys. Acta* 43, 607 (1970).
- [3] B. SIMON, *Proc. A.M.S.*, 42, 395 (1974).
- [4] See, for example, E. U. CONDON and G. H. SHORTLEY, *Theory of Atomic Spectra* (Cambridge University Press 1957).
- [5] H. BETHE, *Quantenmechanik der Ein- und Zwei-Elektronenprobleme*, *Handbuch der Physik* XXIV/1, p. 287 (Springer Verlag, Berlin 1933).

## Notes added in proof

1. The spectrum of  $H$  in subspaces of given angular momentum, parity and symmetry under permutations of identical particles is discussed in E. Balslev, *Annals of Physics* 73, 49 (1972).
2. To derive (6) it suffices to have pointwise exponential bounds for the one-particle densities  $\rho_i$ . Such bounds have already been proved by J. M. Combes and L. Thomas, *Commun. math. Phys.* 34, 251 (1973).
3. The derivation of exponential bounds in subspaces of given symmetry presents some technical difficulties. These are discussed in W. Hunziker, *O'Connors Theorem with Statistics*, preprint 1975.
4. J. M. Combes and, independently, H. Tamura have obtained related results on the asymptotic behaviour of eigenvalues near the continuum (private communications by J. M. Combes and L. Thomas).