

Nuclear spin polarization of hydrogen by collision with optically pumped sodium

Autor(en): **Wakuta, Y. / Ogura, K. / Ueno, A.**

Objektyp: **Article**

Zeitschrift: **Helvetica Physica Acta**

Band (Jahr): **59 (1986)**

Heft 4

PDF erstellt am: **13.09.2024**

Persistenter Link: <https://doi.org/10.5169/seals-115733>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

NUCLEAR SPIN POLARIZATION OF HYDROGEN
BY COLLISION WITH OPTICALLY PUMPED SODIUM

Y. Wakuta, K. Ogura, A. Ueno, K. Takasaki, A. Iida
and K. Oohata*

Kyushu University, Nuclear Engineering Department
Fukuoka, 812 Japan

*Kyushu University, Education Department
Fukuoka, 814 Japan

ABSTRACT

Theoretical calculations have been carried out which results show that the spin-exchange collision in the low magnetic field between laser oriented sodium and hydrogen can be used to produce nuclear spin polarized hydrogen directly. It is shown that after making ~ 8 collisions, more than 90% nuclear spin polarization could be expectable. Nuclear spin polarization decreases as the applied magnetic field increases. On the contrary, electron spin polarization of hydrogen increases with the magnetic field.

1. Introduction

It is no exaggeration to mention that in low and medium energy nuclear physics, the experimental data taken with unpolarized particles are only useful for complementary to those taken with polarized particles. In order to clarify the nuclear reaction mechanism and nuclear structure, a wide variety of the polarized projectiles and targets with high polarization and intensity are indispensable. There exist, however, no single polarized source and target which can satisfy such a wide range of requirements. Therefore, "the right sources (or targets) for the right experiments" philosophy has been observed. So far, several polarization schemes have been proposed and some are brought into routine operation.

The polarized ion sources based on the optical pumping method may open a new horizon for a universal polarized ion source. High polarization and intensity of a wide variety of polarized projectiles may become available with rather low power laser. In the present paper we will present some preliminary results of theoretical calculations on the obtainable nuclear spin polarization of hydrogen by collision with the optically oriented sodium directly in the low magnetic field.

2. Method of Calculations and Results

Knize et al.[1] have shown that laser optical pumping spin-exchange between Rb and hydrogen atoms seems to be an efficient way of producing of large amount of nuclear spin polarized hydrogen with moderate laser power (~ 1 watt).

We have calculated obtainable hydrogen nuclear polarization by spin-exchange with optically orientated sodium ($\text{Na}(\uparrow) + \text{H}(\downarrow) \rightarrow \text{Na}(\downarrow) + \text{H}(\uparrow)$). For the sake of simplification of the calculations, we assumed only one substate ($F = 2$ and $m_F = +2$) of sodium 8 hyperfine structures is completely

populated by optical pumping.

Occupation probability of hydrogen substate n (j = 1,2,3 and 4) after the k collision can be given by following equation

n_j^k = sum_{i=1}^4 sum_{l=1}^8 sum_{m=1}^8 n_j^{k-1} N_l^{k-1} |<j,m| P_0 + e^{-i phi_st} P_1 | i,l >|^2

where i and j; hydrogen substates before and after the collision (i,j = 1~4),

l and m; sodium substates before and after the collision (l,m = 1~4),

P_0 and P_1; the projection operators for total spin angular momentum 0 and 1, respectively.

phi_st; relative phase between singlet and triplet parts of the wave function after the collision,

n_i^{k-1} and N_l^{k-1}; occupation probabilities after (k-1)th collision for hydrogen and sodium, respectively.

Table 1 shows the obtained results of the square of the matrix elements averaged over phi_st. Table 2 shows the results expressing the occupation probabilities of four hydrogen substates after kth collision n_j^k (j = 1~4).

Table 1. Square of the matrix elements averaged over relative phase.

A large grid table with 32 rows and 32 columns. Each cell contains a fraction representing a matrix element value. The rows are labeled with (j,m) pairs from (1,1) to (4,8). The columns are labeled with (i,l) pairs from (1,1) to (4,8). The diagonal elements are all 1. Other elements are fractions like 1/16, 3/32, 5/8, etc.

Table 2. Occupation probabilities of hydrogen substates after k^{th} collision.

j	F	m_F	n_j^k
1	1	1	$(n_1^{k-1} + \frac{1}{4}n_2^{k-1} + \frac{1}{4}n_4^{k-1}) \times N_1^{k-1}$
2	1	0	$(\frac{5}{8}n_2^{k-1} + \frac{1}{4}n_2^{k-1} + \frac{1}{8}n_4^{k-1}) \times N_1^{k-1}$
3	1	-1	$\frac{1}{2}n_3^{k-1} \times N_1^{k-1}$
4	0	0	$(\frac{1}{8}n_2^{k-1} + \frac{1}{4}n_3^{k-1} + \frac{5}{8}n_4^{k-1}) \times N_1^{k-1}$

From these results, the degree of nuclear spin polarization can be obtained and is shown in Fig.1 as a function of the number of collisions assuming $N_1 = 1$. As can be seen from this figure, nuclear polarization increases as the result of successive collision with optically oriented sodium.

Fig.2 shows another interesting results obtained from the present calculations. Nuclear spin polarization of hydrogen becomes almost zero as the applied magnetic field increases. On the contrary, the electron polarization of hydrogen approaches more than 90% after several collisions with sodium as the applied external magnetic field is increased.




Fig.1 Nuclear spin polarization as a function of the number of collisions.




Fig.2 Nuclear spin and electron spin polarization of hydrogen atom as a function of the magnetic field. P_N ; nuclear spin polarization and P_e ; electron spin polarization.

3. Summary

In conclusion, we can expect very high nuclear spin polarization directly through the spin-exchange collision if the time between spin-exchange collision is short compared to either the spin lattice relaxation time or the pumping time for the system. Furthermore, owing to the high spin-exchange cross section, we will be able to obtain highly intense nuclear spin polarized hydrogen by using relatively low power laser.

References

- [1] R.J. Knize, W. Happer and J.L. Cecchi,
Optically Pumping Production of Spin Polarized Hydrogen
PPPL - 2125(1984), Princeton University