

Time-delay operator for a class of singular potentials

Autor(en): **Wang, Xue-Ping**

Objektyp: **Article**

Zeitschrift: **Helvetica Physica Acta**

Band (Jahr): **60 (1987)**

Heft 4

PDF erstellt am: **11.09.2024**

Persistenter Link: <https://doi.org/10.5169/seals-115860>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Time-delay operator for a class of singular potentials

By Xue-Ping Wang*

Institut de Mathématiques, Université de Nantes, 44072 Nantes Cedex, France

(1. IX. 1986)

Abstract. We prove the existence of the time-delay operator defined by taking the large space limit of the approximate sojourn times for a class of singular potentials: $V = V_1 + V_2$, where V_1 is a smooth short range potential and V_2 and $x \cdot \nabla V_2$ are both bounded from H^2 to $L^{2,2+\epsilon_0}$ for some $\epsilon_0 > 0$.

1. Introduction

In [8], we proved the finiteness of time-delay defined by taking the space limit of sojourn times and established its equivalence with Eisenbud–Wigner’s time-delay in scattering theory for smooth short range potentials. In this work we will show that our method developed there can be also applied to a class of singular potentials.

Let $H_0 = -\Delta$ and $H = H_0 + V$ in $L^2(\mathbb{R}^n)$. We suppose that the short range potential V can be decomposed as: $V(x) = V_1(x) + V_2(x)$ where V_1 is C^∞ on \mathbb{R}^n and for some $\epsilon_0 > 0$

$$|\partial_x^\alpha V_1(x)| \leq c_\alpha \langle x \rangle^{-1-\epsilon_0-|\alpha|} \tag{1.1}$$

and the multiplication by V_2 is bounded as operator from H^2 to $L^{2,2+\epsilon_0}$ and so is the distributional derivative $x \nabla_x \cdot V_2$. Here $\langle x \rangle = (1 + |x|)$ and H^s is the usual Sobolev space of order s ; $L^{2,s}$ is the weighted L^2 space with the norm: $\|f\|_s = \|\langle x \rangle^s f\|$. This assumption will be made throughout this work. Under this condition on V , it is well known that the wave operators W_\pm defined by:

$$W_\pm = s - \lim_{t \rightarrow \pm\infty} e^{itH} e^{-itH_0} \text{ in } L^2$$

exist and are complete. Let \tilde{P}_R denote the multiplication by the characteristic function for the ball $\{|x| < R\}$. Then the local time-delay of f in $\{|x| < R\}$ is defined as the difference of the sojourn times:

$$\langle f, T_R f \rangle = \int_{-\infty}^{\infty} (\|\tilde{P}_R e^{-itH} W_- f\|^2 - \|\tilde{P}_R e^{-itH_0} f\|^2) dt \tag{1.2}$$

* Address after February 1st, 1987: Institute of Mathematics, University of Pekin, Pekin, China.

Notice that (1.2) is well defined for $f \in L^2$ such that the Fourier transform \hat{f} has suitable compact support in $\mathbb{R}^n \setminus \{0\}$. Finally the time-delay operator T is defined by:

$$\langle f, Tf \rangle = \lim_{R \rightarrow +\infty} \langle f, T_R f \rangle \tag{1.3}$$

whenever the limit exists. Surely the existence of time-delay operator depends on how much such f 's we can find. As in [8], we consider here a similar question. Let P_R be the multiplication by $P(x/R)$, where $P(\cdot)$ is a smooth, spherically symmetrical function such that $P(x) = 1$ for $|x| \leq 1$ and $P(x) = 0$ for $|x| \geq 2$. It is clear that P_R can be regarded as an approximation of \tilde{P}_R . In the following we denote still T_R the operator defined by (1.2) with \tilde{P}_R replaced by P_R .

Then for smooth short range potentials we proved in [8] that the limit (1.3) exists for a dense subset in L^2 and in the spectral representation of H_0 , the time-delay operator T is given by a family of operators $T(\lambda)$, $\lambda > 0$, where

$$T(\lambda) = -iS(\lambda)^* \frac{d}{d\lambda} S(\lambda) \tag{1.4}$$

$S(\lambda)$ being the scattering matrix. (1.4) is the Eisenbud–Wigner’s formula for time-delay. It reveals that the method and techniques used in [8] are powerful enough. It can be applied to treat time-delay in other scattering theories (see [6]) and to include a class of singular potentials.

Let $A = -i(x \cdot \nabla_x + \nabla_x x)/2$ be the generator of dilation group. We define the set \mathcal{D} by:

$$\mathcal{D} = \{f \in L^2; f \in D(\langle x \rangle) \cap D(A^2) \text{ and } \exists \chi \in C_0^\infty(\mathbb{R}_+ \setminus \sigma_p(H)), \chi(H_0)f = f\} \tag{1.5}$$

In this work we want to prove the following result.

Theorem 1. *Under the above assumption on V , the limit (1.3) exists for T_R defined by (1.2) with \tilde{P}_R replaced by P_R and for $f \in \mathcal{D}$. We have:*

$$\langle f, Tf \rangle = \langle f, -S^*[A, S]f_1 \rangle$$

where f_1 is determined by $2H_0f_1 = f$ and $S = W_+^*W_-$ is scattering operator. The time-delay operator T is essentially selfadjoint with core \mathcal{D} and the Eisenbud–Wigner formula (1.4) is true for $\lambda \in \mathbb{R}_+ \setminus \sigma_p(H)$.

The proof of this result consists in regarding V_2 as a perturbation of the Hamiltonian $H_1 = H_0 + V_1$. In §2, we give some technical preparations, which were mostly proved in [8]. In §3, we achieve the main step of the proof, reducing the existence of the limit (1.3) to that of $\lim_{R \rightarrow +\infty} \int_0^\infty \langle U_0(t)f, S^*P_R S - P_R \rangle f \rangle dt$. We finish the proof of Theorem 1 in §4 by the method of [8]. Very recently Nakamura ([12]) considered the similar problem by a different method. His proof is in the spirit of Lavine [4], while ours is in that of Martin [5].

2. Some preparations

Let $U(t)$ (resp. $U_0(t), U_1(t)$) denote the unitary group associated to H (resp. H_0, H_1). Let $E_{ac}(H_1)$ denote the spectral projector onto the absolute continuous space of H_1 . The wave operators W_{\pm}^1, W_{\pm}^2 are defined by:

$$W_{\pm}^1 = s\text{-}\lim_{t \rightarrow \pm\infty} U_1(t)^* U_0(t)$$

$$W_{\pm}^2 = s\text{-}\lim_{t \rightarrow \pm\infty} U(t)^* U_1(t) E_{ac}(H_1)$$

in $L^2(\mathbb{R}^n)$. By chain rule, $W_{\pm} = W_{\pm}^2 W_{\pm}^1$ (see [1]). Put: $\mathbb{R}_+ =]0, +\infty[$.

Lemma 2.1. *Let $f \in C_0^\infty(\mathbb{R}_+/\sigma_p(H))$. Then for every $0 \leq \mu \leq 1$, one has:*

$$\|\langle A \rangle^{-\mu} f(H) U(t) W_{\pm} \langle A \rangle^{-\mu}\| \leq C(1 + |t|)^{-\mu} \tag{2.1}$$

for $t \in \mathbb{R}$. For every $\mu > 1$, there exists $\rho > 1$ such that

$$\|\langle A \rangle^{-\mu} f(H) U(t) W_{\pm} \langle A \rangle^{-\mu}\| \leq c(1 + |t|)^{-\rho} \text{ for } t \in \mathbb{R} \tag{2.2}$$

Note that this result is proved in [8] for $V = V_1(V_2 = 0)$. But the proof can be carried over, because we used only the short range properties of V and $x \cdot \nabla V$.

Recall that if $V = V_1 + V_2$ with V_1 satisfying (1.1) and V_2 bounded from H^2 to $L^{2,2+\varepsilon_0}$ it is proved in [3] that $S(\lambda)$ is continuously differentiable in $\mathcal{L}(L^2(S^{n-1}))$ for $\lambda \in \mathbb{R}_+/\sigma_p(H)$. Since under the assumptions of Theorem 1, $x \cdot \nabla V$ satisfies still the above conditions, it should be clear that by exterior scaling method, we can easily prove that $S(\lambda)$ is two times differentiable for $\lambda \in \mathbb{R}_+/\sigma_p(H)$. For reader's convenience we give the details of the proof.

Let $\lambda \in \mathbb{R}_+/\sigma_p(H)$. Then we have the following representation for the scattering matrix $S(\lambda)$ ($= S(\lambda, V)$) ([11]):

$$S(\lambda, V) = 1 - i\pi \mathcal{F}(\lambda)(V - VR(\lambda + i0, V)V) \mathcal{F}(\lambda)^*$$

where $R(\lambda \pm i0, V)$ is the boundary values of the resolvent $(H_0 + V - z)^{-1}$ and $\mathcal{F}(\cdot): L^2(\mathbb{R}^n) \mapsto L^2(\mathbb{R}_+, L^2(S^{n-1}))$ is a spectral representation for the free Hamiltonian H_0 . Take $a > 0$ to be sufficiently small. Put: $I =]-a, a[$. We can prove that:

$$S(e^{2\theta}\lambda, V) = S(\lambda, V(\theta)) \text{ for } \theta \in I \tag{S}$$

where $V(\theta) = e^{-2\theta}U(\theta)^* V U(\theta)$ and $U(\theta)$ is the unitary group generated by A . Now we check the derivability of $V(\theta)$ and $R(\lambda + i0, V(\theta))$ for $\theta \in I$. Let $H^{s,m}$ denote the weighted Sobolev space with the norm $\|\langle x \rangle^m (1 - \Delta)^{s/2} f\|$. Put: $\rho = 1 + \varepsilon_0 > 1$. Then the assumptions on V say that $i[A, V_2]$ defines a bounded operator from $H^{s,r}$ to $H^{s-2, r+\rho+1}$ for $0 \leq s \leq 2$ and $r \in \mathbb{R}$. From this we derive that $A[A, V_2]$ and $[A, V_2]A$ are both bounded from $H^{s,r}$ to $H^{s-4, r+\rho}$ for $0 \leq s \leq 3$ and $r \in \mathbb{R}$. Since V_1 satisfies (1.1), we conclude easily from the above remarks that the operator valued function $\theta \mapsto V(\theta)$ is in the class

$$C^1(I; \mathcal{L}(H^{s,r}; H^{s-2, r+\rho+1})) \cap C^2(I; \mathcal{L}(H^{s+1, r}; H^{s-3, r+\rho}))$$

Since $R(\lambda \pm i0, V(\theta))$ is in $\mathcal{L}(H^{0,r}; H^{2,-r})$ if $r > \frac{1}{2}$ ([10]), we can also prove that the map $\theta \mapsto R(\lambda + i0, V(\theta))$ belongs to the class:

$$C^1(I; \mathcal{L}(H^{0,r}; H^{2,-r})) \cap C^2(I; \mathcal{L}(H^{0,r}; H^{0,-r}))$$

for $r > \frac{1}{2}$. This means that the map

$$I \ni \theta \mapsto \langle H_0 \rangle^{-2} V(\theta) (1 - R(\lambda + i0, V(\theta)) V(\theta)) \langle H_0 \rangle^{-2}$$

is in $C^2(I; \mathcal{L}(L^{2,-s}; L^{2,s}))$ for $s \in]\frac{1}{2}, \rho/2[$. Making use of the relation: $\mathcal{F}(\lambda) \langle H_0 \rangle^{-1} = \langle \lambda \rangle^{-1} \mathcal{F}(\lambda)$, we derive from (S) that $S(\lambda, V)$ is twice continuously differentiable in $\mathcal{L}(L^2(S^{n-1}))$ for $\lambda \in \mathbb{R}_+ / \sigma_p(H)$. This proves our assertion.

Since in the spectral representation of H_0 , A is given by a family of operators $A(\lambda) = i(\lambda d/d\lambda + d/d\lambda \cdot \lambda)$, we conclude that the domain of A^2 is invariant by $Sf(H_0)$ for $f \in C_0^\infty(\mathbb{R}_+ / \sigma_p(H))$. Now we can easily prove the following lemma which is important in this work.

Lemma 2.2. *Let \mathcal{D} be defined by (1.5). Then \mathcal{D} is invariant by S . In particular if f belongs to \mathcal{D} , $\langle x \rangle Sf$ and $A^2 Sf$ are both in $L^2(\mathbb{R}^n)$.*

Proof. It remains to show that $\langle x \rangle Sf$ is in L^2 . We can use the same commutator method as in the proof of Prop. 4.2 in [8]. The details are omitted here.

In order to regard V_2 as a perturbation to H_1 , we need some continuity of wave operators W_\pm^1 .

Lemma 2.3. *Let $f \in C_0^\infty(\mathbb{R}_+)$. Under the condition (1.1) on V_1 , the four operators $A^2 W_\pm^1 f(H_0) \langle A \rangle^{-2}$ and $A^2 W_\pm^{1*} f(H_1) \langle A \rangle^{-2}$ are all bounded on L^2 .*

Proof. We prove only the result for $A^2 W_+^{1*} f(H_1) \langle A \rangle^{-2}$. The other cases can be treated in the same way. Put $W(t) = U_0(t)^* U_1(t)$. We can write, as forms on $D(A) \times D(A)$,

$$\begin{aligned} AW(t)f(H_1) &= W(t)f(H_1)A + U_0(t)^*[A, f(H_1)]U_1(t) \\ &\quad + 2tU_0(t)V_1U_1(t)f(H_1) + W(t) \int_0^t U_1(s)^* f(H_1) \tilde{V} U_1(s) ds \end{aligned} \tag{2.3}$$

where $\tilde{V} = i[A, V_1] - 2V_1$. Notice that $i[A, f(H_1)] = 2f'(H_1) + Q$ with Q bounded from L^2 to $L^{2,1+\varepsilon_0}$ (see [8]). Now we need the following result due to Jensen et al.:

$$\| \langle A \rangle^{-r} f(H_1) U_1(t) \langle A \rangle^{-r} \| \leq c(1 + |t|)^{-r+\varepsilon} \quad t \in \mathbb{R} \tag{2.4}$$

for every $r > 0$ and $0 < \varepsilon \ll r$. Take $g \in C_0^\infty(\mathbb{R}_+)$ such that $g = 1$ on $\text{supp} f$. Multiplying (2.3) by $g(H_1) \langle A \rangle^{-2}$ and taking the limit $t \rightarrow +\infty$, applying (2.4), we

get:

$$\begin{aligned}
 AW_+^{1*}f(H_1)\langle A \rangle^{-2} &= W_+^{1*}f(H_1)Ag(H_1)\langle A \rangle^{-2} + W_+^{1*}h(H_1)\langle A \rangle^{-2} \\
 &\quad + W_+^{1*} \int_0^{+\infty} f(H_1)U(-s)\tilde{V}U(s)g(H_1)\langle A \rangle^{-2} dt \tag{2.5}
 \end{aligned}$$

where $h = 2if'g$. Since $AW_+^{1*}f(H_1)\langle A \rangle^{-1}$ is bounded on L^2 , in order to prove the desired result by (2.5), it is sufficient to show that

$$\int_0^{+\infty} Af(H_1)U_1(-s)\tilde{V}U_1(s)g(H_1)\langle A \rangle^{-2} ds \tag{2.6}$$

is bounded on L^2 . To simplify notations, we denote f, g the operators $f(H_1), g(H_1)$ respectively. We have the following relation:

$$\begin{aligned}
 [A, U_1(-s)g\tilde{V}gU_1(s)] &= -2sU_1(-s)g[H_0, \tilde{V}]gU_1(s) \\
 &\quad + U_1(-s)[A, g\tilde{V}g]U_1(s) \\
 &\quad - \int_0^s U_1(t-s)\tilde{V}gU_1(-t)\tilde{V}U_1(s)g dt \\
 &\quad + \int_0^s U_1(-s)g\tilde{V}U_1(s-t)g\tilde{V}U_1(t) dt \tag{2.7}
 \end{aligned}$$

Since $g[H_0, \tilde{V}]$ is continuous from $L^{2,r}$ to $L^{2,r+2+\epsilon_0}$, we can prove as in [8] (Prop. 4.2) that $\int_0^{+\infty} 2sfU_1(-s)[H_0, \tilde{V}]U_1(s)g\langle A \rangle^{-1} ds$ is bounded on $L^2(\mathbb{R}^n)$. Since $[A, g\tilde{V}g]$ is bounded as operator from $L^{2,r}$ to $L^{2,r+1+\epsilon_0}$, it follows from (2.4) that:

$$\|[A, g\tilde{V}g]U_1(s)g_1\langle A \rangle^{-2}\| \leq C(1 + |s|)^{-1-\epsilon_0/2}$$

for $s \in \mathbb{R}$. Here $g_1 = g_1(H_1)$ is chosen so that $g_1g = g$. Therefore the integral $\int_0^\infty fU_1(-s)[A, g\tilde{V}g]U_1(s)g_1\langle A \rangle^{-2} ds$ defines a bounded operator on L^2 . To treat the last two terms in (2.7), we use the local H_1 -smoothness of $\langle x \rangle^{-1/2-\epsilon}$, which implies that the operator $\int_0^s fU_1(t)\tilde{V}U_1(-t)g dt$ is uniformly bounded with respect to $s \in \mathbb{R}$. Applying (2.4), we get the estimate over the third term in (2.7):

$$\left\| \int_0^s fU_1(t-s)\tilde{V}gU_1(-t)\tilde{V}U_1(s)g_1\langle A \rangle^{-2} dt \right\| \leq C(1 + |s|)^{-1-\epsilon_0/2}$$

The last term in (2.7) can be estimated in the same way. Since the commutator $[A, f]$ is bounded, we derive from (2.7) that (2.6) is a bounded operator on L^2 . This proves that $A^2W_+^{1*}f(H_1)\langle A \rangle^{-2}$ is bounded. The lemma is proved.

3. Reduction of the problem

In the proof of the finiteness of time-delay, an important step is to show that

$$\lim_{R \rightarrow +\infty} \left(\langle f, T_R f \rangle - \int_0^{+\infty} \langle U_0(t)f, (S^*P_R S - P_R)U_0(t)f \rangle dt \right) = 0 \tag{3.1}$$

(3.1) makes also clear the close relationship between time-delay operator T and scattering operator S . In this section we will prove the following result which implies (3.1).

Theorem 3.1. *Let $f \in \mathcal{D}$. Put $g = Sf$. Then we have:*

$$\lim_{R \rightarrow +\infty} \int_{-\infty}^0 (\langle U_0(t)f, (W_-^* P_R W_- - P_R)U_0(t)f \rangle) dt = 0 \tag{3.2}$$

$$\lim_{R \rightarrow +\infty} \int_0^{\infty} (\langle U_0(t)g, (W_+^* P_R W_+ - P_R)U_0(t)g \rangle) dt = 0 \tag{3.3}$$

Proof. Since the set \mathcal{D} is invariant by S , it suffices to prove (3.2). Put $h = W_-^1 f$, which is in $D(A^2)$ by Lemma 2.3. The integrand in (3.2) can be written as: $\langle U_1(t)h, (W_-^{2*} P_R W_-^2 - P_R)U_1(t)h \rangle + \langle U_0(t)f, (W_-^{1*} P_R W_-^1 - P_R)U_0(t)f \rangle$. It is proved in [8] by method of pseudo-differential operators that for $f \in \mathcal{D}$, we have:

$$\lim_{R \rightarrow +\infty} \int_{-\infty}^0 \langle U_0(t)f, (W_-^{1*} P_R W_-^1 - P_R)U_0(t)f \rangle dt = 0$$

Therefore we have to prove

$$\lim_{R \rightarrow +\infty} \int_{-\infty}^0 \langle U_1(t)h, (W_-^{2*} P_R W_-^2 - P_R)U_1(t)h \rangle dt = 0 \tag{3.4}$$

The integrand in (3.4) can be written as:

$$\begin{aligned} &\langle P_R U(t)W_- f, (U(t)W_-^2 - U_1(t))h \rangle \\ &+ \langle (U(t)W_-^2 - U_1(t))h, P_R U_1(t)h \rangle \end{aligned} \tag{3.5}$$

Take $\chi \in C_0^\infty(\mathbb{R}_+ / \sigma_p(H))$ such that $\chi(H_0)f = f$. We get:

$$\begin{aligned} (U(t)W_-^2 - U_1(t))h &= - \int_{-\infty}^t \chi(H)U(t-s)V_2 U_1(s)\chi(H_1)h ds \\ &+ (\chi(H) - \chi(H_1))U_1(t)h \end{aligned}$$

By the assumption, $V_2 \chi(H_1)$ is continuous from $L^{2, -\epsilon_0}$ to $L^{2, 2}$. We can easily prove that $\chi(H) - \chi(H_1)$ is continuous from $L^{2, -1-\epsilon_0}$ to L^2 . Thus the first term in (3.5) can be estimated as:

$$\begin{aligned} &|\langle P_R U(t)W_- f, (U(t)W_-^2 - U_1(t))h \rangle| \\ &\leq c \int_{-\infty}^t (1 + |s|)^{-2+\epsilon} \|\langle x \rangle^{-\epsilon_0} \chi(H)U(s-t)P_R U(t)\chi(H)W_- f\| \|\langle A \rangle^2 f\| ds \\ &+ c(1 + |t|)^{-1-\epsilon_0} \|\langle A \rangle^2 f\|^2, \quad \text{for any } \epsilon > 0 \end{aligned} \tag{3.6}$$

Here we have used (2.4) and Lemma 2.3. Before going on with the proof of Theorem 3.1, we need still a lemma.

Lemma 3.2. For every $0 \leq \mu \leq 1$, we have:

$$\|\langle x \rangle^{-\mu} \chi(H) U(s-t) P_R U(t) W_- f\| \leq C(1+|s|)^{-\mu} \|\langle A \rangle W_- f\| \tag{3.7}$$

uniformly in $t \in \mathbb{R}$ and $R \geq 1$.

Proof. Observe first that $|\partial_x^\alpha P_R(x)| \leq c \langle x \rangle^{-|\alpha|}$ uniformly in $R \geq 1$. By the arguments used in the proof of Lemma 2.3, we can show that:

$$\|A \chi(H) U(-t) P_R U(t) \chi(H) \langle A \rangle^{-1}\| \leq C$$

uniformly in $t \in \mathbb{R}$ and $R \geq 1$. (3.7) follows from (2.1) and the fact that $\langle A \rangle \chi(H) \langle x \rangle^{-1}$ is bounded on L^2 .

Now return to the proof of Theorem 3.1. By (3.6) and (3.7) we obtain, for $\varepsilon > 0$ sufficiently small and for $t < 0$,

$$|\langle P_R U(t) W_- f, (U(t) W_-^2 - U_1(t)) h \rangle| \leq c(1+|t|)^{-1-\varepsilon/2} \|\langle A \rangle^2 f\|^2$$

uniformly in $R \geq 1$. The same estimate is also true for the second term in (3.5). Since the integrand in (3.4) tends to 0 as R tends to $+\infty$, by the dominated convergence theorem, (3.4) is proved. This finishes the proof of Theorem 3.1.

We remark that the various constants c appeared in the proof of Theorem 3.1 depend on the function χ , but not $f \in \mathcal{D}$ such that $\chi(H_0)f = f$.

4. Existence of time-delay operator

In this section we prove Theorem 1. Let $\chi \in C_0^\infty(\mathbb{R}_+/\sigma_p(H))$. We put:

$$\mathcal{D}_\chi = \{f \in \mathcal{D}; \chi(H_0)f = f\}$$

Lemma 4.1. Let $f \in \mathcal{D}_\chi$. Then,

$$\int_{R^{5/2}}^{+\infty} |\langle U_0(t)f, (S^* P_R S - P_R) U_0(t)f \rangle| dt \leq C R^{-1/2} \|\langle A \rangle f\|^2$$

Proof. It follows easily from the estimate:

$$\|P_R^{1/2} U_0(t)f\| \leq C_\chi R(1+|t|)^{-1} \|\langle A \rangle f\|$$

for $t \in \mathbb{R}$, $R \geq 1$ and $f \in \mathcal{D}_\chi$.

Lemma 4.2. For $f \in \mathcal{D}_\chi$, we have the asymptotic expansion:

$$\begin{aligned} \int_0^{R^{5/2}} \langle U_0(t)f, P_R U_0(t)f \rangle dt &= R \langle f, a(D)f \rangle \\ &- \langle f, b^w(x, D; \chi)f \rangle + O(R^{-1/2}) \end{aligned} \tag{4.1}$$

where $a(\xi) = c_0 |\xi|^{-1} \chi(|\xi|^2)$ with $c_0 = \frac{1}{2} \int_0^\infty P(s) ds$, $b^w(x, D; \chi)$ is a Weyl pseudo-differential operator with symbol $\frac{1}{2} x \xi |\xi|^{-2} \chi(|\xi|^2)$. The remainder $O(R^{-1/2})$ can be

estimated by

$$|O(R^{-1/2})| \leq C_x R^{-1/2} (\|\langle x \rangle f\| + \|\langle A \rangle^2 f\|)^2 \quad R \geq 1$$

Proof. We use the fact that $U_0(-t)P_R U_0(t)$ is a Weyl pseudo-differential operator with symbol $P((x + 2t\xi)/R)$ and develop the symbol around $2t\xi/R$. Since this result is proved in [8] (Proposition 4.3) for $f \in D(\langle x \rangle) \cap D(\langle A \rangle^3)$, we indicate only the difference and omit the details. Checking the proof of Proposition 4.3([8]), we see that the condition $f \in D(\langle A \rangle^3)$ is only used to get the estimate (see (4.14)[8]):

$$\frac{t}{R^3} \|Q^w(\tau x/R, D)U_0(t/\tau)f\| \leq CR^{-1/2}t^{-3/2} \|\langle A \rangle^3 f\|$$

for $t \geq 1, \tau \in]0, 1]$ and $R \geq 1$, where $Q(x, \xi) = P_1(x)\chi(|\xi|^2)$, $P_1(x)$ is some derivative of $P(x)$. Hence it is supported in $\{1 \leq |x| \leq 2\}$. But this term can be equally estimated as follows: Since all the derivatives of $\tau^2 R^{-2} Q(\tau x/R, \xi)$ are bounded by a constant times $\langle x \rangle^{-2}$ uniformly with respect to $R \geq 1$ and $\tau \in]0, 1]$, we have, by continuity result of pseudo-differential operators ([2]),

$$\begin{aligned} \frac{t}{R^3} \|Q^w(\tau x/R, D)U_0(t/\tau)f\| \\ \leq cR^{-1}\tau^{-2}t \|\langle x \rangle^{-2}U_0(t/\tau)f\| \leq c_x R^{-1}t^{-1} \|\langle A \rangle^2 f\| \end{aligned} \tag{4.2}$$

for $t \geq 1, R \geq 1$ and $\tau \in]0, 1]$. Integrating (4.2) over $[1, R^{5/2}]$, we get:

$$\int_0^{R^{5/2}} tR^{-3} \|Q^w(\tau x/R, D)U_0(t/\tau)f\| dt \leq C_x R^{-1/2} \|\langle A \rangle^2 f\|$$

This gives the desired result. The lemma is proved.

Now we are able to give the proof of Theorem 1.

Proof of Theorem 1. Let f be in \mathcal{D} . Then Sf is also in \mathcal{D} . Take $\chi \in C_0^\infty(\mathbb{R}_+/\sigma_p(H))$ such that $\chi(H_0)f = f$. Since S commutes with $a(D)$, we derive from Lemmas 4.1 and 4.2 that

$$\begin{aligned} \left| \int_0^\infty \langle U_0(t)f, S^*[P_R, S]U_0(t)f \rangle dt + \langle f, S^*[b^w(x, D; \chi), S]f \rangle \right| \\ \leq C_f R^{-1/2} \end{aligned} \tag{4.3}$$

By a simple calculus of Weyl pseudo-differential operators ([2]), we get:

$$\langle f, S^*[b^w(x, D; \chi), S]f \rangle = \langle f_1, S^*[A, S]f \rangle \tag{4.4}$$

where $f_1 = (2H_0)^{-1}\chi(H_0)f$. Theorem 1 is a consequence of (4.3) and (4.4). See also [8].

REFERENCES

[1] W. O. AMREIN, J. M. JAUCH and K. D. SINHA, *Scattering Theory in Quantum Mechanics*, Reading, Benjamin Inc., 1977.

- [2] L. HÖRMANDER, *The Weyl calculus of pseudo-differential operators*, Comm. Pure Appl. Math., 32 (1979), 359–443.
- [3] A. JENSEN, *A stationary proof of Lavine's formula for time-delay*, Lett. in Math. Phys., 7 (1983), 137–143.
- [4] R. LAVINE, *Commutators and local decay*, in *Scattering Theory in Mathematical Physics*, pp. 144–156, ed. J. A. La Vita, J. P. Marchand; Reidel, Dordrecht, 1974.
- [5] PH. MARTIN, *On the time-delay of simple scattering systems*, Commun. Math. Phys., 47 (1976), 221–227.
- [6] D. ROBERT, X. P. WANG, *Time-delay and spectral density for Stark Hamiltonians*, in preparation.
- [7] X. P. WANG, *Opérateurs de temps-retard dans la théorie de la diffusion*, C.R. Acad. Sci. Paris, Sér. I, 301 (1985), 789–791.
- [8] X. P. WANG, *Phase-space description of time-delay in scattering theory*, to appear.
- [9] X. P. WANG, *Low energy resolvent estimates and continuity of time-delay operators*, to appear in Proc. Royal Soc. Edinburgh.
- [10] S. AGMON, *Spectral properties of Schrödinger operators and scattering theory*, Ann. Scuol. Norm. Pisa, 4 (1975), 151–218.
- [11] S. T. KURODA, *Scattering theory for differential operators, I, operator theory*, J. Math. Soc. Japan, 25 (1973), 75–104.
- [12] S. NAKAMURA, *Time-delay and Lavine's formula*, preprint.