

Zeitschrift: Helvetica Physica Acta

Band: 62 (1989)

Heft: 6-7

Artikel: Dissipative mechanics in granular superconductors

Autor: Fazio, R. / Falci, G. / Giaquinta, G.

DOI: <https://doi.org/10.5169/seals-116088>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Siehe Rechtliche Hinweise.

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. See Legal notice.

Download PDF: 29.04.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

DISSIPATIVE MECHANISMS IN GRANULAR SUPERCONDUCTORS

R.Fazio, Faculty of Applied Physics, Delft University, 2600 GA DELFT, The Netherlands, - perm. address : Istituto di Fisica, CATANIA.
G. Falci, Dottorato di Ricerca in Fisica, Universita' di Catania.
G. Giaquinta, Istituto di Fisica, Facolta' di Ingegneria, Universita' di Catania, Viale A.Doria 6, 95129 CATANIA, Italy

Abstract: The role played by dissipation in the coherent transition of granular superconductors is approached from a new viewpoint stressing the link between local and global properties of the system. A novel kind of free energy functional is presented.

1. Introduction

In accounting for the role of dissipation in the physics of a single Josephson Junction(SJJ), it is well known that starting from different dissipative mechanisms we are led to a non completely overlapping understandings of the superconductor-insulator transition, which rests, moreover, on limiting values of some constraints[1]. At present the extension to arrays does not look to be equally well developed and to this end the knowledge gained in the theory of granular superconductors(GS)[2] and related experiments[3] offers as a promising starting point. Indeed, in a GS, coherence results from the competition between the Josephson coupling, that determines the ordering of the phases of the single grain superconductive order parameters(singled out by a non vanishing value of an ordering field Ψ) and the electrostatic effects responsible for quantum fluctuations, dissipative tunneling suppressing this last leading to coherence.

It is our purpose to investigate the mutual interplay between the SJJ and the global transitions, stressing the role of dissipation in the former, by a novel kind of free energy accounting on the same footing for the respective ordering fields.

2. The Free Energy functional

The Euclidean effective phase action obtained in [2] describing a Josephson array exhibits the feature that, besides the spatial Josephson coupling between the single grain phases ϕ_i , dissipation couples on the imaginary time axis the differences of the phases ϕ_{ij} across every SJJ. The spatial decoupling is performed by introducing the Hubbard-Stratenovich field $\Psi_i(\tau)$ conjugated to $\exp(i\phi_i)$. We introduce a novel field $\Phi_{ij}(\tau)$, in order to perform "temporal" decoupling, conjugated with $\exp(i\phi_{ij})$. Integrating out the phase degrees of freedom, after a cumulant expansion in power of the two fields, we get to the following free energy functional form:

$$\begin{aligned} F[\Psi_i, \Phi_{ij}] = & \sum_{ij} \sum_n |\Psi_i(\omega_n) K_{ij}^{-1} \Psi_j(\omega_n) - g(\omega_n) \delta_{ij} \Psi_i(\omega_n) \Psi_j(\omega_n)|^2 + \\ & \sum_{ij} \sum_n ||\Phi_{ij}(\omega_n)||^2 - 2\alpha(\omega_n) g(\omega_n) |\Phi_{ij}(\omega_n)|^2 + \\ & F_1(\Psi^4) + F_2(\Phi^4) + F_3(\Psi^2 \Phi^2) \end{aligned}$$

where K_{ij} is the Josephson matrix and g is the phase-phase correlation function (details are omitted for brevity, they will appear in a forthcoming paper [4]). The fourth order coupling term $F_3(\Psi^2 \Phi^2)$ provides to the aimed link in the sense that it can be viewed as a second order term for the effective free energy describing the Ψ or the Φ transition as alone. The structure of the involved propagators obliges recursion to the RG methodology in accounting for the Φ transition [4] whereas a mean-field approximation is known to lead to reasonable findings as about the phase diagram of the Ψ transition.

3. References

- [1] S.A. Bulgadaev, JETP Lett. 45, 623 (1987).
- [2] U. Eckern, G. Schön and V. Ambegaokar, Phys. Rev. B30, 6914 (1984).
R. Fazio, G. Giaquinta and C. Di Mauro, Progr. Teor. Ph., 76, 592 (1986).
R. Fazio and G. Giaquinta, Phys. Rev. B 34, 4909 (1986).
G. Falci, R. Fazio and G. Giaquinta, Phys. Rev. B, 1 May 1989.
- [3] B.G. Orr et al., Phys. Rev. Lett. 56, 378 (1986).
- [4] R. Fazio, G. Falci and G. Giaquinta, to be published.