

Zeitschrift: Mycologia Helvetica

Band: 2 (1986-1987)

Heft: 1

Artikel: Zelleromyces stephensii, an interesting member of the gasteroid Russulales from Europe

Autor: Miller, Steven L. / Miller, Orson K.

DOI: <https://doi.org/10.5169/seals-1036406>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 09.11.2024

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

MYCOLOGIA HELVETICA

Vol. 2 No 1

pp. 59—66

1986

(Manuscript received on 20th February 1986)

ZELLEROMYCES STEPHENSII AN INTERESTING MEMBER OF THE GASTEROID RUSSULALES FROM EUROPE¹

Steven L. Miller

Department of Forest Science, Oregon State University
Corvallis, Oregon 97331 U.S.A.

and

Orson K. Miller, Jr.

Department of Biology, Virginia Polytechnic Institute and State
University, Blacksburg, Virginia 24061 U.S.A.

Abstract: *Zelleromyces stephensii* was collected near Etzel, Switzerland. Taxonomic notes on morphological features, distribution, and vascular plant associates are provided. *Zelleromyces stephensii* is compared with other species of *Zelleromyces* from North America.

Résumé: *Zelleromyces stephensii* a été récolté en Suisse près d'Etzel dans le canton de Schwitz. Sa morphologie, sa distribution géographique, son association avec des plantes vasculaires et sa taxonomie sont discutées ici. *Zelleromyces stephensii* est comparé avec des espèces nordaméricaines du genre *Zelleromyces*.

Zusammenfassung: *Zelleromyces stephensii* wurde in der Schweiz auf dem Etzel, Kanton Schwyz gefunden. Die Morphologie des Pilzes, seine geographische Verbreitung, Vergesellschaftung mit Gefäßpflanzen sowie seine Taxonomie werden besprochen. *Zelleromyces stephensii* wird mit weiteren nordamerikanischen Vertretern der Gattung *Zelleromyces* verglichen.

¹Funded in part by the USDA Forest Service, Pacific Northwest Research Station, Corvallis, Oregon 97331 U.S.A.

Hypogeous fungi form sporocarps at or below the surface of the ground and are therefore often rarely observed or collected. Taxonomic notes on hypogeous fungi, including fresh morphological features, distribution, associated soil types and vascular plant associates, are irreplaceable in establishing a proper systematic perspective for these fungi. Recently *Zelleromyces stephensii* (Berk.) A. H. Smith was collected by the junior author near Etzel, Switzerland. Although known throughout Europe, it appears that *Z. stephensii* has only been reported briefly in the literature from Switzerland (Froidevaux and Schwärzel 1977). The present paper provides detailed descriptive information on *Z. stephensii* in order to more carefully characterize this species of gasteroid Russulales. It is hoped that additional ecological information on the species, including fruiting period, host range and associated soil type, will result from data presented here.

Colour nomenclature and coding is taken from the Methuen Handbook of Colour (Kornerup and Wanscher 1981). Herbarium names are abbreviated according to Lanjouw and Stafleu (1964).

Zelleromyces stephensii (Berk.) A. H. Smith, Mycologia 54: 635 (1962).

Hydnangium stephensii Berk., Ann. Mag. Nat. Hist. 13: 352 (1844).

Octaviania stephensii (Berk.) Tul., Fung. Hypog. 78 (1851).

Arcangeliella stephensii (Berk.) Z. & D. in Dodge, Ann. Mo. Bot. Gard. 18: 463 (1931).

Basidiomes 1.5-3 cm diam., subglobose, ellipsoid, or irregularly flattened and depressed, often fused together into compound, lobed structures; with a small basal knot of mycelium or sterile base at the point of attachment. **Peridium** orange red (8A-8, 8B-8) when young, darkening to brownish red (8C-8) or reddish brown (8D-8) and mottled with darker areas in age, pale red (7A-3) to greyish red (7B-4) below and in crevices; drying light brown (6D-6); smooth, unpolished at first, subtomentose below, polished above with age; 0.5-0.7 mm thick in section, usually slightly thicker below. **Gleba** labyrinthoid, of unfilled, ellipsoid or irregular chambers, 2-4 per mm at maturity, smaller when young; orange white (6A-2), pale orange (6A-3) to reddish white (7A-2) young, dull red (8B-4) to greyish red (8B-5) in age; drying greyish orange (5B-4). **Tramal plates** 70-300 μ m thick, with a broad hymeno-

phoral trama and poorly developed subhymenial layers, loosely joined, easily movable fresh. **Columella** present, thin, evident only in young gasterocarps, partially percurrent from a small sterile base, dendritic; brownish red (8C-8). **Laticiferous elements** abundant in both hymenophoral trama and lower layer of the peridium, cylindrical to swollen and contorted, 7-11 μm diam., thin-walled, with yellowish refractive contents in KOH; exuding a latex when cut, latex milky white, thin, abundant from around peridium and near columella, also present in tramal plates. **Spores** 9.5-16(18) x 8.4-13.5(15.5) μm (excl. orn.), Q= 1.18; orthotropic to subheterotropic, subglobose to broadly ellipsoid, thin-walled; ornamentation of isolated verrucae and cylindric spines 0.5-1.5 μm high, frequently recurved, spines absent or infrequent near the hilar appendix; subhyaline to dull ochraceous in KOH, weakly amyloid overall in Melzer's reagent, verrucae amyloid, individual spines only partially amyloid, or frequently nonamyloid; hilar appendix 1.5-3 x 0.5-1 μm , tapering or attached to a straight, thin-walled segment of sterigmatal appendage, often with a broad terminal hilar scar. **Basidia** 37-50 x 14-17 μm , inflated-clavate, bearing 2, 3 or 4 spores; sterigmata 6-13 μm long, cylindrical, straight or slightly bowed, often constricted near the hilar appendix. **Cystidia** none. **Hymenophoral** trama regular or nearly so, of interwoven sub-parallel hyphae, 3-5 μm ; laticiferous hyphae abundant. **Subhymenial layer** present but poorly developed, 1-3 cells thick, cells angular, 15-20 x 7-9 μm . **Peridiopellis** a trichodermial palisade of clavate to subulate, often mucronate dermatocystidia, 10-30 x 2-4 μm , arising from a thin epithelium of tightly interwoven, inflated and irregularly shaped cells; hypodermium thick, of loosely interwoven, cylindrical to slightly inflated hyphae, 3-7 μm , laticiferous hyphae abundant, periclinally arranged. (Figs. 1-6).

Material examined

Switzerland: Etzel, hypogeous under *Picea abies*, 12 Aug. 1984, O.K. Miller, Jr. 20180, (VPI).

England: Leigh Woods, Bristol, Aug. 1844, Broome, (K), Holotype; Bristol, Lloyd 7192 (BPI).

Germany: Saar, Rubenheim, hypogeous in calcareous soil under *Carpinus*, *Acer*, *Crataegus*, 27 Aug. 1970, G. Gross 386 (M); Saar, Merzig, hypogeous in calcareous soil, under *Cornus*, *Populus*, *Pinus*, 14 Nov. 1970, G. Gross 4168 (M); Saar, Merzig, hypogeous in calcareous soil under *Pinus*, *Populus*, 4 Nov. 1972, V. Demoulin 4619 (UC).

The hypogeous or erumpant basidiomes grow in small clusters. *Zelleromyces stephensii* undoubtedly forms an ectomycorrhizal symbiosis with particular tree hosts. To the authors' knowledge, no experiments have confirmed its ability to form ectomycorrhizae in pure culture. *Zelleromyces stephensii* has been reported under both conifers and hardwoods (Gross 1980, Froidevaux and Schwärzel 1977), however, it is unusual for a particular species in the genus to be mycorrhizal with both conifer and hardwood tree hosts. More thorough observations are required to determine its ectomycorrhizal hosts and host range. *Zelleromyces stephensii* is distinguished by spores that are weakly amyloid to nearly pseudoamyloid with unconnected verrucae and blunt or recurved, rarely totally amyloid spines, together with the trichodermial peridopellis, filamentous hypodermium, and abundant laticiferous hyphae in the hymenophoral trama and hypodermium. *Zelleromyces stephensii* is apparently endemic to Europe and has been reported from England, Germany, France, Belgium, Czechoslovakia, and Romania (Hawker 1954, Smith 1962, Pegler and Young 1979, Gross 1980, Jülich 1984, Pázmány and Laszlo 1985). Collections examined from North America and labeled as *Z. stephensii* (= *Arcangeliella stephensii*, *Octavianina stephensii*) are misidentified and are *Z. gilkeyae* Singer and Smith, which has larger spores with isolated, uniform, blunt spines and no verrucae. *Zelleromyces ravenelii* (B. & C.) Singer and Smith (= *Arcangeliella ravenelii* (B. & C.) Dodge, *Octaviania ravenelii* Lloyd, *Hydnangium stephensii* var. *ravenelii* Berkeley, *Octaviania stephensii* var. *ravenelii* B. & C.) is a similarly colored species from North America which was once thought to be a variety of *Z. stephensii*, but has heavily amyloid, reticulate ornamentation on the spores.


Literature Cited

- Froidevaux, L. and C. Schwärzel. 1977. Aspects qualitatifs et quantitatifs des champignons hypogés truffoïdes mycorrhiziques en forêt. *Schweizerische Zeitschrift für Forstwesen* 128: 800-813.
- Gross, G., A. Runge and W. Winterhoff. 1980. Bauchpilze (Gasteromycetes s.l.) in der Bundesrepublik und Westberlin. Beihefte zur *Zeitschrift für Mykologie* 2: 1-220.
- Hawker, L. E. 1954. British Hypogeous Fungi. *Philosophical Transactions of the Royal Society of London*. B 237: 429-546.

- Jülich, W. 1984. Die Nichtblätterpilze, Gallertpilze und Bauchpilze. Kleine Kryptogamenflora, vol. 2b, part 1. Gustav Fischer Verlag, Stuttgart, pp.1-626.
- Kornerup, A. and J. H. Wanscher. 1981. Methuen Handbook of Colour 3rd ed. Eyre Methuen, London, pp. 1-252.
- Lanjouw, J., and F. A. Stafleu. 1964. Index Herbariorum part I. The Herbaria of the World. *Regnum Vegetabile* 31: 1-251.
- Smith, A. H. 1962. Notes on Astrogastraceous fungi. *Mycologia* 54: 626-639.
- Pázmány, D. and K. Laszlo. 1985. Seltene Pilze aus Rumänien V. *Notulae Botanicae Horti Agrobotanici* 15: 33-40.
- Pegler, D. N., and T. W. K. Young. 1979. The gasteroid Russulales. *Transactions of the British mycological Society* 72: 353-388.



Figs. 1-2. *Zelleromyces stephensii* basidiomes (OKM 21180). Bars = 1 cm. 1. Habit. 2. Freshly cut to show abundant latex (L).



Figs. 3-6. Microscopic details of *Zelleromyces stephensii* (OKM 21180). Bars = 10 μ m. 3. Spores mounted in Melzer's reagent. 4. Four-spored basidium and portion of subhymenial zone. 5. Two-, three-, and four-spored basidia. 6. Section of peridiopellis showing dermatocystidia and tightly interwoven epithelium.

