

The bug fauna (Heteroptera) of agricultural grasslands in the Schaffhauser Randen (SH) and Rottal (LU), Switzerland, with updated checklists of Heteroptera of the Cantons Luzern and Schaffhausen

Autor(en): **Di Giulio, Manuela / Heckmann, Ralf / Schwab, Andrea**

Objektyp: **Article**

Zeitschrift: **Mitteilungen der Schweizerischen Entomologischen Gesellschaft = Bulletin de la Société Entomologique Suisse = Journal of the Swiss Entomological Society**

Band (Jahr): **73 (2000)**

Heft 3-4

PDF erstellt am: **29.06.2024**

Persistenter Link: <https://doi.org/10.5169/seals-402782>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

The bug fauna (Heteroptera) of agricultural grasslands in the Schaffhauser Randen (SH) and Rottal (LU), Switzerland, with updated checklists of Heteroptera of the Cantons Luzern and Schaffhausen

MANUELA DI GIULIO¹, RALF HECKMANN² & ANDREA SCHWAB¹

We investigated 45 meadows of three different management types in the Schaffhauser Randen (Canton Schaffhausen) and 31 extensively managed meadows in Rottal (Canton Luzern). We recorded 140 species of bugs in the Canton Schaffhausen and 78 in the Canton Luzern. Based on the literature and our own findings we give species-lists for the Cantons of Schaffhausen and Luzern. From the Canton Schaffhausen 207 bug species are known and 201 from the Canton Luzern. The biogeography of some species of special interest is discussed in detail. Three species from the Schaffhauser Randen are new records for Switzerland: *Physatocheila harwoodi* CHINA, 1936, *Polymerus microphthalmus* (WAGNER, 1951) and *Strongylocoris steganoides* (J. SAHLBERG, 1875). The previous records of the latter should be revised.

Keywords: grassland, Heteroptera, Canton Luzern, Canton Schaffhausen, Switzerland

INTRODUCTION

True bugs (Heteroptera) are an ecologically very diverse group, including phytophagous, saprophagous and predatory species (DOLLING, 1991). Furthermore, some species are generalists while others are specialists. The larval stages and adult individuals live in the same habitat and respond sensitively to environmental changes (MORRIS, 1969, 1979; ACHTZIGER, 1995; OTTO, 1996). Some families, especially the Miridae, are very susceptible to chemical sprays used in agricultural production and may therefore be good indicators of ecological change (FAUVEL, 1999). Previous studies have shown that the richness of the bug fauna correlates strongly with total insect diversity (DUELLI & OBRIST, 1998). These properties and the fact that, despite their diversity, they are a manageable group in terms of numbers of species, make the heteropteran bugs an attractive insect group for ecological studies in agricultural systems. Unfortunately, little is known about the distribution and biology of these insects and only few studies are concerned with the complete heteropteran fauna of individual habitats or regions (FAUVEL, 1999). In Switzerland, the number of publications concerning Heteroptera is less than half the corresponding number for Baden-Württemberg in Germany (OTTO, 1994; RIEGER, 1996). According to the literature, 758 species of Heteroptera are known in Switzerland (OTTO, pers. comm., 1997) and 724 in Baden-Württemberg, Germany (RIEGER, 1996).

The aim of this paper is to contribute to the knowledge of the true bugs of Switzerland, with a focus on the regions of Schaffhausen and Rottal (Luzern). The

¹ Swiss Federal Research Station for Agriculture and Agroecology, Reckenholzstrasse 191, CH-8046 Zürich

² Brandenburger Strasse 30, D-78467 Konstanz




Fig. 1. The research areas are located in the North of Switzerland (Canton Schaffhausen) and the Central part of Switzerland (Canton Luzern).

faunistic records presented here come from various studies at the Federal Research Station for Agriculture and Agroecology Zürich-Reckenholz about the influence of management techniques on biodiversity in rural landscapes (financial support by the Swiss National Science Foundation is acknowledged). The ecological results will be presented in separate papers (DI GIULIO, in prep.; SCHWAB, in prep.).

MATERIAL AND METHODS

One research area is located in the Schaffhauser Randen (Canton Schaffhausen, Tab. 1 and Fig. 1) and is part of the Swiss Jura. The soils are nutrient-poor limestone. The average yearly precipitation in Schaffhausen (437 m a.s.l.) is 866 mm, the highest rainfall occurring in summer and the lowest in late winter. The average mean annual temperature is 7.8 °C, with maximum in July (23.2 °C) and minimum in January (-3.9 °C) (SMA, 1998). From 1997 to 1999, we investigated 45 meadows of three grassland management types (medium intensive, moderately intensive and extensive) in four different locations of the Schaffhauser Randen (Tab. 1 and Fig. 1). In all four locations all three grassland types were investigated. The medium intensive meadows are regularly fertilized with slurry and cut 2–3 times per year. The vegetation is dominated by *Arrhenatherum elatius* and *Trisetum flavescens*. The moderately intensive meadows are fertilized occasionally with manure and cut twice a year. No single plant species is dominant but *Trifolium pratense* and *Trisetum flavescens* reach the highest mean cover values. The extensive meadows are not fertilized and are cut once or twice a year. The community is dominated by *Bromus erectus* (STUDER, unpublished data). All sites were sampled during at least one summer by sweep-netting, but frequency and number of sweeps per site varied. Besides, 10 meadows were sampled with a suction sampler of the type «Vortis» (ARNOLD, 1994) and 18 with Barber pitfall traps (CURTIS, 1980).

Tab. 1. The five investigated areas situated in the Canton Schaffhausen and Canton Luzern.

Area	Altitude (m asl)	Municipality	Swiss Coordinates	Geographical Coordinates
Zelgli/Mösli (SH)	800	Hemmental	683'500/ 289'000	8° 34'E / 47° 45' N
Hinterranden (SH)	830	Hemmental Siblingen	683'000/ 287'500	8° 34'E / 47° 44' N
Klosterfeld (SH)	670	Hemmental	687'000/ 288'000	8° 36'E / 47° 44' N
Merishausen (SH)	780-830	Merishausen	683'000/ 291'500- 687'500/ 293'000	8° 36'E / 46° 44' N
Rottal heights (LU)	750-830	Ruswil Buttisholz	650'000/ 220'000	8° 08'E / 47° 07' N

The other study sites are located in the eastern hills of Rottal (Canton Luzern, Tab. 1 and Fig. 1). The area is composed of hilly molasse and a few moraines. The soils tend to be wet and various forms of gleyic brown soils (gleyic camisoles) can be found. The average annual precipitation in Luzern (456 m a.s.l.) is 1154 mm, with highest rainfall in summer and lowest in late winter. The mean annual temperature is 8.3 °C, with the maximum mean monthly temperature in July (17.4 °C) and the minimum in January (-0.7 °C) (SMA, 1998). The whole area is used very intensively for agriculture. During 1999, 31 extensively managed meadows were investigated in an area of approximately 8 km² in the communities of Ruswil and Buttisholz. The vegetation is very heterogeneous, in some sites dominated by grass and in others by forbs. Overall, *Trifolium repens*, *Poa trivialis* and *Ranunculus repens* are the most abundant plants, all of them typical of intensively used meadows which are heavily fertilized and cut early and frequently. Also common are *Lolium multiflorum* and *Dactylis glomerata*, both of them characteristic species of intensive management and wet soils. All sites were sampled monthly from May to August with a standardized sweep net method and with Barber pitfall traps during 10 weeks.

If not specified in the text, the keys of WAGNER (1952, 1966, 1967, 1971, 1973, 1975) and PÉRICART (1972, 1990) were used for the determination of species. R. HECKMANN discussed the faunistic records. A reference collection of the species is located in the entomological collection of the Swiss Federal Institute of Technology (ETHZ).

RESULTS

In total, 167 species have been recorded in both regions, which amounts to 22 % of the 758 (OTTO, pers. comm.) known species of Switzerland. Most species were collected with a sweep net (156) and only 11 species were caught only in pitfall traps or suction samplers (Tab. 2). As there is no Red List of endangered species for Switzerland, those of Baden-Württemberg (RIEGER, 1986), Bayern (ACHTZIGER *et al.*, 1992) and Germany (GÜNTHER *et al.*, 1998) were taken as a basis for assessing the conservation values of the investigated sites (Tab. 2). In the Schaffhauser Randen, we recorded 140 species (12'080 specimens). 29 species (21 %) are listed in the Red Lists for Germany. In Rottal, 78 bug species (1'129 specimens)

Tab. 2. Species list of the investigated meadows in the Schaffhauser Randen (Canton Schaffhausen) and the Rottal heights (Canton Luzern).

Columns: M: Merishausen, H: Hinterranden, K: Klosterfeld, Z: Zelgli. SH and LU.: Total number of individuals recorded in the two Cantons. Met.: Sampling method used; n: Sweep net, s: Suction sampler, p: Pit fall traps. R.L.: 1: Baden-Württemberg (D), 2: Bayern (D), 3: Germany. “*”: Species specialized on plants not normally occurring in meadows (trees, shrubs, ferns) and which probably have been drifted by the wind. The nomenclature of the species list from Ceratocombidae to Miridae (excluding the arrangement in the subfamily Mirinae) follows the Catalogue of the Heteroptera of the Palaearctic Region (AUKEMA & RIEGER, 1995, 1996, 1999). The subfamily Mirinae (changed names adapted from the above catalogue) of Miridae and the Pentatomorpha follow the Verzeichnis der Wanzen Mitteleuropas (GÜNTHER & SCHUSTER, 1990).

SPECIES	M	H	K	Z	SH	LU	Met.	R.L.
CERATOCOMBIDAE								
<i>Ceratocombus coleopratus</i> (Zetterstedt, 1819)	+				6		s	2
SALDIDAE								
<i>Saldula c-album</i> (Fieber, 1859)						3	p	1
<i>Saldula orthochila</i> (Fieber, 1859)	+				1		n	
<i>Saldula saltatoria</i> (Linnaeus, 1758)	+	+			2	11	n/p	
TINGIDAE								
<i>Acalypta carinata</i> (Panzer, 1806)	+				4	1	p	1,2
<i>Acalypta marginata</i> (Wolff, 1804)	+		+		90	3	n/p	2
<i>Campylosteira verna</i> (Fallén, 1826)	+	+			18		n/s/p	2,3
<i>Catoplatys fabricii</i> (Stål, 1868)	+	+	+		71		n/p	2,3
<i>Kalama tricornis</i> (Schrank, 1801)	+	+	+	+	22	9	n/s/p	2
* <i>Physatocheila harwoodi</i> China, 1936				+	1		n	2,3
<i>Tingis reticulata</i> Herrich-Schäffer, 1835						5	n/p	
NABIDAE								
<i>Himacerus major</i> (A. Costa, 1842)						1	p	2
<i>Himacerus mirmicoides</i> (O. Costa, 1834)	+		+		7	20	n/p	
* <i>Himacerus apterus</i> (Fabricius, 1798)						3	n	
<i>Nabis flavomarginatus</i> (Scholtz, 1847)				+	3		n	
<i>Nabis brevis brevis</i> Scholtz, 1847	+		+		34	4	n/p	
<i>Nabis ferus</i> (Linnaeus, 1758)	+	+	+	+	5	18	n/p	
<i>Nabis pseudoferus pseudoferus</i> Remane, 1949	+	+	+	+	131	2	n/s/p	
<i>Nabis punctatus punctatus</i> A. Costa, 1847	+	+	+		18		n	1,2
<i>Nabis rugosus</i> (Linnaeus, 1758)	+		+	+	10	11	n/s/p	
ANTHOCORIDAE								
* <i>Acomporis alpinus</i> Reuter, 1875	+				1	1	n	2
<i>Anthocoris nemorum</i> (Linnaeus, 1761)	+			+	2	40	n	
* <i>Temnostethus pusillus</i> (Herrich-Schäffer, 1835)	+				1	1	p	
<i>Orius majusculus</i> (Reuter, 1879)			+		2	9	n	
<i>Orius minutus</i> (Linnaeus, 1758)	+			+	3	6	n	

THE HETEROPTERAN FAUNA OF SCHAFFHAUSEN AND LUZERN

SPECIES	M	H	K	Z	SH	LU	Met.	R.L.
MIRIDAE								
* <i>Bryocoris pteridis</i> (Fallén, 1807)						1	n	
<i>Campyloneura virgula</i> (Herrich-Schäffer, 1835)			+		1	1	n	
<i>Dicyphus annulatus</i> (Wolff, 1804)			+		10		n	
<i>Dicyphus globulifer</i> (Fallén, 1829)	+				2		n	
<i>Dicyphus errans</i> (Wolff, 1804)						1	n	
* <i>Dicyphus pallidus</i> (Herrich-Schäffer, 1836)						6	n/p	
* <i>Dicyphus stachydis stachydis</i> J. Sahlberg, 1878						1	n	
<i>Deraeocoris morio</i> (Boheman, 1852)	+		+		9		n	1,2,3
<i>Deraeocoris ruber</i> (Linnaeus, 1758)	+			+	2	10	n	
<i>Pithanus maerkelii</i> (Herrich-Schäffer, 1838)						15	n/p	
<i>Leptopterna dolobrata</i> (Linnaeus, 1758)	+	+	+	+	4152		n/p	
<i>Stenodema calcarata</i> (Fallén, 1807)						40	n	
<i>Stenodema holsata</i> (Fabricius, 1787)	+			+	7	78	n/p	
<i>Stenodema laevigata</i> (Linnaeus, 1758)	+	+	+	+	164	8	n/pi	
<i>Stenodema virens</i> (Linnaeus, 1767)						3	n	
<i>Megaloceraea recticornis</i> (Geoffroy, 1785)	+	+		+	222		n/p	
<i>Notostira elongata</i> (Geoffroy, 1785)	+				1	1	n	
<i>Notostira erratica</i> (Linnaeus, 1758)	+	+	+	+	1065	177	n/s/p	
<i>Trigonotylus caelestialium</i> (Kirkaldy, 1902)	+	+	+	+	157	1	n/s	
<i>Phytocoris longipennis</i> Flor, 1860	+				1		n	
<i>Adelphocoris lineolatus</i> (Goeze, 1778)	+	+	+		473	4	n	
<i>Adelphocoris seticornis</i> (Fabricius, 1775)	+	+	+	+	696		n/p	
<i>Brachycoleus pilicornis pilicornis</i> (Panzer, 1805)				+	1		n	2,3
<i>Calocoris affinis</i> (Herrich-Schäffer, 1835)	+				2	3	n	
* <i>Calocoris alpestris</i> (Meyer-Dür, 1843)						8	n	2
<i>Closterotmus biclavatus</i> (Herrich-Schäffer, 1835)	+				1		n	
<i>Closterotomus norvegicus</i> (Gmelin, 1790)	+		+		12	322	n/p	
* <i>Grypocoris sexguttatus</i> (Fabricius, 1777)	+				1		n	2
* <i>Rhodomiris striatellus striatellus</i> (Fabricius, 1794)						1	n	
<i>Hadrodemus m-flavum</i> (Goeze, 1778)	+	+	+	+	239		n	
<i>Stenotus binotatus</i> (Fabricius, 1794)	+				1	1	n	
* <i>Dichrooscytus intermedius</i> Reuter, 1885	+			+	2		n	
<i>Apolygus lucorum</i> (Meyer-Dür, 1843)			+		1		n	
<i>Apolygus spinolae</i> (Meyer-Dür, 1841)			+		2		n	
<i>Lygocoris pabulinus</i> (Linnaeus, 1761)	+				4	35	n	
* <i>Lygocoris viridis</i> (Fallén, 1807)						3	n	
<i>Lygus pratensis</i> (Linnaeus, 1758)	+	+	+	+	677	1	n/s	
<i>Lygus rugulipennis</i> Poppius, 1911	+	+	+	+	371	161	n/s/p	
<i>Lygus wagneri</i> Remane, 1955	+			+	4		n	
<i>Orthops basalıs</i> (A. Costa, 1853)	+	+	+		6	4	n	
<i>Orthops campestris</i> (Linnaeus, 1758)	+				1	2	n	
<i>Orthops kalmii</i> (Linnaeus, 1758)	+	+			10		n	
* <i>Pinalitus atomarius</i> (Meyer-Dür, 1843)						1	n	1,3
* <i>Pinalitus visicola</i> (Puton, 1888)						1	n	1,2
<i>Charagochilus gyllenhalii</i> (Fallén 1807)	+		+	+	19	1	n/s/p	
<i>Polymerus nigrita</i> (Fallén, 1829)	+		+	+	13		n	

SPECIES	M	H	K	Z	SH	LU	Met.	R.L.
<i>Polymerus microphthalmus</i> (Wagner, 1951)	+				1		n	
<i>Polymerus unifasciatus</i> (Fabricius, 1794)	+	+	+	+	745	2	n/s/p	
<i>Capsodes gothicus gothicus</i> (Linnaeus, 1758)				+	2		n	
<i>Capsus ater</i> (Linnaeus, 1758)	+	+	+	+	11		n	
<i>Halticus apterus apterus</i> (Linnaeus, 1758)	+		+	+	44	1	n/s	
<i>Orthocephalus coriaceus</i> (Fabricius, 1777)	+				2	3	n	
<i>Orthocephalus saltator</i> (Hahn, 1835)			+		3		n	
<i>Strongylocoris steganoides</i> (J. Sahlberg, 1875)	+		+	+	39		n/p	
* <i>Blepharidopterus angulatus</i> (Fallén, 1807)						1	n	
<i>Globiceps flavomaculatus</i> (Fabricius, 1794)	+				2	2	n	
<i>Globiceps fulvicollis</i> Jakovlev, 1877	+		+		23		n/p	
* <i>Heterocordylus tumidicornis</i> (Herrich-Schäffer, 1835)			+		1		n	
<i>Heterotoma planicornis</i> (Pallas, 1772)						1	n	
* <i>Malacocoris chlorizans</i> (Panzer, 1794)	+				1		n	
* <i>Cremnocephalus alpestris</i> Wagner, 1941						1	n	
<i>Hallodapus rufescens</i> (Burmeister, 1835)	+				2		n	1,2,3
<i>Ambylytus nasutus</i> (Kirschbaum, 1856)	+				2	2	n	
* <i>Atractotomus magnicornis</i> (Fallén, 1807)	+			+	1	8	n	
* <i>Atractotomus parvulus</i> Reuter, 1878	+				3		n	
<i>Campylomma verbasci</i> (Meyer-Dür, 1843)	+				1		n	
<i>Chlamydatus pulicarius</i> (Fallén, 1807)	+	+	+	+	215	1	n/s/p	
<i>Chlamydatus pullus</i> (Reuter, 1870)		+	+		2		n	
<i>Criocoris crassicornis</i> (Hahn, 1834)				+	1		n	
* <i>Harpocera thoracica</i> (Fallén, 1807)						2	n	
<i>Macrotylus herrichi</i> (Reuter, 1873)	+	+		+	11		n	2
<i>Megalocoleus molliculus</i> (Fallén, 1807)	+			+	2	1	n	
* <i>Phoenicocoris obscurellus</i> (Fallén, 1829)	+				1		n	
<i>Plagiognathus arbustorum arbustorum</i> (Fabricius, 1794)	+		+		4	10	n/p	
<i>Plagiognathus chrysanthemi</i> (Wolff, 1804)	+	+	+	+	1274		n/p	
* <i>Plesiodema pinetella</i> (Zetterstedt, 1828)	+				1		n	
* <i>Psallus perrisi</i> (Mulsant & Rey, 1852)						3	n	
* <i>Psallus piceae</i> Reuter, 1878						2	n	
<i>Psallus haematodes</i> (Gmelin, 1790)						3	n	
* <i>Psallus varians varians</i> (Herrich-Schäffer, 1841)						4	n	
ARADIDAE								
* <i>Aradus cinnamomeus</i> (Panzer, 1794)	+				4	2	p	
PIESMATIDAE								
<i>Piesma maculatum</i> (Laporte, 1832)	+			+	2		n	
BERYTIDAE								
<i>Berytinus clavipes</i> (Fabricius, 1775)	+		+		10		n/p	
<i>Berytinus minor</i> (Herrich-Schäffer, 1835)	+	+	+	+	43	9	n/s/p	
<i>Berytinus crassipes</i> (Herrich-Schäffer, 1835)				+	1		n	2
<i>Berytinus montivagus</i> (Meyer-Dür, 1841)	+	+	+		16		n/p	1,2
<i>Berytinus signoreti</i> (Fieber, 1859)			+		1		n	2
<i>Gampsocoris punctipes</i> (Germar, 1822)			+		1		n	

THE HETEROPTERAN FAUNA OF SCHAFFHAUSEN AND LUZERN

SPECIES	M	H	K	Z	SH	LU	Met.	R.L.
LYGAEIDAE								
<i>Nysius ericae</i> (Schilling, 1829)			+		1	2	n	
* <i>Kleidocerys resedae</i> (Panzer, 1797)						1	n	
<i>Cymus melanocephalus</i> Fieber, 1861						1	n	
<i>Ischnodemus sabuleti</i> (Fallén, 1829)		+			2		n	2
<i>Platylax salviae</i> (Schilling, 1829)	+		+		19		n	
<i>Plinthisus brevipennis</i> (Latreille, 1807)	+		+		14		n/p	
<i>Tropistethus holosericeus</i> (Scholtz, 1845)	+		+		3		n	
<i>Drymus latus</i> Douglas & Scott, 1871	+				2		p	1,2,3
<i>Drymus ryeii</i> Douglas & Scott, 1865	+				2		n/s	
<i>Drymus sylvaticus</i> (Fabricius, 1775)	+				8		n/p	
* <i>Gastrodes abietum</i> Bergroth, 1914	+				1	2	n	
<i>Scolopostethus affinis</i> (Schilling, 1829)	+				1		p	
<i>Scolopostethus thomsoni</i> Reuter, 1874	+				1	1	n	
<i>Stygnocoris rusticus</i> (Fallén, 1807)	+				2		s/p	
<i>Stygnocoris sabulosus</i> (Schilling, 1829)	+				2		s	
<i>Graptopeltus lynceus</i> (Fabricius, 1775)	+	+			53		n/p	
<i>Rhyparochromus pini</i> (Linnaeus, 1758)	+				46		p	
<i>Peritrechus gracilicornis</i> Puton, 1877	+	+	+	+	57		n/p	1,2
<i>Megalonotus chiragra chiragra</i> (Fabricius, 1794)	+	+			97	2	n/s/p	
<i>Emblethis verbasci</i> (Fabricius, 1803)			+		1		n	1
PYRRHOCORIDAE								
<i>Pyrrhocoris apterus</i> (Linnaeus, 1758)		+			1		n	
COREIDAE								
<i>Syromastus rhombeus</i> (Linnaeus, 1767)	+				2		n	1
<i>Coreus marginatus</i> (Linnaeus, 1758)			+		1		n	
<i>Coriomeris denticulatus</i> (Scopoli, 1763)	+		+	+	11		n	
ALYDIDAE								
<i>Alydus calcaratus</i> (Linnaeus, 1758)	+		+		6		n	
RHOPALIDAE								
<i>Corizus hyoscyami</i> (Linnaeus, 1758)	+		+		3		n	
<i>Rhopalus parumpunctatus</i> Schilling, 1829	+		+		2		n	
<i>Rhopalus subrufus</i> (Gmelin, 1790)	+				3		n	
<i>Stictopleurus abutilon abutilon</i> (Rossi, 1790)	+				3		n	
<i>Stictopleurus punctatonervosus</i> (Goeze, 1778)	+				1		n	
PLATASPIDAE								
<i>Coptosoma scutellatum</i> (Geoffroy, 1785)			+		16		n	
CYDNIDAE								
<i>Sehirus luctuosus</i> Mulsant & Rey, 1866	+	+			17		n/p	
<i>Canthophorus impressus</i> Horvath, 1881	+			+	2		n	1
<i>Adomerus biguttatus</i> (Linnaeus, 1758)	+	+	+	+	90		n/p	1
<i>Legnotus picipes</i> (Fallén, 1807)		+	+		3		n	2
THYREOCORIDAE								
<i>Thyreocoris scarabaeoides</i> (Linnaeus, 1758)	+				1		n	

SPECIES	M	H	K	Z	SH	LU	Met.	R.L.
SCUTELLERIDAE								
<i>Eurygaster maura</i> (Linnaeus, 1758)		+	+		22	1	n	
<i>Eurygaster testudinaria</i> (Geoffroy, 1758)			+		2		n	
PENTATOMIDAE								
<i>Sciocoris macrocephalus</i> Fieber, 1851	+				3		n	1,3
<i>Sciocoris microphthalmus</i> Flor, 1860	+				50		n	2
<i>Aelia acuminata</i> (Linnaeus, 1758)			+		3		n	
<i>Neottiglossa pusilla</i> (Gmelin, 1789)	+				2		n	
<i>Rubiconia intermedium</i> (Wolff, 1811)	+				1		n	2,3
<i>Palomena prasina</i> (Linnaeus, 1761)	+		+		2	6	n	
<i>Palomena viridissima</i> (Poda, 1761)			+		3	4	n	
<i>Dryocoris vernalis</i> (Wolff, 1804)			+		1		n	
<i>Carpocoris fuscispinus</i> (Boheman, 1849)	+	+	+	+	97	8	n/s/p	
<i>Carpocoris purpureipennis</i> (De Geer, 1773)	+	+	+	+	23	3	n	
<i>Dolycoris baccarum</i> (Linnaeus, 1758)	+	+	+	+	135		n/s/p	
<i>Eurydema oleraceum</i> (Linnaeus, 1758)	+	+	+		8		n/p	
* <i>Pentatoma rufipes</i> (Linnaeus, 1758)	+				2		n	
<i>Troilus luridus</i> (Fabricius, 1775)	+				1		n/p	
* <i>Arma custos</i> (Fabricius, 1794)						1	n	
<i>Zicrona caerulea</i> (Linnaeus, 1758)	+	+	+	+	18		n/p	2
Total number of species	167	112	41	70	46	140	78	

were recorded. Ten species account for more than 80 % of all specimens recorded and only 10 species (13 %) are listed in at least one of the Red Lists for Germany.

Altogether 31 species have been found that do not belong to grassy habitats (see "*" in Tab. 2). Fourteen of them were found in Schaffhausen and 22 in Luzern. Most of these species were probably drifted by wind from trees and shrubs in the nearby woodlands and the forest edge. That species characteristic of the herbal layer of the forest edge can also live in moist meadows is confirmed by a much higher rate of vagrants in the Luzern than in the Schaffhauser meadows (JUDEX, 1999).

DISCUSSION

General discussion

Recent faunistic publications concerning the heteropteran fauna of Switzerland mostly deal with the entomologically more interesting southern part of the country, especially the Canton Ticino where till now 395 species have been found (PÉRICART, 1984; GÖLLNER-SCHIEDING & REZBANYAI-RESER, 1992; OTTO, 1992, 1994, 1995, 1996; REZBANYAI-RESER, 1993, 1997; OTTO & BÜRKI, 1996; OTTO & REZBANYAI-RESER, 1996; RAMPAZZI & DETHIER, 1997). The other Cantons are not well investigated and the fauna is much less well known than that of various regions of Baden-Württemberg. Tab. 3 gives a survey of the faunistic knowledge in the various Cantons of Switzerland in relation to data available from Central Europe (GÜNTHER & SCHUSTER, 1990), Switzerland (OTTO, unpublished list), Baden-Württemberg (RIEGER, 1996) and the Landkreis Konstanz in Baden-Württemberg (HECKMANN, 1989, 1990, 1993, 1996, 1999; HECKMANN & RIEGER, in prep.).

Tab. 3. List of absolute and relative species numbers of ecologically most important families caught in different regions.

Columns: EU: Central Europe (GÜNTHER & SCHUSTER, 1990). CH: Switzerland (OTTO, unpublished list and pers. comm., 1997: uncontrolled literature data). BW: Baden-Württemberg (RIEGER, 1996: literature data). KN: Landkreis Konstanz (HECKMANN, 1989, 1990, 1993, 1996, 1999; HECKMANN & RIEGER, in prep., unpublished verified records). TI: Canton Ticino (PÉRICART, 1984; REZBANYAI-RESER, 1993; OTTO, 1994, 1995, 1996; OTTO & BÜRKI, 1996; REZBANYAI-RESER, 1997: the 12 unpublished records of DETHIER excluded). SH: Canton Schaffhausen (Tab. 5). LU: Canton Luzern (Tab. 4). TG: Canton Thurgau (BLÖCHLINGER, verified records of unpublished list & pers. comm., 1999 partly based on HOFMÄNNER, 1928). Other families: Hydrocorisae, Amphibiocorisae and the "rest" of Geocorisae.

	ABSOLUTE NUMBERS								PERCENTAGE NUMBERS							
	EU	CH	BW	KN	TI	SH	LU	TG	EU	CH	BW	KN	TI	SH	LU	TG
Tingidae	74	49	45	22	20	10	4	10	7,0	6,5	6,2	5,4	5,2	4,8	2,3	3,8
Nabidae	18	16	14	11	7	8	8	7	1,7	2,1	1,9	2,7	1,8	3,9	4,5	2,7
Anthocoridae	52	36	33	23	18	7	9	12	4,9	4,7	4,6	5,6	4,7	3,4	5,1	4,6
Reduviidae	18	14	13	7	7	5	1	6	1,7	1,8	1,8	1,7	1,8	2,4	0,6	2,3
Miridae	401	271	284	164	156	76	100	95	37,8	35,8	39,0	39,9	40,8	36,7	56,5	36,1
Berytidae	14	10	10	7	2	7	1	4	1,3	1,3	1,4	1,7	0,5	3,4	0,6	1,5
Lygaeidae	155	116	110	62	58	25	19	30	14,6	15,3	15,2	15,1	15,2	12,1	10,7	11,4
Coreidae	27	23	17	7	10	5	1	5	2,5	3,0	2,3	1,7	2,6	2,4	0,6	1,9
Rhopalidae	17	15	14	10	10	5	2	8	1,6	2,0	1,9	2,4	2,6	2,4	1,1	3,0
Pentatomidae	70	52	43	26	28	18	12	23	6,6	6,9	5,9	6,3	7,3	8,7	6,8	8,7
Other families	251	157	141	72	67	31	44	63	23,7	20,1	19,5	17,5	17,5	15,0	21,9	24,0
Species number	1061	758	724	411	383	207	201	263	100	71,4	68,2	38,7	36,0	19,5	18,9	24,8

One way to assess the ecological differences between regions of a country is to look at differences in species numbers of families known to be indicators for various climate types. The numbers of species in families like the Tingidae, Reduviidae, Lygaeidae, Rhopalidae, Coreidae, Berytidae and the Pentatomidae increase from boreal and alpine areas to mediterranean and tropical habitats. These taxa can therefore be used as indicator families of xero-thermophily. On the other hand, species numbers of Miridae and Anthocoridae decrease in the same direction, being indicators for more temperate climates (GÖLLNER-SCHEIDING, 1989a). Although the data are not complete, it is interesting to note that there is a good fit of the data from SH, KN and even TG which are located close together (Tab. 3). Moreover, there are big differences between these areas and the Canton Luzern which has the highest proportions of Miridae and the lowest proportions of xero-thermophilic Tingidae, Reduviidae, Berytidae, Lygaeidae, Coreidae and Rhopalidae.

We found more than one fifth of all known species of Switzerland in the meadows of the Schaffhauser Randen and the Rottal. The sites in the region of Schaffhausen, however, are much more diverse than those in Luzern, and we found nearly twice as many species and a much higher proportion of rare or endangered species. Most of the latter are xero-thermophilic species, which are restricted to dry and extensively managed meadows (DI GIULIO, in prep.). The climatic conditions in the Schaffhauser Randen not only favour these species directly, but in combination with the soil conditions, have limited the intensification of agricultural production. It is known for other taxa (e.g. butterflies and flowering plants) that many species can persist here which have disappeared from the other regions of Switzerland (ISLER-HÜBSCHER, 1980; SCHIESS-BÜHLER & SCHIESS, 1997).

Canton Luzern

Faunistic data are available from the Canton Luzern for the upland moor Balmoos near Hasle (REZBANYAI, 1980; GÖLLNER-SCHIEDING, 1981) and for the surroundings of the Vogelwarte Sempach (GÖLLNER-SCHIEDING, 1982). Later the Vogelmoos near Neudorf (GÖLLNER-SCHIEDING, 1990) was investigated and occasional records for different places in the Canton of Luzern were published (GÖLLNER-SCHIEDING, 1989b). The record of *Panaorus adspersus* MULS. & REY was added

Tab. 4. Check list of Heteroptera for the Canton Luzern. Nomenclature as in Tab. 2. The number of species in the different families are noted. The asterisks indicate the 26 new species for the Canton.

CHECK LIST OF HETEROPTERA FOR THE CANTON LUZERN		201	
NEPIDAE	2	GERRIDAE	6
<i>Nepa cinerea</i> L.		<i>Aquarius paludum</i> F.	
<i>Ranatra linearis</i> L.		<i>Gerris argentatus</i> Schum.	
CORIXIDAE	12	<i>Gerris gibbifer</i> Schum.	
<i>Micronecta scholtzi</i> Fieb.		<i>Gerris lacustris</i> L.	
<i>Callicorixa praeusta</i> Fieb.		<i>Gerris odontogaster</i> Zett.	
<i>Corixa punctata</i> Ill.		<i>Gerris thoracicus</i> Schum.	
<i>Hesperocorixa linnaei</i> Fieb.		SALDIDAE	4
<i>Hesperocorixa sahlbergi</i> Fieb.		<i>Chartoscirta cincta</i> H.-S.	
<i>Paracorixa concinna</i> Fieb.		<i>Macrosaldula variabilis</i> H.-S.	
<i>Sigara nigrolineata</i> Fieb.		* <i>Saldula c-album</i> Fieb.	
<i>Sigara semistriata</i> Fieb.		<i>Saldula saltatoria</i> L.	
<i>Sigara striata</i> L.		TINGIDAE	4
<i>Sigara distincta</i> Fieb.		* <i>Acalypta carinata</i> Pz.	
<i>Sigara falleni</i> Fieb.		* <i>Acalypta marginata</i> Wff.	
<i>Sigara lateralis</i> Lch.		* <i>Kalama tricornis</i> Schrank	
NAUCORIDAE	1	* <i>Tingis reticulata</i> H.-S.	
<i>Ilyocoris cimicoides</i> L.		NABIDAE	8
NOTONECTIDAE	1	* <i>Himacerus major</i> A. C.	
<i>Notonecta glauca</i> L.		<i>Himacerus mirmicoides</i> O. C.	
<i>Notonecta maculata</i> F.		* <i>Himacerus apterus</i> F.	
<i>Notonecta viridis</i> Delc.		<i>Nabis limbatus</i> Dhlbm.	
PLEIDAE	1	* <i>Nabis brevis</i> Sz.	
<i>Plea minutissima</i> Lch.		<i>Nabis ferus</i> L.	
MESOVELIIDAE	1	<i>Nabis pseudoferus</i> Rem.	
<i>Mesovelia furcata</i> Muls. & Rey		<i>Nabis rugosus</i> L.	
HYDROMETRIDAE	2	ANTHOCORIDAE	9
<i>Hydrometra gracilentata</i> Horv.		<i>Acompocoris alpinus</i> Reut.	
<i>Hydrometra stagnorum</i> L.		<i>Anthocoris confusus</i> Reut.	
VELIIDAE	4	<i>Anthocoris limbatus</i> Fieb.	
<i>Microvelia pygmaea</i> Duf.		<i>Anthocoris nemorum</i> L.	
<i>Microvelia reticulata</i> Burm.		<i>Temnostethus pusillus</i> H.-S.	
<i>Velia caprai</i> Tam.		<i>Orius laticollis</i> Reut.	
<i>Velia saulii</i> Tam.		<i>Orius majusculus</i> Reut.	
		<i>Orius minutus</i> L.	
		<i>Orius vicinus</i> Rib.	

CIMICIDAE	1	
<i>Cimex lectularius</i> L.		
REDUVIIDAE	1	
<i>Reduvius personatus</i> L.		
MIRIDAE	100	
* <i>Bryocoris pteridis</i> Fall.		
<i>Monalocoris filicis</i> L.		
* <i>Campyloneura virgula</i> H.-S.		
<i>Dicyphus epilobii</i> Reut.		
<i>Dicyphus errans</i> Wff.		
<i>Dicyphus hyalinipennis</i> Burm.		
* <i>Dicyphus pallidus</i> H.-S.		
<i>Dicyphus stachydis</i> J. Shlbg.		
<i>Alloeotomus germanicus</i> E. Wgn.		
<i>Alloeotomus gothicus</i> Fall.		
<i>Deraeocoris ruber</i> L.		
<i>Deraeocoris lutescens</i> Schill.		
<i>Pithanus maerkelii</i> H.-S.		
<i>Leptopterna dolobrata</i> L.		
<i>Teratocoris paludum</i> J. Shlbg.		
<i>Stenodema calcarata</i> Fall.		
<i>Stenodema holsata</i> F.		
<i>Stenodema laevigata</i> L.		
<i>Stenodema virens</i> L.		
<i>Megaloceraea recticornis</i> Geoffr.		
<i>Notostira elongata</i> Geoffr.		
<i>Notostira erratica</i> L.		
<i>Trigonotylus caelestialium</i> Kirk.		
<i>Phytocoris dimidiatus</i> Kb.		
<i>Phytocoris intricatus</i> Flor		
<i>Phytocoris longipennis</i> Flor		
<i>Phytocoris tiliae tiliae</i> F.		
<i>Phytocoris pini</i> Kb.		
<i>Pantilius tunicatus</i> F.		
* <i>Adelphocoris lineolatus</i> Gz.		
<i>Adelphocoris seticornis</i> F.		
<i>Calocoris affinis</i> H.-S.		
<i>Calocoris alpestris</i> Mey.-D.		
<i>Calocoris roseomaculatus</i> De G.		
<i>Closterotmus biclavatus</i> H.-S.		
<i>Closterotomus norvegicus</i> Gm.		
<i>Grypocoris sexguttatus</i> F.		
<i>Rhabdomiris striatellus</i> F.		
<i>Stenotus binotatus</i> F.		
<i>Apolygus lucorum</i> Mey.-D.		
<i>Lygocoris pabulinus</i> L.		
<i>Lygocoris rugicollis</i> Fall.		
<i>Lygocoris contaminatus</i> Fall.		
<i>Lygocoris viridis</i> Fall.		
		<i>Lygus punctatus</i> Zett.
		<i>Lygus pratensis</i> L.
		<i>Lygus rugulipennis</i> Popp.
		<i>Lygus wagneri</i> Rem.
		<i>Orthops basalis</i> A. C.
		* <i>Orthops campestris</i> L.
		<i>Orthops montanus</i> Schill.
		* <i>Pinalitus atomarius</i> Mey.-D.
		<i>Pinalitus cervinus</i> H.-S.
		<i>Pinalitus rubricatus</i> Fall.
		* <i>Pinalitus visicola</i> Put.
		<i>Agnocoris rubicundus</i> Fall.
		<i>Liocoris tripustulatus</i> F.
		<i>Camptozygum aequale</i> Vill.
		<i>Charagochilus gyllenhalii</i> Fall.
		<i>Polymerus holosericeus</i> Hahn
		* <i>Polymerus unifasciatus</i> F.
		<i>Capsus ater</i> L.
		* <i>Halticus apterus</i> L.
		* <i>Orthocephalus coriaceus</i> F.
		<i>Blepharidopterus angulatus</i> Fall.
		<i>Blepharidopterus diaphanus</i> Kb.
		<i>Dryophilocoris flavoquadrimaculatus</i> De G.
		* <i>Globiceps flavomaculatus</i> F.
		* <i>Heterotoma planicornis</i> Pall.
		<i>Mecomma ambulans</i> Fall.
		<i>Orthotylus marginalis</i> Reut.
		<i>Orthotylus nassatus</i> F.
		<i>Orthotylus prasinus</i> Fall.
		<i>Pseudoloxops coccineus</i> Mey.-D.
		<i>Cremnocephalus alpestris</i> E. Wgn.
		<i>Pilophorus clavatus</i> L.
		<i>Pilophorus confusus</i> Kb.
		<i>Pilophorus perplexus</i> Doug. & Sc.
		* <i>Ambylytus nasutus</i> Kb.
		<i>Atractotomus kolenatii</i> Flor
		<i>Atractotomus magnicornis</i> Fall.
		<i>Compsidolon salicellum</i> H.-S.
		* <i>Chlamydatus pulicarius</i> Fall.
		<i>Harpocera thoracica</i> Fall.
		* <i>Megalocoleus molliculus</i> Fall.
		<i>Orthonotus rufifrons</i> Fall.
		<i>Phoenicocoris obscurellus</i> Fall.
		<i>Phylus coryli</i> L.
		<i>Phylus melanocephalus</i> L.
		<i>Placochilus seladonicus</i> Fall.
		<i>Plagiognathus arbustorum</i> F.
		<i>Psallus betuleti</i> Fall.
		<i>Psallus ambiguus</i> Fall.
		* <i>Psallus perrisi</i> Muls. & Rey

MIRIDAE	CONT.	
* <i>Psallus piceae</i> Reut. <i>Psallus falleni</i> Reut. <i>Psallus haematodes</i> Gm. <i>Psallus lepidus</i> Fieb. <i>Psallus mollis</i> Muls. & Rey <i>Psallus varians varians</i> H.-S.		
ARADIDAE		1
* <i>Aradus cinnamomeus</i> Pz.		
BERYTIDAE		1
<i>Berytinus minor</i> H.-S.		
LYGAEIDAE		19
<i>Spilostethus saxatilis</i> Scop. <i>Nysius cymoides</i> Spin. * <i>Nysius ericae</i> Schill. <i>Kleidocerys resedae</i> Pz. <i>Cymus glandicolor</i> Hahn <i>Cymus aurescens</i> Dist. <i>Cymus melanocephalus</i> Fieb. <i>Ischnodemus sabuleti</i> Fall. <i>Gastrodes abietum</i> Bgrth. <i>Gastrodes grossipes</i> De G. <i>Scolopostethus affinis</i> SCHILL. <i>Scolopostethus thomsoni</i> REUT. <i>Stygnocoris rusticus</i> FALL. <i>Pachybrachius fracticollis</i> SCHILL. <i>Panaorus adpersus</i> MULS. & REY <i>Rhyparochromus pini</i> L. <i>Peritrechus geniculatus</i> HAHN <i>Peritrechus nubilus</i> FALL. <i>Megalonotus chiragra</i> F.		
COREIDAE		1
<i>Coreus marginatus</i> L.		
RHOPALIDAE		2
<i>Rhopalus subrufus</i> Gm. <i>Stictopleurus crassicornis</i> L.		
SCUTELLERIDAE		2
<i>Eurygaster maura</i> L. <i>Eurygaster testudinaria</i> Geoffr.		
PENTATOMIDAE		12
<i>Eysarcoris fabricii</i> Kirk. <i>Palomena prasina</i> L. <i>Palomena viridissima</i> Pd. <i>Carpocoris fuscispinus</i> Boh. <i>Carpocoris purpureipennis</i> De G. <i>Dolycoris baccarum</i> L. <i>Eurydema dominulum</i> Scop. <i>Rhaphigaster nebulosa</i> Pd. <i>Pentatoma rufipes</i> L. <i>Picromerus bidens</i> L. <i>Arma custos</i> F. <i>Zicrona caerulea</i> L.		
ACANTHOSOMATIDAE		4
<i>Acanthosoma haemorrhoidale</i> L. <i>Elasmotethus minor</i> Horv. <i>Elasmotethus interstinctus</i> L. <i>Elasmucha grisea</i> L.		

from the Museum of Natural History in Paris (PÉRICART, 1998) and 35 species by an investigation near Rottal (JUDEX, 1999). Thirty species of water and semiaquatic bugs (Nepomorpha and Gerromorpha) were found in the Wauwiler Ebene and 18 in the Möslweiher near Schötz (WIPRÄCHTIGER, 1999a, 1999b). We found 78 species for Rottal, 26 of them being new records for this Canton (see "*" in Tab. 4). Altogether 201 species have been recorded in the Canton Luzern.

In the publications of GÖLLNER-SCHIEDING (1981, 1982) *Orthops basalis* A.C. was published as *Orthops kalmii* L. The misidentification of these two species was apparently due to a mistake in the key of WAGNER (1967, 1971) which later was revised (RIEGER, 1985).

Canton Schaffhausen

The earliest records for the Canton Schaffhausen include 23 species of Miridae collected by S. SEILER (MEYER-DÜR, 1843). A list of 76 species was published by FREY-GESSNER (1864–1866), most of them also collected by S. SEILER in the mid 19th century. In his list he included a revised list of the Mirids recorded by MEYER-

DÜR (1843). Much later the occurrence of two Saldid species was reported (DETHIER & PÉRICART, 1990). Together with our results, we present here a list of 207 species of Heteroptera for the Canton Schaffhausen, 118 of them being new records for the Canton (see “*” in Tab. 5).

We were not able to check the determinations of FREY-GESSNER because the collection was destroyed by bombing of Schaffhausen during World War II (A. MÜL-

Tab. 5. Check list of Heteroptera for the Canton Schaffhausen. Nomenclature as in Tab. 2. “?”: doubtful historical records. The asterisks indicate the 118 new species for the Canton.

CHECK LIST OF HETEROPTERA FOR THE CANTON SCHAFFHAUSEN		207
CERATOCOMBIDAE	1	<i>*Nabis rugosus</i> L.
<i>*Ceratocombus coleopratus</i> Zett.		ANTHOCORIDAE
NEPIDAE	1	7
<i>Ranatra linearis</i> L.		<i>*Acomporis alpinus</i> Reut.
CORIXIDAE	4	? <i>Acomporis pygmaeus</i> Fall. ?
<i>Hesperocorixa moesta</i> Fieb.		<i>*Anthocoris nemorum</i> L.
<i>Sigara striata</i> L.		<i>*Temnostethus pusillus</i> H.-S.
<i>Sigara falleni</i> Fieb.		<i>*Orius majusculus</i> Reut.
<i>Sigara fossarum</i> Lch.		<i>*Orius minutus</i> L.
VELIIDAE	2	<i>Lyctocoris campestris</i> F.
<i>Microvelia reticulata</i> Burm.		REDUVIIDAE
? <i>Velia caprai caprai</i> Tam. ?		5
GERRIDAE	1	<i>Empicoris culiciformis</i> De G.
<i>Gerris argentatus</i> Schum.		<i>Empicoris vagabundus</i> L.
SALDIDAE	3	<i>Phymata crassipes</i> F.
<i>Saldula c-album</i> Fieb.		<i>Pygolampis bidentata</i> Gz.
<i>*Saldula orthochila</i> Fieb.		<i>Rhynocoris iracundus</i> Pd.
<i>Saldula saltatoria</i> L.		MIRIDAE
TINGIDAE	10	76
<i>*Acalypta carinata</i> Pz.		<i>*Campyloneura virgula</i> H.-S.
<i>*Acalypta marginata</i> Wff.		<i>*Dicyphus annulatus</i> Wff.
<i>Acalypta parvula</i> Fall.		<i>*Dicyphus globulifer</i> Fall.
<i>Agramma laetum</i> Fall.		<i>*Deraeocoris morio</i> Boh.
<i>*Campylosteira verna</i> Fall.		<i>*Deraeocoris ruber</i> L.
<i>*Catoplatus fabricii</i> Stål		<i>Deraeocoris lutescens</i> Schill.
<i>Dictyla echii</i> Schrank		<i>Leptopterna dolobrata</i> L.
<i>Dictyla lupuli</i> H.-S.		<i>Stenodema holsata</i> F.
<i>*Kalama tricornis</i> Schrank		<i>*Stenodema laevigata</i> L.
<i>*Physatocheila harwoodi</i> China		<i>*Megaloceraea recticornis</i> Geoffr.
NABIDAE	8	<i>*Notostira elongata</i> Geoffr.
<i>Prostemma guttula</i> F.		<i>*Notostira erratica</i> L.
<i>*Himacerus mirmicoides</i> O.C.		<i>*Trigonotylus caelestialium</i> Kirk
<i>*Nabis flavomarginatus</i> Sz.		<i>*Phytocoris longipennis</i> Flor
<i>*Nabis brevis</i> Sz.		<i>*Adelphocoris lineolatus</i> Gz.
<i>*Nabis ferus</i> L.		<i>*Adelphocoris seticornis</i> F.
<i>*Nabis pseudoferus</i> Rem.		<i>*Brachycoleus pilicornis</i> Pz.
<i>*Nabis punctatus</i> A. C.		<i>*Calocoris affinis</i> H.-S.
		<i>Calocoris roseomaculatus</i> De G.
		<i>*Closterotomus biclavatus</i> H.-S.
		<i>Closterotomus fulvomaculatus</i> De G.
		<i>Closterotomus norvegicus</i> Gm.

MIRIDAE	cont.	
<p><i>Grypocoris sexguttatus</i> F. <i>Rhabdomiris striatellus</i> F. <i>*Hadrodemus m-flavum</i> Gz. <i>Stenotus binotatus</i> F. <i>*Dichrooscytus intermedius</i> Reut. <i>Apolygus lucorum</i> Mey.-D. <i>*Apolygus spinolae</i> Mey.-D. <i>*Lygocoris pabulinus</i> L. <i>*Lygus pratensis</i> L. <i>*Lygus rugulipennis</i> Popp. <i>*Lygus wagneri</i> Rem. <i>*Orthops basalis</i> A. C. <i>*Orthops campestris</i> L. <i>*Orthops kalmii</i> L. <i>Pinalitus rubricatus</i> Fall. <i>Charagochilus gyllenhalii</i> Fall. <i>Polymerus holosericeus</i> Hahn <i>*Polymerus nigrita</i> Fall. <i>*Polymerus unifasciatus</i> F. <i>*Polymerus microphtalmus</i> E. Wgn. <i>*Capsodes gothicus gothicus</i> L. <i>*Capsus ater</i> L. <i>*Halticus apterus apterus</i> L. <i>*Orthocephalus coriaceus</i> F. <i>*Orthocephalus saltator</i> Hahn <i>Strongylocoris luridus</i> Fall. <i>*Strongylocoris steganooides</i> J. Shlbg. <i>Cyllecoris histrionius</i> L. <i>Globiceps sphaegiformis</i> Rossi <i>*Globiceps flavomaculatus</i> F. <i>*Globiceps fulvicollis</i> Jak. <i>Heterocordylus tumidicornis</i> H.-S. <i>*Malacocoris chlorizans</i> Pz. <i>Orthotylus nassatus</i> F. <i>Orthotylus viridinervis</i> Kb. <i>*Hallodapus rufescens</i> Burm. <i>*Ambylytus nasutus</i> Kb. <i>*Atractotomus magnicornis</i> Fall. <i>*Atractotomus parvulus</i> Reut. <i>Campylomma verbasci</i> Mey.-D. <i>*Chlamydatus pulicarius</i> Fall. <i>*Chlamydatus pullus</i> Reut. <i>Criocoris crassicornis</i> Hahn <i>Hoplomachus thunbergii</i> Fall. <i>*Macrotylus herrichi</i> Reut. <i>*Megalocoleus molliculus</i> Fall. <i>Monosynamma bohemanni</i> Fall. <i>Oncotylus viridiflavus</i> Gz. <i>*Phoenicocoris obscurellus</i> Fall.</p>		
ARADIDAE		3
<p><i>*Aradus cinnamomeus</i> Pz. <i>Aradus conspicuus</i> H.-S. <i>Aradus depressus</i> F.</p>		
PIESMATIDAE		2
<p><i>Piesma capitatum</i> Wff. <i>*Piesma maculatum</i> Lap.</p>		
BERYTIDAE		7
<p><i>Berytinus clavipes</i> F. <i>Berytinus minor</i> H.-S. <i>*Berytinus crassipes</i> H.-S. <i>*Berytinus montivagus</i> Mey.-D. <i>*Berytinus signoreti</i> Fieb. <i>*Gampsocoris punctipes</i> Germ. <i>Neides tipularius</i> L.</p>		
LYGAEIDAE		25
<p><i>*Nysius ericae</i> Schill. <i>*Ischnodemus sabuleti</i> Fall. <i>Macroplax preysleri</i> Fieb. <i>Heterogaster urticae</i> F. <i>*Platyplax salviae</i> Schill. <i>Plinthisus brevipennis</i> Latr. <i>Tropistethus holosericeus</i> Sz. <i>*Drymus latus</i> Doug. & Sc. <i>*Drymus ryeii</i> Doug. & Sc. <i>*Drymus sylvaticus</i> F. <i>*Gastrodes abietum</i> Bgrth. <i>*Scolopostethus affinis</i> Schill. <i>*Scolopostethus thomsoni</i> Reut. <i>?Taphropeltus contractus</i> H.-S. ? <i>*Stygnocoris rusticus</i> Fall. <i>Stygnocoris sabulosus</i> Schill. <i>Pachybrachius fracticollis</i> Schill. <i>Aellopus atratus</i> Gz. <i>*Graptopeltus lynceus</i> F. <i>*Rhyparochromus pini</i> L. <i>Xanthochilus quadratus</i> F. <i>*Peritrechus gracilicornis</i> Put. <i>*Megalonotus chiragra</i> F. <i>Emblethis verbasci</i> F. <i>Pterotmetus staphiliniformis</i> Schill.</p>		
PYRRHOCORIDAE		1
<p><i>*Pyrrhocoris apterus</i> L.</p>		

<p>COREIDAE 5 <i>Gonocerus acuteangulatus</i> Gz. <i>Syromastus rhombeus</i> L. <i>Enoplops scapha</i> F. *<i>Coreus marginatus</i> L. *<i>Coriomeris denticulatus</i> Scop.</p>	<p>ACANTHOSOMATIDAE 2 <i>Elasmucha ferrugata</i> F. <i>Cyphostethus tristriatus</i> F.</p>
<p>ALYDIDAE 1 <i>Alydus calcaratus</i> L.</p>	
<p>RHOPALIDAE 5 *<i>Corizus hyoscyami</i> L. *<i>Rhopalus parumpunctatus</i> Schill. *<i>Rhopalus subrufus</i> Gm. *<i>Stictopleurus abutilon</i> Rossi *<i>Stictopleurus punctatonervosus</i> Gz.</p>	
<p>PLATASPIDAE 1 <i>Coptosoma scutellatum</i> Geoffr.</p>	
<p>CYDNIDAE 5 <i>Sehirus luctuosus</i> Muls. & Rey <i>Tritomegas bicolor</i> L. *<i>Canthophorus impressus</i> Horv. *<i>Adomerus biguttatus</i> L. *<i>Legnotus picipes</i> Fall.</p>	
<p>THYREOCORIDAE 1 *<i>Thyreocoris scarabaeoides</i> L.</p>	
<p>SCUTELLERIDAE 3 <i>Odontoscelis fuliginosa</i> L. *<i>Eurygaster maura</i> L. *<i>Eurygaster testudinaria</i> Geoffr.</p>	
<p>PENTATOMIDAE 18 *<i>Sciocoris macrocephalus</i> Fieb. *<i>Sciocoris microphthalmus</i> Flor <i>Sciocoris umbrinus</i> Wff. *<i>Aelia acuminata</i> L. *<i>Neottiglossa pusilla</i> Gm. *<i>Rubiconia intermedium</i> Wff. *<i>Palomena prasina</i> L. *<i>Palomena viridissima</i> Pd. <i>Dryocoris vernalis</i> Wff. <i>Chlorochroa juniperina</i> L. *<i>Carpocoris fuscispinus</i> Boh. *<i>Carpocoris purpureipennis</i> De G. *<i>Dolycoris baccarum</i> L. *<i>Eurydema oleraceum</i> L. *<i>Pentatoma rufipes</i> L. *<i>Troilus luridus</i> F. <i>Arma custos</i> F. *<i>Zicrona caerulea</i> L.</p>	

LER, ETH collection, pers. comm.). Therefore, we have only updated the name (STICHEL, 1955–1962; HEISS & PÉRICART, 1983; PÉRICART, 1984, 1998; AUKEMA & RIEGER, 1995, 1996, 1999). However, the following four records are problematic:

1. *Velia currens* F. was published in the lists for the Canton of FREY-GESSNER (1864–1866). Much later, *Velia caprai* TAM. and *Velia saulii* TAM. were separated from that species as distinct species (TAMANINI, 1947). It is therefore not clear which of the three species had actually been found. Probably it was *Velia caprai* TAM., which is also very common in the southern part of Baden-Württemberg, while *Velia currens* F. occurs only in Ticino (DETHIER & MATTHEY, 1977).

2. *Acomporis alpinus* REUT. was described in 1875 and separated from *Acomporis pygmaeus* FALL., so we do not know which of the two species had been found.

3. *Taphropeltus hamulatus* THOMS. was described in 1870. After being considered as a synonym of *Taphropeltus contractus* H.-S. it was later separated again; we do not know which of the two species was really collected.

4. *Sehirus morio* L. was published in the lists for the Canton of FREY-GESSNER (1864–1866). At that time *Sehirus luctuosus* MULSANT & REY had not been described as a separate species from *S. morio*. We assume that the record probably deals with *S. luctuosus* MULS. & REY which is found frequently, while *S. morio* L. is seldom found in Central Europe.

Biogeography of rare species

In the following, we discuss our own findings in detail with a critical focus on species on the Red List and those rarely found in Switzerland. Some species on the Red Lists of Germany are very common and widespread in both Baden-Württemberg and Switzerland and will not be discussed here. These are *Saldula c-album* FIEB., *Saldula orthochila* FIEB., *Acalypta marginata* WFF., *Catoplatus fabricii* STAL., *Kalama tricornis* SCHR., *Acomporis alpinus* REUT., *Calocoris alpestris* MEY.-D., *Grypocoris sexguttatus* F., *Macrotylus herrichi* REUT., *Berytinus signoreti* FIEB., *Ischnodemus sabuleti* FALL., *Adomerus biguttatus* L., *Legnotus picipes* FALL., *Rubiconia intermedium* WFF. and *Zicrona caerulea* L.

Canton Schaffhausen – We compared our results for Schaffhausen (Tab. 5) with those of Baden-Württemberg and the dry habitats of the Hegau (Landkreis Konstanz). For the Landkreis Konstanz 411 bug species have been recorded (HECKMANN, 1989, 1990, 1993, 1996, 1999; HECKMANN & RIEGER, in prep.; HECKMANN, unpublished data). In total 21 species of the Schaffhausen bug fauna have not been found in the Landkreis Konstanz so far.

Eight of our recently recorded species have not been found in the Landkreis Konstanz, though it is a well investigated region. All species except *Campylosteira verna* FALL., *Physatocheila harwoodi* CHINA and *Sciocoris microphthalmus* FLOR. are xero-thermophilic species and most are on the Red List of Germany: *Saldula orthochila* FIEB., *Campylosteira verna* FALL., *Physatocheila harwoodi* CHINA, *Brachycoleus pilicornis* PZ., *Hallodapus rufescens* BURM., *Syromastus rhombeus* L., *Sciocoris macrocephalus* FIEB. and *Sciocoris microphthalmus* FLOR.

Thirteen species are mentioned in the early records but were neither found by us nor in the Landkreis Konstanz: *Hesperocorixa moesta* FIEB., *Sigara fossarum* LCH., *Dictyla lupuli* H.-S., *Calocoris roseomaculatus* DE G., *Strongylocoris luridus* FALL., *Globiceps sphaegiformis* ROSSI, *Neides tipularius* L., *Aellopus atratus* F., *Xanthochilus quadratus* F., *Syromastus rhombeus* L., *Sciocoris umbrinus* WFF.,




Fig. 2. *Physatocheila harwoodi*. The arrow shows the narrow black area of the elytra which is much wider in the other species of the genus. The body length is about 3.5 mm. (Photo: M. DI GIULIO).

Chlorochroa juniperina L. and *Elasmucha ferrugata* F. Also *Neides tipularius* L., *Aellopus atratus* F., *Xanthochilus quadratus* F., *Syromastus rhombeus* L., and *Chlorochroa juniperina* L. are xero-thermophilic and are mostly Red List species in Germany. *Strongylocoris luridus* FALL. is found in the Black Forest (RIEGER & STRAUSS, 1992) and lives on *Jasione montana*. *Elasmucha ferrugata* F. also is known from the nearby Wutachschlucht in the Black Forest (KLESS, 1961) and lives on *Vaccinium myrtillus*. The distribution of the other eleven species in Baden-Württemberg corresponds with hot and dry habitats along the Rhine Valley north of Basel (HECKMANN, 1996).

The species of special interest found by us are:

1. *Ceratocombus coleoptratus* (ZETTERSTEDT, 1819): In Switzerland this species has been found only recently in the Canton Ticino (OTTO, 1994). It was found several times in Baden-Württemberg and is common in the Landkreis Konstanz (HECKMANN & RIEGER, in prep.). It is rarely collected because of its small size and the fact that it occurs hidden in moss. Mostly it is recorded by pitfall traps. In Bayern it has been put on the Red List.

2. *Acalypta carinata* (PANZER, 1806) is often found in the same habitats as *Ceratocombus coleoptratus* ZETT. and can be caught easily with pitfall-traps. Records are known for Vaud, Bern, Zürich and Wallis (PÉRICART, 1983). It is on the Red List for Baden-Württemberg and Bayern but seems to be common in the south of Baden-Württemberg (RIEGER, 1981) and the Landkreis Konstanz (HECKMANN, 1990).

3. *Campylosteira verna* (FALLÉN, 1826) lives hidden between mosses and lichens. In Switzerland it is known from Bern and Graubünden (PÉRICART, 1983)

but few records are known for Baden-Württemberg (HÜEBER, 1891; MEESS, 1907; FISCHER, 1961; RIEGER, 1981). It is on the Red List of Bayern and Germany.

4. *Physatocheila harwoodi* CHINA, 1936 (Fig. 2): New record for Switzerland! The species lives on *Acer pseudoplatanus* and is recorded very rarely. It is known from several places (HEISS, 1978) in Tyrol (Austria), from Baden-Württemberg (RIEGER, 1981; VOIGT, 1983) and from Bayern (SINGER, 1952). This new record corresponds to the known distribution area.

5. *Deraeocoris morio* (BOHEMAN, 1852): This xero-thermophilic species lives on *Thymus* and is known only from the Ticino (GÖLLNER-SCHIEDING & REZBANYAI-RESER, 1992) and several places in Baden-Württemberg and in the Landkreis Konstanz (HECKMANN & RIEGER, in prep.).

6. *Brachycoleus pilicornis pilicornis* (PANZER, 1805): This thermophilic species lives on *Euphorbia verrucosa*. It is known from the Cantons of Basel, Zürich, Aargau (FREY-GESSNER, 1864–1866) and Thurgau (BLÖCHLINGER, pers. comm.). In Baden-Württemberg it is known from several places and also from warm places in the nearby Wutachschlucht in the Black Forest (KLESS, 1961).

7. *Polymerus microphthalmus* (WAGNER, 1951): New record for Switzerland! This species lives on *Galium*, often together with *P. unifasciatus* F., and is common in the Hegau and rarely found in other parts of Baden-Württemberg. Probably there is no Swiss record because it has been confused with *P. unifasciatus* F.

8. *Strongylocoris steganoides* (J. SAHLBERG, 1875) was synonymised with *S. leucocephalus* by REUTER (1888). KIRITSHENKO (1951) revised the genus and separated these two species. WAGNER (1973) treats it as a subspecies, and it might therefore have been misidentified (RIEGER, 1997). In Baden-Württemberg *S. steganoides* is widespread and very common whereas *S. leucocephalus* is a rare species. In the Landkreis Konstanz only *S. steganoides* has been found (HECKMANN & RIEGER, in prep.). We assume that the same is true for Switzerland and the records of *S. leucocephalus* should be checked.

9. *Hallodapus rufescens* (BURMEISTER, 1835) is known in Switzerland only from the Cantons Vaud, Bern (FREY-GESSNER, 1864–1866), Wallis (CERUTTI, 1937a) and Graubünden (Valbella; DI GIULIO, unpublished data). In Baden-Württemberg it is only known from the Suebian Alb (RIEGER, 1976) and from the Kniebis in the northern Black Forest (RIEGER, pers. comm.).

10. *Berytinus crassipes* (HERRICH-SCHÄFFER, 1835) lives on *Cerastium*. Records are known from Aargau and Vaud in Switzerland (PÉRICART, 1984), the Wollmatinger Ried in the Landkreis Konstanz (HECKMANN, 1990) and a few other places in Baden-Württemberg (KLESS, 1961; RIEGER, 1976; HECKMANN, 1996).

11. *Berytinus montivagus* (MEYER-DÜR, 1841): This xero-thermophilic species is known in Switzerland only from Graubünden (FREY-GESSNER, 1871), Aargau (PÉRICART, 1984) and Neuchâtel (BARBALAT, 1991) but is probably more widespread. It is known from the Hohentwiel in the Landkreis Konstanz and from other places in Baden-Württemberg (SCHMID, 1967; HECKMANN & RIEGER, in prep.).

12. *Drymus latus* DOUGLAS & SCOTT, 1871: This is the second record for Switzerland. The species has only been recorded in the Canton Basel (PÉRICART, 1998). There are few records in the Landkreis Konstanz (HECKMANN, 1990; HECKMANN & RIEGER, in prep.) and from few other places in Baden-Württemberg (RIEGER & STRAUSS, 1992).

13. *Peritrechus gracilicornis* PUTON, 1877: This xerophilic species is known in Switzerland from Wallis and Ticino (PÉRICART, 1998). In Baden-Württemberg it




Fig. 3. *Sciocoris macrocephalus* FIEB. (upper part) can be distinguished from *Sciocoris microphthalmus* FLOR (lower part) by the sinuated lateral margins of the head (arrows) and by the longer eye-stalks. The body length is about 6 mm. (Photo: R. HECKMANN and M. DI GIULIO).

is mostly found along the valleys of the Rhine and the Neckar and also in the Hegau (HECKMANN, 1996).

14. *Canthophorus impressus* HORVATH, 1881: This is the third reliable record of this species for Switzerland. It has been found previously in the Cantons Obwalden (GÖLLNER-SCHIEDING, 1989b) and Ticino (GÖLLNER-SCHIEDING & REZBANYAI-

RESER, 1992). The species lives on *Thesium* and is widespread in Baden-Württemberg (RIEGER & STRAUSS, 1992) and has been found in the Wollmatinger Ried in the Landkreis Konstanz (HECKMANN, 1999). Before RIEGER (1997) the determination of the females was unsure and therefore the records of *C. dubius* SCOP. in FREY-GESSNER (1864–1866) were probably *C. impressus* as this is the more common species in Baden-Württemberg.

15. *Sciocoris microphthalmus* FLOR, 1860 (Fig. 3): This species is known from Ticino (RAMPAZZI & DETHIER, 1997) and Tyrol (HEISS, 1977); however, in Baden-Württemberg it is common (RIEGER, pers. comm)

16. *Sciocoris macrocephalus* FIEBER, 1851 (Fig. 3): Reliable records exist only from Wallis (VOELLMY & SAUTER, 1983), Neuchâtel (BARBALAT, 1991), Ticino (GÖLLNER-SCHIEDING & REZBANYAI-RESER, 1992) and Thurgau (Coll. BLÖCHLINGER, Müllheim). There are no checked records from Baden-Württemberg (RIEGER, 1996) and the Fürstentum Liechtenstein (BERNHARDT, 1992), but the species is found frequently in Tyrol (HEISS, 1977). Reports also exist from Vaud, Wallis, Aargau, Zürich and Graubünden (FREY-GESSNER, 1864–1866). However, it remains uncertain whether these reports differentiate between *S. macrocephalus* and *S. microphthalmus*.

Canton Luzern – Four interesting species were found in the Canton Luzern:

1. *Acalypta carinata* (PANZER, 1806): see discussion of Canton Schaffhausen.

2. *Himacerus major* (A. COSTA, 1842): In Switzerland there are only two records of this species, from the Cantons Wallis and Basel. A further record (PÉRICART, 1987; DETHIER & PÉRICART, 1988) is attributed to a mistake of the name Kaiserstuhl. The species was actually found in Baden-Württemberg (Kaiserstuhl in Baden) and not in the Canton Aargau. Several records come from Baden-Württemberg along the Rhine Valley (HECKMANN, 1996) and from the Landkreis Konstanz. This is the first record for Central Switzerland.

3. *Pinalitus atomarius* (MEYER-DÜR, 1843): In Switzerland this alpine-boreal species which lives on *Picea* is known from Bern, Basel and Aargau (FREY-GESSNER, 1864–1866) and recently from Ticino (RAMPAZZI & DETHIER, 1997). The distribution in Baden-Württemberg is restricted to the Black Forest (RIEGER & STRAUSS, 1992; HECKMANN, 1996) and to the Suebian Alb (RIEGER, pers. comm.).

4. *Pinalitus visicola* (PUTON, 1888): This species lives on *Viscum album* and has been recorded in Switzerland only in the Canton Wallis (CERUTTI, 1937b). In the Landkreis Konstanz, the distribution corresponds with that of *Viscum*. Its distribution for Baden-Württemberg has been described in HECKMANN & RIEGER (in prep.). The sparse records reflect the dependence of the species on *Viscum* and the fact that its habitat mostly in the canopy, is rarely sampled.

CONCLUSIONS

The Red List of Baden-Württemberg is very helpful in interpreting the rare species of Switzerland. Some xero-thermophilic species recorded more or less frequently in the south of Baden-Württemberg especially in the valley of the Rhine and Hegau also occur in the Canton Schaffhausen. The same species are recorded in the southern part of Switzerland and in the Cantons Basel and Aargau, but not in the Innerschweiz and Mittelland. These species, which are good candidates for a Red List of Switzerland, include: *Deraeocoris morio* BOH., *Hallodapus rufescens* BURM., *Berytinus crassipes* H.-S., *Berytinus montivagus* MEY.-D., *Berytinus signoreti* FIEB., *Drymus latus* DOUG. & SC., *Peritrechus gracilicornis* PUT., *Emblethis ver-*

basci F., *Sciocoris macrocephalus* Fieb., and *Sciocoris microphthalmus* FLOR. From our findings in Luzern *Himacerus major* A. C. and *Pinalitus visicola* PUT. should be added to the list.

Many species found in the ancient records of MEESS (1900, 1907) and FREY-GESSNER (1864–1866) appear to have decreased in abundance during the last century, e.g. *Neides tipularius* L., *Aellopus atratus* F., *Xanthochilus quadratus* F., *Sciocoris umbrinus* WFF., *Chlorochroa juniperina* L. and *Elasmucha ferrugata* F. However, the sparse data available on the distribution of heteropteran bugs do not allow a thorough analysis of the situation in Switzerland. The data of Baden-Württemberg, however, support the conclusion that these species decreased in the last century (HECKMANN, 1996). We assume that this decline was a consequence of increasing intensification of agricultural production in both Germany and Switzerland.

ACKNOWLEDGMENTS

We thank B. Merz for general assistance, H. BLÖCHLINGER, Ch. RIEGER and S. STUDER for permission to use unpublished data, to P. J. EDWARDS, Ch. RIEGER and L. RESER for their critical reading of the manuscript, to A. MÜLLER for the permission to use the material of the Entomological Collection of the Swiss Federal Institute of Technology (ETHZ).

This work was funded by the Swiss National Science Foundation (5001-044634/1 and 5001-44636) and the Canton Schaffhausen (Kantonales Planungs- und Naturschutzamt).

ZUSAMMENFASSUNG

Im Schaffhauser Randen (Kanton Schaffhausen) wurden zwischen 1997 und 1999 drei verschiedene Wiesentypen untersucht. Dabei wurden verschiedene Sammelmethode angewendet, wie Kescherfang, Bodenfallen und ein Sauggerät. Insgesamt wurden 45 Wiesen untersucht und 140 Wanzenarten gefunden. Auf dem Gemeindegebiet Ruswil/Buttisholz (Kanton Luzern) wurden 31 extensiv bewirtschaftete Wiesen untersucht und dabei 78 Arten festgestellt. In unseren Untersuchungen fanden wir 118 neue Arten für den Kanton Schaffhausen und 26 für den Kanton Luzern. Anhand der Literatur und aufgrund eigener Funde stellten wir für beide Kantone Artenlisten zusammen. Aus dem Kanton Schaffhausen sind bisher 207 Wanzenarten bekannt, aus dem Kanton Luzern 201 Arten. Die Biogeographie ausgewählter Arten wird diskutiert.

Drei Arten aus dem Schaffhauser Randen sind Neunachweise für die Schweiz: *Physatocheila harwoodi* CHINA, 1936, *Strongylocoris steganoides* (J. SAHLBERG, 1875) und *Polymerus microphthalmus* (WAGNER, 1951). Die bisherigen Funde von *Strongylocoris leucocephalus* sollten überarbeitet werden.

REFERENCES

- ACHTZIGER, R. 1995. Die Struktur von Insektengemeinschaften an Gehölzen: Die Hemipteren-Fauna als Beispiel für die Biodiversität von Hecken- und Waldrandökosystemen. *Bayreuther Forum Ökologie* 20: 1–183.
- ACHTZIGER, R., SCHOLZE, W., & SCHUSTER, G. 1992. *Rote Liste gefährdeter Landwanzen (Heteroptera, Geocorisae) Bayerns*. In: Schriftenreihe Bayer. Landesamt für Umweltschutz 111, pp. 87–95. München.
- ARNOLD, A.J. 1994. Insect suction sampling without nets, bags or filters. *Crop prot.* 13(1): 73–76.
- AUKEMA, B. & RIEGER, C. 1995. *Catalogue of the Heteroptera of the Palaearctic Region. Vol. 1. Enicocephalomorpha, Dipsocoromorpha, Nepomorpha, Gerromorpha and Leptopodomorpha*. The Netherlands Entomological Society, 222 pp.
- AUKEMA, B. & RIEGER, C. 1996. *Catalogue of the Heteroptera of the Palaearctic Region. Vol 2. Cimicomorpha I*. The Netherlands Entomological Society, 361 pp.
- AUKEMA, B. & RIEGER, C. 1999. *Catalogue of the Heteroptera of the Palaearctic Region. Vol 3. Cimicomorpha II*. The Netherlands Entomological Society, 577 pp.
- BARBALAT, S. 1991. Inventaire des Coléoptères Carabidés et des Hétéroptères de cinq talus du Val-de-Ruz (Canton de Neuchâtel, Suisse). *Bull. Soc. Neu. Sci. Nat.* 114: 7–17.
- BERNHARDT, K.-G. 1992. Die Wanzen (Heteroptera) des Fürstentums Liechtenstein. I. Teil: Die Wanzenfauna des ausseralpinen Raumes. *Ber. Bot.-Zool. Ges. Liechtenstein-Sargans-Werdenberg* 19: 295–325.

- CERUTTI, N. 1937a. Captures intéressantes d'Hémiptères du Valais. *Mitt. Schweiz. Ent. Ges.* 17(1/2): 30–32.
- CERUTTI, N. 1937b. Captures intéressantes d'Hémiptères du Valais (2e liste). *Mitt. Schweiz. Ent. Ges.* 17(4): 168–172.
- CURTIS, D.J. 1980. Pitfalls in spider community studies (Arachnida, Araneae). *J. Archnol.* 8: 271–280.
- DETHIER, M. & MATTHEY, W. 1977. Contribution à la connaissance des Hétéroptères aquatiques de Suisse. *Revue suisse Zool.* 84 (3): 583–591.
- DETHIER, M. & PÉRICART, J. 1988. Les Hétéroptères Nabidae de Suisse. *Mitt. Schweiz. Ent. Ges.* 61: 157–166.
- DETHIER, M. & PÉRICART, J. 1990. Les Hétéroptères Leptodomorpha de Suisse. *Mitt. Schweiz. Ent. Ges.* 63: 33–42.
- DI GIULIO, M., EDWARDS, P.J., & MEISTER, E. (in prep.) Enhancing insect diversity in agricultural grasslands: The roles of management and landscape structure.
- DOLLING, W.R. 1991. *The Hemiptera*. Oxford University Press, Oxford, 266 pp.
- DUELLI, P. & OBRIST, M. 1998. In search of the best correlates for local organismal biodiversity in cultivated areas. *Biodiversity and Conservation* 7: 297–309.
- FAUVEL, G. 1999. Diversity of Heteroptera in agroecosystems: role of sustainability and bioindication. *Agricult. Ecosys. Environ.* 74: 275–303.
- FISCHER, H. 1961. Die Tierwelt Schwabens I. Teil. Die Wanzen. 13. *Ber. Naturf. Ges. Augsburg* 72: 1–32.
- FREY-GESSNER, E. 1864–1866. Verzeichnis der schweizerischen Insekten. 1. Hemiptera. *Mitt. Schweiz. Ent. Ges.* 1(6): 195–203; 1(7): 225–244; 1(9): 304–310; 2(1): 7–30; 2(3): 115–133.
- FREY-GESSNER, E. 1871. Sammelbericht aus den Jahren 1869 und 1870. *Mitt. Schweiz. Ent. Ges.* 3(7): 313–326.
- GÖLLNER-SCHIEDING, U. 1981. Die Insektenfauna des Hochmoores Balmoos bei Halse, Kanton Luzern. X. Heteroptera (Wanzen). *Ent. Ber. Luzern* 5: 83–85.
- GÖLLNER-SCHIEDING, U. 1982. Zur Insektenfauna der Umgebung der Vogelwarte Sempach, Kanton Luzern. XI. Heteroptera (Wanzen). *Ent. Ber. Luzern* 8: 83–86.
- GÖLLNER-SCHIEDING, U. 1989a. Ergebnisse von Lichtfängen in Berlin aus den Jahren 1981–1986 .1. Heteroptera. Teil I: Landwanzen (Cimicomorpha et Pentatomorpha) (Insecta). *Faun. Abh. Mus. Tierkd. Dresden* 16: 111–123.
- GÖLLNER-SCHIEDING, U. 1989b. Angaben zur Wanzenfauna der Zentralschweiz (Heteroptera). *Ent. Ber. Luzern* 22: 107–116.
- GÖLLNER-SCHIEDING, U. 1990. Zur Insektenfauna vom Vogelmoos (775 m) bei Neudorf, Kanton Luzern. III. Heteroptera (Wanzen). *Ent. Ber. Luzern* 24: 115–122.
- GÖLLNER-SCHIEDING, U. & REZBANYAI-RESER, L. 1992. Zur Wanzenfauna des Monte Generoso, Kanton Tessin, Südschweiz (Heteroptera). *Ent. Ber. Luzern* 28: 15–36.
- GÜNTHER, H. & SCHUSTER, G. 1990. Verzeichnis der Wanzen Mitteleuropas (Heteroptera). *Dtsch. ent. Z., N.F.* 37: 361–396.
- GÜNTHER, H., HOFFMANN, H.-J., MELBER, A., REMANE, R., SIMON, H., & WINKELMANN, H. 1998. Rote Liste der Wanzen (Heteroptera). In: Bundesamt für Naturschutz (ed.): Rote Liste gefährdeter Tiere Deutschlands. *Schriftenreihe Landschaftspflege Naturschutz* 55: 235–242.
- HECKMANN, R. 1989. Wanzen (Heteroptera). In: DIENST, M. & JACOBY, H. (eds), *Jahresbericht 1988 über das Naturschutzgebiet Wollmatinger Ried-Untersee-Gnadensee*, pp. 21–26. Konstanz.
- HECKMANN, R. 1990. Wanzen (Heteroptera). In: DIENST, M. & JACOBY, H. (eds), *Jahresbericht 1989 über das Naturschutzgebiet Wollmatinger Ried-Untersee-Gnadensee*, pp. 22–25. Konstanz.
- HECKMANN, R. 1993. Wanzen (Heteroptera). In: DIENST, M. & JACOBY, H. (eds), *Jahresbericht 1992 über das Naturschutzgebiet Wollmatinger Ried-Untersee-Gnadensee*, pp. 25–29. Konstanz.
- HECKMANN, R. 1996. Katalog der Wanzen aus Baden-Württemberg in der Sammlung des Staatlichen Museums für Naturkunde Karlsruhe (Insecta, Heteroptera). *Carolinea Beiheft* 10, Karlsruhe, 146 pp.
- HECKMANN, R. 1999. Wanzen (Heteroptera). In: KLEIN, E. & JACOBY, H. (eds), *Jahresbericht 1998 über das Naturschutzgebiet Wollmatinger Ried-Untersee-Gnadensee*, pp. 24–27. Konstanz.
- HECKMANN, R. & RIEGER, C. (in prep.) Neunachweise seltener und bisher nicht bekannter Wanzen in Baden-Württemberg II (Insecta, Heteroptera). *Carolinea*.
- HEISS, E. 1977. Zur Heteropterenfauna Nordtirols (Ins.: Het.) VI: Pentatomoidea. *Veröff. Mus. Ferdinandeum* 57: 53–77.
- HEISS, E. 1978. Zur Heteropterenfauna Nordtirols (Ins.: Het.) VII: Tingidae. *Ber. nat.-med. Verein Innsbruck* 65: 73–84.
- HEISS, E. & PÉRICART, J. 1983. Revision of Palaearctic Piesmatidae (Heteroptera). *Mitt. Münch. Ent. Ges.* 73: 61–171.

- HOFMÄNNER, B. 1928. Beiträge zur Kenntnis der thurgauischen Halbflügler (Hemipteren). *Mitt. Thurg. Natforsch. Ges.* 27: 1–16.
- HÜEBER, T. 1891. ROSER's Württembergische Hemipteren-Fauna. *Jh. Ver. vaterl. Naturkde. Württemberg* 47: 149–169.
- ISLER-HÜBSCHER, K. 1980. Beiträge 1976 zu Georg Kummers "Flora des Kantons Schaffhausen" mit Berücksichtigung der Grenzgebiete. *Mitt. naturf. Ges. Schaffhausen* 31: 7–121.
- JUDEX, M. 1999. Einfluss der Wald-Distanz auf die Diversität der Heteroptera in extensiv bewirtschafteten Wiesen. Diplomarbeit ETH Zürich.
- KIRITSHENKO, A.N. 1951. True bugs of the European Part of the USSR (Hemiptera): Key and Bibliography. *Opredel. Faune SSSR* 42, Moskva/Leningrad [russ.], pp. 423.
- KLESS, J. 1961. Die Käfer und Wanzen der Wutachschlucht. *Mitt. bad. Landesver. Naturkde. Naturschutz N.F.* 8(1): 79–152.
- MEESS, A. 1900. Erster Beitrag zur Hemipteren-Fauna Badens. *Mitt. Bad. Zool. Ver.* 2: 37–93.
- MEESS, A. 1907. 2. Beitrag zur Kenntnis der Hemipteren-Fauna Badens. *Mitt. Bad. Zool. Ver.* 18: 130–151.
- MEYER-DÜR, L.R. 1843. *Verzeichnis der in der Schweiz einheimischen Rhynchoten (Hemiptera LINN.). I. Die Familie der Capsini.* Jent & Gassmann, Solothurn, 120 pp.
- MORRIS, M.G. 1969. Differences between the invertebrate faunas of grazed and ungrazed chalk grasslands. III. The heteropterous fauna. *J. Appl. Ecol.* 6: 475–487.
- MORRIS, M.G. 1979. Responses of grassland invertebrates to management by cutting. II. Heteroptera. *J. Appl. Ecol.* 16: 417–432.
- OTTO, A. 1992. Zur Landwanzenfauna der Magadino-Ebene, Kanton Tessin (Heteroptera: Geocoridae). *Ent. Ber. Luzern* 28: 37–44.
- OTTO, A. 1994. Für die Schweiz neue oder selten gesammelte Wanzen-Arten (Heteroptera). *Mitt. schweiz. Ent. Ges.* 67: 189–197.
- OTTO, A. 1995. Für die Schweiz neue oder selten gesammelte Wanzen-Arten (Heteroptera) – Zweiter Beitrag. *Mitt. schweiz. Ent. Ges.* 68: 137–142.
- OTTO, A. 1996. *Die Wanzenfauna montaner Magerwiesen und Grünbrachen im Kanton Tessin (Insecta: Heteroptera).* Dissertation ETH Zürich Nr. 11457, 212 pp.
- OTTO, A. & BÜRKI, H.-M. 1996. *Phytocoris (Ktenocoris) singeri* E. WAGNER, eine neue Art für die schweizerische Wanzenfauna (Heteroptera). *Ent. Ber. Luzern* 35: 47–48.
- OTTO, A. & REZBANYAI-RESER, L. 1996. Zur Wanzenfauna der Insel Brissago, Kanton Tessin (Heteroptera). *Ent. Ber. Luzern* 35: 49–58.
- PÉRICART, J. 1972. *Hémiptères Anthocoridae, Cimicidae et Microphysidae de l'Ouest-Paléarctique.* Faune de l'Europe et du Bassin Méditerranéen 7, Paris, 402 pp.
- PÉRICART, J. 1983. *Hémiptères Tingidae Euro-méditerranéens.* Faune de France 69. Féd. franç. Soc. Sci. nat., Paris, 620 pp.
- PÉRICART, J. 1984. *Hémiptères Berytidae Euro-méditerranéens.* Faune de France 70. Féd. franç. Soc. Sci. nat., Paris, 171 pp.
- PÉRICART, J. 1987. *Hémiptères Nabidae d'Europe occidentale et du Maghreb.* Faune de France 71. Féd. franç. Soc. Sci. nat., Paris, 185 pp.
- PÉRICART, J. 1990. *Hémiptères Saldidae et Leptopodidae d'Europe occidentale et du Maghreb.* Faune de France 77. Féd. franç. Soc. Sci. nat., Paris, 238 pp.
- PÉRICART, J. 1998. *Hémiptères Lygaeidae Euro-méditerranéens Vol. 1, 2 & 3.* Faune de France 84 A, B, C. Féd. franç. Soc. Sci. nat., Paris.
- RAMPAZZI, F. & DETHIER, M. 1997. Gli Eterotteri (Insecta: Heteroptera) delle torbiere a sfagni del Cantone Ticino e del Moesano (Val Calanca e Val Mesolcina – GR), Svizzera. *Mitt. Schweiz. Ent. Ges.* 70: 419–439.
- REUTER, O.M. 1888. Revisio synonymica Heteropterorum palaearticorum quae descripserunt auctores vetustiores (Linnaeus 1758 – Latreille 1806). *Acta Soc. Scient. Fenn.* 15: 241–315.
- REZBANYAI, L. 1980. Fauna Centrohelveticica. Die Insektenfauna des Hochmoores Balmoos bei Hasle, Kanton Luzern I. Allgemeines. *Ent. Ber. Luzern* 3: 3–14.
- REZBANYAI-RESER, L. 1993. Elenco attuale degli Eterotteri del canton Ticino, Svizzera meridionale (Insecta: Heteroptera). *Boll. Soc. Tic. Sci. Natur.* 81(1): 97–105.
- REZBANYAI-RESER, L. 1997. Nachträge zur Wanzenfauna des Monte Generoso, Kanton Tessin, Südschweiz (Heteroptera). *Ent. Ber. Luzern* 37: 109–110.
- RIEGER, C. 1976. Die Wanzenfauna des mittleren Neckartales und der angrenzenden Albhochfläche (Landkreise Nürtingen, Reutlingen, Tübingen) 3. Nachtrag. *Veröff. Naturschutz Landschaftspflege Bad.-Württ.* 43: 162–169.
- RIEGER, C. 1981. Ergänzungen zur Faunistik und Biologie einiger Netzwanzen in Baden-Württemberg (Heteroptera, Tingidae). *Jh. Ges. Naturkde. Württemberg* 136: 231–240.

- RIEGER, C. 1985. Zur Systematik und Faunistik der Weichwanzen *Orthops kalmii* K. und *Orthops basalis* COSTA (Heteroptera, Miridae). *Veröff. Naturschutz Landschaftspflege Bad.-Württ.* 59/60: 457–465.
- RIEGER, C. 1986. *Vorschlag für eine Rote Liste der Wanzen in Baden-Württemberg (Heteroptera)*. In: *Rote Listen der gefährdeten Tiere und Pflanzen in Baden-Württemberg*. Arbeitsblätter Naturschutz 5, pp. 56–59. Karlsruhe.
- RIEGER, C. 1996. Verzeichnis der bisher in Baden-Württemberg (Bundesrepublik Deutschland) aufgefundenen Wanzen (Insecta: Heteroptera) 1. Fassung. *Jh. Ges. Naturkde. Württemberg* 152: 231–265.
- RIEGER, C. 1997. Ergänzungen zur Faunistik und Systematik einiger Wanzen in Baden-Württemberg (Insecta, Heteroptera) II. *Carolinea* 55: 43–48.
- RIEGER, C. & STRAUSS, G. 1992. Neunachweise seltener und bisher nicht bekannter Wanzen in Baden-Württemberg (Insecta, Heteroptera). *Jh. Ges. Naturkde. Württ.* 147: 247–263.
- SCHIESS-BÜHLER, C. & SCHIESS, H. 1997. Die Tagfalterfauna des Schaffhauser Randens und ihr Wandel im 20. Jahrhundert. *Mitt. natf. Ges. Schaffhausen* 42: 35–106.
- SCHMID, G. 1967. Wanzen aus Baden-Württemberg. *Veröff. Landesst. Nat.schutz u. Landschaftspf. Bad.-Württ.* 35: 89–107.
- SINGER, K. 1952. Die Wanzen (Hemiptera-Heteroptera) des unteren Maingebietes von Hanau bis Würzburg mit Einschluß des Spessarts. *Mitt. naturwiss. Mus. Aschaffenburg, n.F.* 5: 1–128.
- SMA. 1998. *Annalen der Schweizerischen Meteorologischen Anstalt 1997*. SMA, Zürich.
- STICHEL, W. 1955–1962. *Illustrierte Bestimmungstabellen der Wanzen 2. Europa. Bd. 1–4*. Selbstverlag, Berlin-Hermsdorf, 2173 pp.
- TAMANINI, L. 1947. Contributo ad una revisione del genere *Velia* Latr. e descrizione di alcune specie nuove. *Memorie Soc. ent. ital.* 26: 17–74.
- VOELLMY, H. & SAUTER, W. 1983. Wanzen (Heteroptera). *Ergebn. wiss. Unters. Schweiz. Nationalparks N.F.* 12(9): 69–100.
- VOIGT, K. 1983. Erstnachweis einiger Wanzenarten für Baden-Württemberg (Hemiptera, Heteroptera). *Carolinea* 41: 130–131.
- WAGNER, E. 1952. *Blindwanzen oder Miriden*. Die Tierwelt Deutschlands und der angrenzenden Meeresteile 41. G. Fischer Verlag, Jena, 218 pp.
- WAGNER, E. 1966. *Wanzen oder Heteropteren. I. Pentatomorpha*. Die Tierwelt Deutschlands und der angrenzenden Meeresteile 54. G. Fischer Verlag, Jena, 235pp.
- WAGNER, E. 1967. *Wanzen oder Heteropteren. II. Cimicomorpha*. Die Tierwelt Deutschlands und der angrenzenden Meeresteile 55. G. Fischer Verlag, Jena. 179 pp.
- WAGNER, E. 1971. Die Miridae HAHN, 1831, des Mittelmeerraumes und der Makaronesischen Inseln (Hemiptera, Heteroptera). Teil 1. *Ent. Abh. Mus. Tierk. Dresden* 37 (Suppl.): 1–484.
- WAGNER, E. 1973. Die Miridae HAHN, 1831, des Mittelmeerraumes und der Makaronesischen Inseln (Hemiptera, Heteroptera). Teil 2. *Ent. Abh. Mus. Tierk. Dresden* 39 (Suppl.): 1–421.
- WAGNER, E. 1975. Die Miridae HAHN, 1831, des Mittelmeerraumes und der Makaronesischen Inseln (Hemiptera, Heteroptera). Teil 3. *Ent. Abh. Mus. Tierk. Dresden* 40 (Suppl. 1): 1–483.
- WIPRÄCHTIGER, P. 1999a. Die Wasserwanzen in der Wauwiler Ebene. *Ent. Ber. Luzern* 36: 125–133.
- WIPRÄCHTIGER, P. 1999b. Beitrag zur Kenntnis der Wasserwanzenfauna des Möslweiher in Schötz, Kt. Luzern (Heteroptera: Nepomorpha, Gerromorpha). *Ent. Ber. Luzern* 42: 87–90.

(received March 22, 2000; accepted after minor revision July 7, 2000)