

Zeitschrift: Mitteilungen der Schweizerischen Entomologischen Gesellschaft =
Bulletin de la Société Entomologique Suisse = Journal of the Swiss
Entomological Society

Herausgeber: Schweizerische Entomologische Gesellschaft

Band: 80 (2007)

Heft: 3-4

Artikel: Photomicrographs of the male terminalia of Scandinavian Drosophilidae
(Diptera)

Autor: Bächli, Gerhard / Vilela, Carlos R.

DOI: <https://doi.org/10.5169/seals-402949>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 23.12.2024

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Photomicrographs of the male terminalia of Scandinavian Drosophilidae (Diptera)

GERHARD BÄCHLI¹ & CARLOS R. VILELA²

¹ Zoologisches Museum, Universität Zürich, Winterthurerstrasse 190, CH-8057, Zürich, Switzerland.
E-mail: baechli@zm.uzh.ch

² Departamento de Genética e Biologia Evolutiva, Instituto de Biociências, Universidade de São Paulo,
Caixa Postal 11461, São Paulo - SP, 05422-970, Brazil. E-mail: crvilela@ib.usp.br

Black and white photomicrographs of the internal male terminalia of 77 species of Scandinavian Drosophilidae, which were redescribed and illustrated in a book in the series «Fauna Entomologica Scandinavica», are presented. A list of corrections to mistakes in the same book is also included.

Keywords: Internal male genitalia, aedeagus, hypandrium, Fennoscandia, pictures

INTRODUCTION

While preparing illustrations of the male terminalia used in the redescriptions of Scandinavian Drosophilidae (Bächli *et al.* 2004), photomicrographs of all microscope slide mounts were taken as well. Usually, the modern descriptions of drosophilid species are accompanied by line drawings which are two-dimensional interpretations of the three-dimensional terminalia, giving some hints about the connections and potential articulations among its different sclerites, their relative position and detailed structure. For the majority of species, however, the overall shape of the internal terminalia allows their prompt identification. Such standardized views can also be obtained by photomicrographs, which must always be regarded as complementary sources to the published line drawings. We are convinced that the pictures presented below, most of them combined in plates according to genus or species group and enlarged to the same magnification, will be helpful in this regard.

In addition, we take the occasion to present a list of corrections, and clarifications of ambiguities, to the data in the same book.

MATERIAL AND METHODS

Refer to Bächli *et al.* (2004) for details regarding terminalia preparations. The terminalia mounted on microscope slides were photomicrographed in left lateral views using Agfa PAN 25 ASA negative film and a Zeiss Photomicroscope with an 6.3x objective. Unless otherwise stated in the figure captions, all the photomicrographs were taken from the same specimens used for preparing the line drawings. Additional specimens are deposited in the collections of the Zoologisches Museum der Universität Zürich. Detailed label were listed in Tab. 1 of Bächli *et al.* (2004: 332–336), and in the present paper only approximate collection sites, followed by the respective countries, are cited in the captions, except for the additional speci-

mens where detailed data are provided here. Whenever in the same plate, all illustrations were obtained from negatives scanned and enlarged to the same magnification.

RESULTS

An alphabetical list of species epithets dealt with here is presented in Tab. 1, containing references to the figure and page numbers of their respective line drawings published by Bächli *et al.* (2004). For some species, additional pictures of the aedeagus and associated sclerites only, i.e. not including the hypandrium, are included aiming to add clearer views which may be helpful.

Three species were intentionally excluded here, because their respective photomicrographs have already been published: *Drosophila subarctica* Hackman, 1969 and *Drosophila vireni* Bächli, Vilela and Haring, 2002 (see Bächli *et al.* 2002: 319), and *Scaptodrosophila rufifrons* (Loew, 1873) (see Bächli *et al.* 2005: 360).

CORRECTIONS TO BÄCHLI *ET AL.* (2004)

1) In the key to *Drosophila* species, four awkward errors have occurred:

- p. 129 left-hand column, couplet 37, line 14: 42 should be 50
- p. 129, left-hand column, couplet 38, line 22: 50 should be 42
- p. 129, right-hand column, couplet 42, line 31: 42(37) should be 42(38)
- p. 130, right-hand column, couplet 50, line 12: 50(38) should be 50(37)

2) Minor corrections:

- p. 1, right-hand column, line 15: read <http://www.faunaeur.org>
- p. 15, left-hand column, line 10: read ... and is ... instead of ... which are ...
- p. 17, right-hand column, line 12: read ... and therefore could not be ...
- p. 17, right-hand column, lines 46–47: read ... an exact ... instead of ... axed ...
- p. 19, left-hand column, line 36: read *Cacoxenus perspicax* (Knab, 1914)
- p. 19, right-hand column, line 14: read *Phortica semivirgo* (Máca, 1977)
- p. 27, right-hand column, line 14: read ... weakly sclerotized... instead of ... hardly sclerotised ...
- p. 30, right-hand column, lines 20–21: delete (microtrichose in some Oriental species)
- p. 48, right-hand column, line 13: read (Figs 61, 65, 66, 106–109)
- p. 68, Figs 150 and 151: refer to the Figs 146 and 147 (p. 65) for labelling of the sclerites which are from top to bottom in the same sequence; additionally read 'posterior ejaculatory duct' instead of 'ejaculatory'
- p. 71, left-hand column, line 33: read *P. semivirgo* (Máca)
- p. 71, left-hand column, line 43: read *P. variegata* (Fallén)
- p. 115, right-hand column, line 43: read (Figs 236–238, 262–266)
- p. 146, left-hand column, line 18: read ... high ... instead of ... long ...
- p. 151, left-hand column, line 23: read ... 1996). instead of ... 1992).
- p. 151, left-hand column, line 41: read (Figs 334, 413–417, 463)
- p. 159, left-hand column, line 42: read (Figs 366–369, 426–430)
- p. 162, right-hand column, line 40: read (Figs 370–373, 431, 434–438)
- p. 166, left-hand column, line 16: read ... forwards.
- p. 166, left-hand column, line 17: read Aedeagus has ... instead of and has
- p. 169, left-hand column, line 37: read (Figs 374–377, 433, 444–447)
- p. 178, left-hand column, line 16: read ... medially interrupted dark ... instead of ... dark ...
- p. 178, right-hand column, line 45: read (Figs 457–460)
- p. 182, right-hand column, line 6: read (Figs 341, 342, 465–469)
- p. 185, right-hand column, line 3: read (Figs 349, 350, 462, 470–474)
- p. 188, right-hand column, line 16: read (Figs 345, 346, 475–479)
- p. 191, right-hand column, line 14: read (Figs 347, 348, 480–484)
- p. 195, left-hand column, line 35: (Figs 286, 287, 289, 291, 485–489)
- p. 199, left-hand column, line 7: read (Figs 290, 292, 490, 493–496)
- p. 202, left-hand column, line 30: read (Figs 299, 316, 491, 497–501)
- p. 205, right-hand column, line 15: read (Figs 324, 492, 502–506)

- p. 208, right-hand column, line 39: read (Figs 309, 322, 507–511, 515)
- p. 213, left-hand column, line 16: read (Figs 308, 323, 512, 516–520)
- p. 216, left-hand column, line 41: read (Figs 304, 305, 315, 513, 521–524)
- p. 218, right-hand column, line 47: read (Figs 300, 310, 317, 321, 514, 525–529)
- p. 222, left-hand column, line 22: read (Figs 295, 301, 302, 312, 530–534, 538)
- p. 226, right-hand column, line 6: read (Figs 306, 307, 314, 319, 535, 539–543)
- p. 230, left-hand column, line 5: read (Figs 293, 294, 296, 313, 320, 536, 544–548)
- p. 252, right-hand column, line 37: read (Figs 586–588, 595–599)
- p. 256, left-hand column, line 39: read (Figs 592–594, 600, 604–607)
- p. 259, left-hand column, line 8, read: (Figs 589–591, 601, 608–611)
- p. 261, right-hand column, line 42: read (Figs 583–585, 602, 612–615)
- p. 264, right-hand column, line 10: read (Figs 578, 581, 603, 616–619)
- p. 275, left-hand column, line 10: read (Figs 56, 625, 630, 633–635, 638–641)
- p. 286, left-hand column, line 37: read (Figs 652, 656–658, 686–690)
- p. 289, right-hand column, line 37: read (Figs 651, 653–655, 691, 694–697)
- p. 292, right-hand column, line 43: read (Figs 50, 51, 648–650, 692, 698–701)
- p. 288, Fig. 693: this illustration may refer to a different species (Máca, pers. comm.)
- p. 284, Fig. 673: this illustration may refer to a different species (Máca, pers. comm.)
- p. 298, left-hand column, lines 17–52: this description may refer to a different species (Máca, pers. comm.)
- p. 296, left-hand column, line 12: read (Figs 693, 671, 672, 702–705); see comment above
- p. 298, right-hand column, line 31: read (Figs 662, 663, 666–668, 706–710)
- p. 301, left-hand column, line 39: read (Figs 52, 664, 665, 677–679, 711, 714–717)
- p. 305, left-hand column, line 8: read (Figs 680–682, 712, 718–721)
- p. 307, right-hand column, line 16: read (Figs 669, 670, 722–725)
- p. 310, left-hand column, line 6: read (Figs 674–676, 713, 726–729)

3) For some taxa names mentioned, the respective references are missing. Instead of including them in this publication, we suggest to refer to the online database TAXODROS (Bächli 2007).

ACKNOWLEDGEMENTS

Pierre Bächli helped with preparing the photomicrographs for publication; Elsa Obrecht (Bern), Milan Chvála (Praha) and in particular Jan Máca (České Budějovice) informed us about some errors in Bächli *et al.* (2004) to be corrected.

REFERENCES

- Bächli, G. 2007. TAXODROS, the online database on taxonomy of Drosophilidae. Version June 2007. www.taxodros.uzh.ch
- Bächli, G., Vilela, C.R. & Haring, E. 2002. Four new species of West Palaearctic Drosophilidae (Diptera). *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 75: 299–333.
- Bächli, G., Vilela, C.R., Andersson Escher, S., & Saura, A. 2004. The Drosophilidae (Diptera) of Fennoscandia and Denmark. *Fauna Entomologica Scandinavica*, vol. 39, 362 pp. Brill, Leiden.
- Bächli, G., Haring, E. & Vilela, C.R. 2005. On the phylogenetic relationships of *Scaptodrosophila rufifrons* and *S. lebanonensis* (Diptera, Drosophilidae). *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 78: 349–364.


(received October 31, accepted November 15, 2007)

Table 1. List of the specific epithets of Scandinavian drosophilids, their respective male internal terminalia photomicrographs, and previously published line drawings.


Specific epithet	this publication	Bächli <i>et al.</i> 2004
<i>acuminata</i>	Fig. 70	p. 253 Fig. 597
<i>albilabris</i>	Fig. 4	p. 35 Fig. 85
<i>alboguttata</i>	Fig. 6	p. 37 Fig. 89
<i>alpina</i>	Fig. 56	p. 203 Fig. 499
<i>ambigua</i>	Fig. 57	p. 206 Fig. 504
<i>amoena</i>	Fig. 25	p. 110 Fig. 255
<i>andalusiaca</i>	Fig. 73	p. 257 Fig. 606
<i>argyreator</i>	Fig. 9	p. 56 Fig. 125
<i>baechlii</i>	Fig. 16	p. 85 Fig. 196
<i>basdeni</i>	Fig. 1	p. 46 Fig. 104
<i>bifasciata</i>	Fig. 58	p. 209 Fig. 509
<i>busckii</i>	Fig. 29	p. 136 Fig. 388
<i>cameraria</i>	Fig. 69	p. 243 Fig. 568
<i>caudatula</i>	Fig. 24	p. 103 Fig. 242
<i>coleoptrata</i>	Fig. 17	p. 86 Fig. 200
<i>confusa</i>	Fig. 67	p. 246 Fig. 572
<i>congesta</i>	Fig. 76	p. 268 Fig. 620
<i>constimilis</i>	Fig. 81	p. 297 Fig. 704
<i>costata</i>	Fig. 26	p. 107 Fig. 250
<i>deflexa</i>	Fig. 75	p. 278 Fig. 644
<i>distigma</i>	Fig. 10	p. 61 Fig. 137
<i>distincta</i>	Fig. 27	p. 113 Fig. 259
<i>eskoii</i>	Fig. 59	p. 214 Fig. 518
<i>ezoana</i>	Fig. 46	p. 183 Fig. 467
<i>fenestrarum</i>	Fig. 74	p. 260 Fig. 610
<i>flava</i>	Fig. 82	p. 299 Fig. 708
<i>flavopruinosa</i>	Fig. 3	p. 42 Fig. 96
<i>formosus</i>	Fig. 7	p. 29 Fig. 59
<i>funnebris</i>	Fig. 30	p. 140 Fig. 396
<i>furta</i>	Fig. 15	p. 81 Fig. 190
<i>fuscimana</i>	Fig. 28	p. 116 Fig. 264
<i>graminum</i>	Fig. 83	p. 304 Fig. 716
<i>griseola</i>	Fig. 84	p. 306 Fig. 720
<i>helvetica</i>	Fig. 60	p. 217 Fig. 523
<i>hexasticha</i>	Fig. 71	p. 262 Fig. 614
<i>histrion</i>	Fig. 31	p. 144 Fig. 401
<i>hydei</i>	Fig. 42	p. 173 Fig. 450
<i>hypoleuca</i>	Fig. 18	p. 89 Fig. 204
<i>immigrans</i>	Fig. 32	p. 147 Fig. 406
<i>indagator</i>	Fig. 8	p. 54 Fig. 121
<i>ingrica</i>	Fig. 55	p. 234 Fig. 551
<i>kuntzei</i>	Figs 34, 35	p. 160 Fig. 428
<i>limbata</i>	Figs 36, 37	p. 163 Fig. 436
<i>littoralis</i>	Fig. 47	p. 186 Fig. 472
<i>longifibula</i>	Fig. 19	p. 91 Fig. 209
<i>lummei</i>	Fig. 48	p. 189 Fig. 477
<i>lundstroemi</i>	Fig. 65	p. 237 Fig. 555
<i>maculata</i>	Fig. 13	p. 65 Fig. 146
<i>mehadiae</i>	Fig. 20	p. 93 Fig. 213
<i>melanogaster</i>	Figs 50, 51, 52	p. 196 Figs 487, 488
<i>montana (Drosophila)</i>	Fig. 49	p. 192 Fig. 482
<i>montana (Scaptomyza)</i>	Fig. 85	p. 308 Fig. 724
<i>nigricolor</i>	Fig. 72	p. 265 Fig. 618
<i>nigrithorax</i>	Fig. 21	p. 96 Fig. 218
<i>obscura</i>	Fig. 61	p. 219 Fig. 527
<i>oldenbergi</i>	Fig. 66	p. 241 Fig. 564

PHOTOMICROGRAPHS OF MALE TERMINALIA OF SCANDINAVIAN DROSOPHILIDAE


<i>pallida</i>	Fig.	80	p. 293	Fig.	700
<i>phalerata</i>	Figs	38, 39	p. 167	Fig.	441
<i>picta</i>	Fig.	33	p. 157	Fig.	424
<i>quinquemaculata</i>	Fig.	14	p. 68	Fig.	150
<i>repleta</i>	Figs	43, 44	p. 176	Fig.	455
<i>rufescens</i>	Fig.	5	p. 49	Fig.	108
<i>semivirgo</i>	Fig.	12	p. 73	Fig.	162
<i>similis</i>	Figs	22, 23	p. 98	Fig.	222
<i>simulans</i>	Figs	53, 54	p. 200	Figs	495, 496
<i>subobscura</i>	Fig.	62	p. 223	Fig.	532
<i>subsilvestis</i>	Fig.	63	p. 227	Fig.	541
<i>subtusradiata</i>	Fig.	2	p. 44	Fig.	100
<i>teinoptera</i>	Fig.	86	p. 311	Fig.	728
<i>testacea</i>	Fig.	45	p. 179	Fig.	459
<i>transversa</i>	Figs	40, 41	p. 170	Fig.	446
<i>tristis</i>	Fig.	64	p. 231	Fig.	546
<i>trivittata</i>	Fig.	68	p. 249	Fig.	576
<i>trochanterata</i>	Fig.	78	p. 287	Fig.	688
<i>unipunctum</i>	Fig.	79	p. 290	Fig.	696
<i>variegata</i>	Fig.	11	p. 75	Fig.	166
<i>zetterstedti</i>	Fig.	77	p. 271	Fig.	626




Figs 1–6. Photomicrographs of internal male terminalia of species of *Amiota*, left lateral view. 1: *A. basdeni*, Biel, Switzerland; 2: *A. subtusradiata*, Tvärminne, Finland; 3: *A. flavopruinosa*, Finges, Switzerland; 4: *A. albilabris*, Tjentište, Serbia; 5: *A. rufescens*, Parc National Suisse, Switzerland; 6: *A. alboguttata*, Aucelon, France.




Figs 7–14. Photomicrographs of internal male terminalia of species of Steganinae, left lateral view. 7: *Acletoxenus formosus*, K. Meiron, Israel; 8: *Cadoxenus indagator*, Fazel, Israel; 9: *Cadoxenus argyreator*, Pfywald, Switzerland; 10: *Gitona distigma*, Gyón, Hungary; 11: *Phortica variegata*, Popovica, Serbia; 12: *Phortica semivirgo*, Arcegno, Switzerland; 13: *Leucophenga maculata*, Seelisberg, Switzerland; 14: *Leucophenga quinquemaculata*, Someo, Switzerland.




Figs 15–23. Photomicrographs of internal male terminalia of species of *Stegana*, left lateral view. 15: *S. furta*, Finges, Switzerland; 16: *S. baechlii*, Aarau, Switzerland; 17: *S. coleoprata*, Karislojo, Finland; 18: *S. hypoleuca*, Dyrehaven, Denmark; 19: *S. longifibula*, Biasca, Switzerland; 20: *S. mehadiae*, Beeram, Denmark; 21: *S. nigrithorax*, Dietikon, Switzerland; 22: *S. similis*, Höggerberg, Switzerland; 23: *S. similis*, Höggerberg, Switzerland.




Figs 24–29. Photomicrographs of internal male terminalia of species of Drosophilidae, left lateral view, all collected in Switzerland. 24: *Chymomyza caudatula*, Someo; 25: *Chymomyza amoena*, Dietikon; 26: *Chymomyza costata*, Il Fuorn; 27: *Chymomyza distincta*, Faido; 28: *Chymomyza fuscimana*, Faido; 29: *Drosophila busckii*, Rheineck.




Figs 30–33. Photomicrographs of internal male terminalia of *Drosophila* species, left lateral view. 30: *D. funebris*, Seelisberg, Switzerland; 31: *D. histrio*, Rheineck, Switzerland; 32: *D. immigrans*, Gomel, Belarus; 33: *D. picta*, Bordils, Spain.




Figs 34–45. Photomicrographs of internal male terminalia (only aedeagus+aedeagal apodeme in Figs 35, 37, 39, 41, and 44) of seven species of *Drosophila*, left lateral view. 34: *D. kuntzei*, Würenlingen, Switzerland; 35: idem [Würenlingen, Switzerland, 6–11.6.73, G. Bächli coll.]; 36: *D. limbata*, Stams, Austria; 37: idem [Rheineck, Switzerland, 14–17.8.73, G. Bächli coll.]; 38: *D. phalerata*, Dietikon, Switzerland; 39: idem [Dietikon, Switzerland, Mai 1974, G. Bächli coll.]; 40: *D. transversa*, Delémont, Switzerland; 41: idem [Rheineck, Switzerland, 14–17.8.73, G. Bächli coll.]; 42: *D. hydei*, Alexandria, Egypt; 43: *D. repleta*, Bachs, Switzerland; 44: idem [Cueva Chirrepeck, Alta Verapaz, Guatemala, 6–8.4.1973, Strinetti coll.]; 45: *D. testacea*, Dietikon, Switzerland.




Figs 46–55. Photomicrographs of internal male terminalia of *Drosophila* species, mainly of the *virilis* and *melanogaster* groups, left lateral view (except Figs 51 and 54: posterior view). 46: *D. ezoana*, Kemi, Finland; 47: *D. littoralis*, Popovica, Serbia; 48: *D. lummei*, Hokkaido, Japan; 49: *D. montana*, Kemi, Finland; 50: *D. melanogaster*, Hönngerberg, Switzerland; 51: same specimen as 50; 52: *idem*, unvert aedeagus [Landquart, Switzerland, 9–12.8.74, G. Bächli coll.]; 53: *D. simulans*, Rheineck, Switzerland; 54: same specimen as 53; 55: *D. ingrlica*, Suistamo, Russia.




Figs 56–64. Photomicrographs of internal male terminalia of *Drosophila* species of the *obscura* group, left lateral view, all from Switzerland (except *D. eskoi*). 56: *D. alpina*, Les Reussilles; 57: *D. ambigua*, Geeren; 58: *D. bifasciata*, Arcegno; 59: *D. eskoi*, Oulu, Finland; 60: *D. helvetica*, Dietikon; 61: *D. obscura*, Würenlingen; 62: *D. subobscura*, Würenlingen; 63: *D. subsilvestris*, Niesen; 64: *D. tristis*, Dietikon.



Figs 65–72. Photomicrographs of internal male terminalia of species of the genera *Hirtodrosophila* and *Lordiphosa*, left lateral view. 65: *Hirtodrosophila lundstroemi*, Gödöllő, Hungary; 66: *Hirtodrosophila oldenbergi*, Hönggerberg, Switzerland; 67: *Hirtodrosophila confusa*, Bergdietikon, Switzerland; 68: *Hirtodrosophila trivittata*, Hluboka, Czech Republic; 69: *Hirtodrosophila cameraria*, Seelisberg, Switzerland; 70: *Lordiphosa acuminata*, Pichelsberg, Germany; 71: *Lordiphosa hexasticha*, Rappersdorf, Germany; 72: *Lordiphosa nigricolor*, Albisgütli, Switzerland.



Figs 73–77. Photomicrographs of internal male terminalia (plus external terminalia in Figs. 76 and 77) of species of Drosophilidae, left lateral view. 73: *Lordiphosa andalusiaca*, Vargös, Turkey; 74: *Lordiphosa fenestrarum*, Stams, Austria; 75: *Scaptodrosophila deflexa*, Veyrier, Switzerland; 76: *Microdrosophila congesta*, Aarau, Switzerland; 77: *Microdrosophila zetterstedti*, Smolandia, Sweden [lectotype].



Figs 78–86. Photomicrographs of internal male terminalia of species of *Scaptomyza*, left lateral view. 78: *S. trochanterata*, Urjala, Finland; 79: *S. unipunctum*, Paltamo, Finland; 80: *S. pallida*, Glockner, Austria; 81: *S. consimilis*, Joutseno, Finland; 82: *S. flava*, Bielefeld, Germany; 83: *S. graminum*, Klöntal, Switzerland; 84: *S. griseola*, Eičiai Taurage distr., Lithuania; 85: *S. montana*, Merkinė, Lithuania; 86: *S. teinoptera*, Messuby, Finland.