

**Zeitschrift:** Mitteilungen der Schweizerischen Entomologischen Gesellschaft =  
Bulletin de la Société Entomologique Suisse = Journal of the Swiss  
Entomological Society

**Band:** 84 (2011)

**Heft:** 3-4

**Artikel:** Atrichops crassipes (Meigen, 1820) (Diptera, Athericidae), a species not  
so new for Switzerland

**Autor:** Sartori, Michel / Lods-Crozet, Brigitte / Derleth Sartori, Pascale

**DOI:** <https://doi.org/10.5169/seals-403029>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

**Download PDF:** 15.10.2024

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

*Atrichops crassipes* (Meigen, 1820) (Diptera, Athericidae), a species not so new for Switzerland

MICHEL SARTORI<sup>1</sup>, BRIGITTE LODS-CROZET<sup>1,2</sup>, PASCALE DERLETH SARTORI<sup>1,2</sup>, JEAN PERFETTA<sup>3</sup>, PASCAL STUCKI<sup>4</sup> & HEINRICH VICENTINI<sup>5</sup>

<sup>1</sup> Musée cantonal de zoologie, Palais de Rumine, Place Riponne 6, CH-1005 Lausanne; michel.sartori@vd.ch

<sup>2</sup> Laboratoire du Service des Eaux, Sols et Assainissement (SESA), Ch. des Boveresses 155, CH-1066 Epalinges

<sup>3</sup> Service de l'écologie de l'eau (SECOE-DIM), case postale 78, CH-1211 Genève 8

<sup>4</sup> Aquabug, case postale 1643, CH-2001 Neuchâtel

<sup>5</sup> Gewässerökologie, Wasserschöpfli 48, CH-8055 Zürich

The genus *Atrichops* Verrall, 1909, with the species *A. crassipes* (Meigen, 1820) is documented for the first time in Switzerland, although its first mention dates back to 1995, but remained unnoticed.

Keywords: new record, Switzerland, aquatic insect, Diptera.

#### INTRODUCTION

Athericidae (watersnipe-flies) is a small family of primitive Brachycera related to Tabanidae and Rhagionidae (Stuckenberg 1973; Rozkosny & Nagatomi 1997). The larvae are always aquatic and live in fast flowing streams or in more lentic habitats (Thomas 1981, 1997). They are top predators and feed on other invertebrates such as Diptera (Chironomidae), Ephemeroptera or Plecoptera (Thomas 1975).

Up to now, only two species, out of the 10 known to occur in Europe, have been reported from Switzerland (Merz 1998): *Atherix ibis* (Fabricius, 1798) and *Ibisia marginata* (Fabricius, 1781) which are both relatively common in stony streams. A third one, *Atrichops crassipes* (Meigen, 1820) has been mentioned by Merz (1998) as a potential candidate to live in Switzerland. In fact, we discovered during the writing of this note, that it has been already cited by Oertli (1995) in a species list of macroinvertebrates living in a pond near Geneva. This citation remained unnoticed up to now, although published in an entomological journal. Here we document further on the presence of this species in several Swiss streams and rivers, its most common habitat.

#### *Atrichops crassipes* (Meigen, 1820)

#### Material:

**Geneva:** **Seymaz**, embouchure, 502970/115070, 392 m, 1 larva; Claparede, 503220/116000, 403 m, 3 larvae; De Haller, 504095/117390, 417 m, 2 larvae, II. 2007. **Nant du Paradis**, embouchure, 507215/12065, 428 m, 3 larvae, II. 2007. **Hermance**, Pont de Bouringe, 508225/128395, 380 m, 1 larva, XI. 2007; Pont de

Crevy, 507695/126490, 415 m, 2 larvae, V. 2007. Marnot, embouchure, 508600/127400, 395 m, 1 larva, V. 2007. **Rhone**, Touvière, 488100/114740, 347 m, 1 larva, XII. 2008. **Arve**, Ecole-de-Médecine, 499440/116815, 375 m, 1 larva, V. 2008. All coll. SECOE.

*Vaud*: **Venoge**, Bussigny/Lausanne, 531710/ 155175, 375 m, 3 larvae; Vufflens-la-Ville, 530415/ 158980, 405 m, 2 larvae; Denges, 531505/152915, 380 m, 3 larvae, all coll. 16.II.2010 SESA. Denges, 1 larva, 16.IV.2011, coll. P. Derleth.

*Jura*: **Vendline**, Beurnevésin aval, frontière, 576345/261000, 419 m, 2 larvae, 31.III.2011, coll. P. Stucki.

*Lucerne*: **Suhre**, Oberkirch, 663459/225401, 506 m, 1 larva, 25. IV. 2007, 3 larvae, 28. VIII. 2007, 2 larvae, 16. IV. 2003, 3 larvae, 14. VII. 2003; Büron, 648740/228840, 487 m, 1 larva, 15. IV. 2003, coll. H. Vicentini. **Aabach**, Hitzkirch, 661390/230220, 473 m, 2 larvae, 15. IV. 2003, coll. V. Lubini.

*A. crassipes* is known from Western Europe, extending its range up to Poland and Romania (Roskosny 2011). It has been recently mentioned from Hungary for the first time (Muranyi *et al.* 2009). The larvae are psammophilous, living in sand and fine substrate deposited by the current in streams and rivers, where they feed largely on Chironomidae (Thomas 1975). They also colonize the deposits of some dam-lakes (Gagneur 1981) or ponds (Oertli 1995). The morphology of the larvae is very characteristic and cannot be confused with any other species, even among watersnipe-flies. Abdominal segments 6 and 7 bear each four tracheal projections, (two dorsal and two lateral), and the anal segment two more. They are oriented rearwards and upwards to the surface of sediments (Fig. 1).

This merolimnic species goes through five aquatic larval instars before pupation in the banks, and the single available study reports a semivoltine cycle, i.e. development lasts two years (Gerke & Böttger 2001). At the moment, no adults have been caught but according to Thomas (1981) adults fly between June and July. As for other Athericidae, females lay their eggs in batches on the underside of leaves above the watercourse (Gerke & Böttger 2001), and first instar larvae drop into the water shortly after hatching.

In the Canton de Vaud, no specimens have been caught before 2010. Up to now, the Venoge River is the only watercourse to be colonized by *A. crassipes*. At the mentioned localities, the river is between 8 and 10 m wide, the current speed is around 0.2 and 0.4 m/s. Substrate composition and bank vegetation strongly vary between localities, suggesting that the availability of adequate fine substrate is the main parameter for its setting up.

It has been reported that *A. crassipes* belongs to species on which global climate change has a positive impact (Daufresne *et al.* 2003). This could explain the actual scarcity of data for this species in Switzerland, its low altitudinal range (all localities below 510 m), and the fact that it could be a new immigrant, as documented for instance for the mayfly *Habrophlebia eldae* (Wagner *et al.* 2007).

#### ACKNOWLEDGEMENTS

Our sincere thanks to Alain Thomas (Toulouse) for his comments and to Verena Lubini (Zurich) for sharing her data.






Fig. 1 : *Atrichops crassipes* (Meigen, 1820), larva. A: general habitus, B: Larva digging in the sediment, C: close view of the mouthparts, especially the mandibles.

## REFERENCES

- Daufresne, M., Roger, M.C., Capra, H. & Lamouroux, N. 2003. Long-term changes within the invertebrate and fish communities of the Upper Rhône River: effects of climatic factors. — *Global Change Biology* 10: 124-140.
- Gagneur, J. 1981. Sur la biologie d'*Atrichops crassipes* Meigen (Diptères, Athericidae) dans le réservoir de Cajarc (Lot). — *Bulletin de la Société d'Histoire naturelle de Toulouse* 117: 231-237.
- Gerke, N. & Böttger, K. 2001. The life cycle of *Atrichops crassipes* Meigen, 1820 (Diptera: Athericidae) at the lower Schierensee brook, a lake outflow in the North German Lowland. — *Aquatic Insects* 23: 85-92.
- Merz, B. 1998. Athericidae. *In*: Merz, B., Bächli, G., Haenni, J.-P. & Gonseth, Y. (eds), *Diptera – Checklist*, p. 149. — *Fauna Helvetica* 1, CSCF-SEG, Neuchâtel.
- Muranyi, D., Tarjanyi, N. & Schöll, K. 2009. First record of the genus *Atrichops* Verrall, 1909 in Hungary (Diptera: Athericidae). — *Opuscula Zoologica, Budapest* 40: 103-105.
- Oertli, B. 1995. Impact de la création d'un biotope sur la conservation de la biodiversité : les macroinvertébrés de l'étang du Bois-Vieux (Genève). — *Bulletin romand d'Entomologie* 13: 11-24.
- Rozkosny, R. 2011. Fauna Europaea: Athericidae. *In*: Papp, T. & Beuk, P. (eds), *Fauna Europaea: Diptera, Brachycera. Fauna Europaea version 2.4*, <http://www.faunaeur.org>
- Rozkosny, R. & Nagatomi, A. 1997. Family Athericidae. *In*: Papp, L. & Darvas, B. (eds), *Contributions to a manual of Palearctic Diptera*, volume 2, pp. 439-446. — *Science Herald, Budapest*.
- Stuckenberg, B.R. 1973. The Athericidae, a new family in the lower Brachycera (Diptera). — *Annals of the Natal Museum* 21: 649-673.
- Thomas, A.G.B. 1975. Diptères torrenticoles peu connus: III.- Les Athericidae du sud de la France (Régime alimentaire des larves: aspect qualitatif) (Brachycera, Orthorrhapha). — *Annales de Limnologie* 11: 169-188.
- Thomas, A.G.B. 1981. Travaux sur la taxonomie, la biologie et l'écologie d'insectes torrenticoles du Sud-ouest de la France (Ephéméroptères et Diptères: Dixidae, Cecidomyiidae, Rhagionidae et Athericidae), avec quelques exemples de perturbations par l'homme. — Thèse de doctorat, n° 988, Université Paul Sabatier, Toulouse, 330 pp.
- Thomas, A.G.B. 1997. Rhagionidae and Athericidae, Snipe-flies. *In*: Nilsson, A.N. (ed.), *Aquatic Insects of North Europe*, volume 2, pp. 311-320. — *Apollo Books, Stenstrup*.
- Wagner, A., Lubini, V. & Vicentini, H. 2007. *Habrophlebia eldae* Jacob & Sartori, 1984 (Ephemeroptera: Leptophlebiidae) neu für die Schweizer Fauna. — *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 80: 91-98.

(received September 15, 2011, accepted November 7, 2011)