Zeitschrift:	Schweizerische mineralogische und petrographische Mitteilungen = Bulletin suisse de minéralogie et pétrographie
Band:	50 (1970)
Heft:	2
Artikel:	Geological observations in the Bergell Area (SE-Alps). II, Contact minerals from Mt. Sissone-cima di Vazzeda (a mineralogical note)
Autor:	Wenk, Hans-Rudolf / Maurizio, Remo
DOI:	https://doi.org/10.5169/seals-39261

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. <u>Siehe Rechtliche Hinweise.</u>

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. <u>Voir Informations légales.</u>

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. <u>See Legal notice.</u>

Download PDF: 29.04.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Geological Observations in the Bergell Area (SE-Alps) II. Contact Minerals from Mt. Sissone-Cima di Vazzeda (A Mineralogical Note)

By Hans-Rudolf Wenk (Berkeley)*) and Remo Maurizio (Vicosoprano)

With 2 figures and 3 tables in the text

Abstract. Chemical data and unit cell dimensions are given for some newly found minerals occurring in the contact zone of carbonate and calc-silicate rocks and amphibolites with the Bergell granite in the Sissone-Vazzeda area. Diopsides show a linear relation between lattice constants and Mg/Fe ratio. Of special interest are a deep blue diopside and scheelite.

The northern contact zone of the granitic rocks of the Bergell with metamorphic rocks of the Margna nappe has been studied geologically by GYR (1967). Discordant contacts of calc-silicates and amphibolites with granite and well preserved lenses of marble which are floating in the granite are the most conspicuous features. These rocks belong to the amphibolite metamorphic facies and show mineral assemblages labradorite-hornblende (E. WENK and KELLER, 1969), and anorthite-diopside-calcite with wollastonite and clinohumite (E. WENK, 1962; E. WENK, 1963; TROMMSDORFF, 1966; SCHWANDER and E. WENK, 1967; E. WENK et al., 1967). The Bergell is known for its abundance of rare minerals, some of them unique for Switzerland. It is surprising that they have been treated only superficially. In addition to the papers mentioned above STAUB (1920), DÉVERIN (1937), NIGGLI et al. (1940), WEIBEL and LOCHER (1964) are the only references we were able to find in the literature on the mineralogy of the area. In recent years (1966–1969) some unusual finds have been made in the area Val Sissone-Cima di Vazzeda-Vadrec del Forno which warrant a description in a short note. Localities are listed in Table 1. Most of the specimens were found in moraines. The original samples are in the mineralogical collection of the museum in Stampa (Ciäsa Granda). The results reported in

^{*)} University of California, Dept. of Geology and Geophysics, Berkeley, California 94720.

Diopside 1	Val Sissone, 2000 m	777.9/129.5
2	Alpe Sissone, 2600 m	777.1/130.9
3	Val Sissone, 2200 m	777.4/129.4
4	Val Sissone, 2200 m	777.4/129.4
5 6	Alpe Vazzeda, 2600 m moraine from Pizzo Vazzeda,	777.2/131.8
	Vadrec del Forno, 2600 m	774.7/132.4
Tremolite	same specimen as diopside 4	
Spinel	moraine from Pizzo Vazzeda, Vadrec del Forno, 2600 m	774.7/132.4
Scheelite	Val Sissone, 2500 m	777.0/130.0

Table 1. Localities (coordinates of Swiss maps are used)All specimens were found in blocks in local moraines

this study include microprobe analyses (done by F. Brown at Berkeley: Table 2), and precise determination of lattice constants with a Jagodzinskitype Guinier camera (Fe K α_1 -radiation, Si as internal standard, least squares refinement: Table 3).

This short mineralogical note is meant as part of a series of contributions of data to the geology of the Bergell. No effort for completeness is made, nor do we try to propose ultimate interpretations. (See also WENK, 1970a.)

Diopside

TROMMSDORFF (1966) shows a map of the distribution of diopside associated with calcite in the Lepontine Alps. Its occurrence coincides with the zone of highest regional metamorphism. The Bergell is located in the peripheral part of this metamorphic complex. We describe six finds of diopside of various chemical compositions in this area.

1. Dark green prismatic crystals, up to two inches long. Good cleavage. Inclusions in pegmatitic dykes cutting across amphibolites.

2. Tabular idiomorphic crystals in a cavity associated with quartz, feldspar and calcite. Light green color. Many crystals clear. 1/8-11/2''. These crystals were used by LEUNG (1969) to study deformation effects. Under nonhydrostatic stress submicroscopic twinning on (100) was produced (cf. WENK, 1970b).

3. Aggregate of prismatic greyish-green milky crystals in a cavity. Individual crystals reach a size of $1^{1}/_{2}$ ". They are chemically very heterogeneous. The diopside is associated with calcite.

4. Fine-grained turquoise blue diopside intermixed with coarser-grained colorless diopside occurs as the central part of a concentrically zoned lens 2' long and 3'' wide (Fig. 1). The blue diopside is surrounded by 1'' of white pure tremolite (cf. WENK, 1970c) and 1-2'' dark green diopside, rich in Fe. The zoned lens is included in white calcite marble. Due to its conspicuous color the diopside has first been mistaken for lazulite which occurs in the lower meta-morphic rocks of the Suretta nappe along the north side of the Bergell Valley.



Fig. 1. Schematic sketch of the lens of blue diopside in Val Sissone.



Fig. 2. Relation between lattice constants and Fe/Mg ratio. The fine lines are the ones given by RUTSTEIN and YUND (1969); the dashed lines indicate the dependence found in this study.

Microprobe analyses showed that the coexisting colorless and blue diopside are very similar in chemical composition, almost Fe and Al free. However, there is a significant difference in the vanadium content. The electron beam of the microprobe produces bright blue fluorescence in the blue diopside and a pale pink one in the colorless-type. It is not impossible that this minute chemical difference be due to metasomatic replacement of fossiliferous limestone, as trace elements are often enriched in organic material.

5. In the moraine east of Passo di Vazzeda which is rich in calc-silicate rocks, large white crystals up to 1' long are common. The mineral which could be mistaken for wollastonite with which it is associated, has been identified by GYR (1967) as diopside. It is chemically extremely pure.

6. Similar crystals as are found on the Italian part of P. Vazzeda are carried down to the Forno glacier: they are very common in the northeastern moraine near Cap. Forno which is characterized by colorful carbonate and calc-silicate rocks originating from P. Vazzeda.

The chemical composition of diopsides from the same region varies considerably. A plot of chemical composition (Mg/Fe ratio) versus lattice constants of these genetically related specimens shows a nearly linear correlation which resembles but nevertheless differs from the results of RUTSTEIN and YUND (1969) for synthetic diopsides. It is uncertain whether these differences are caused by conditions during formation or by a more complicated chemistry in the natural crystals (e. g., Al). It indicates, however, that the curves should be used with caution. It is obviously unjustified to generalize results from these few observations, but it appears that:

- -- the diopsides from carbonate rocks are extremely pure and homogeneous;
- --- diopsides in cavities (Alpine-type fissures) have an average of 15% Mg (by weight) and show even in clear single crystals large variations in chemical composition;
- diopsides in amphibolites have a composition of close to 50% hedenbergite (see also WENK, 1970b).

Spinel

In marbles and calc-silicate rocks of the Vazzeda region very pure Al-Mg-Zn spinels occur quite frequently. Octahedral crystals which are associated with anorthite, calcite and grossularite are black or dark green and approximately 1/8'' in size. The marble is fairly coarse-grained.

Scheelite

Scheelite is rare in the Alps. NIGGLI et al. (1940) describe finds of scheelite in the Aarmassiv and in the Simplon railway tunnel. The mineral was so far unknown in the Bergell. In 1967 a specimen of yellow scheelite with idio-

Table 2. (Jhemical datı	ı (detern	ined wit	sh an AF	L microl	probe by F	. Brown).	Values are	in weight _I	bercent		
	TiO ₂	Si02	V	l ₂ O ₃	FeO	Ca(MgO	OuM	V_2O_3	ZnO	
Diopside			6						t.			
1 Sissone	0.02	52.6	_	0.8	11.4	24.	2	10.8	0.5			_
2 Vazzeda	0	55.0		0.5	2.9	25.	++	16.3	0.4			_
3 Sissone	0.1	54.4		0.73	3.46	25.	4	15.9	0.2			1.0
4a Sissone blue ¹)	0.03	55.2		0.1	0.08	25.	7	18.8	0	0.11		-
4 b Sissone colorless ²)	0	55.3	_	0.1	0.1	25.	6	18.6	< 0.01	0.03		
5 Vazzeda-Sissone	0.11	55.4		0.83	0.14	25.		18.0	0.03			
3	(0.03-0.12)		(0.4)	9-1.3)	(2.9-4.0) (25.2-5	25.5) (15	.2-16.2) (0.19-0.27)	3		
Tremolite	0.01	56.8		0.1	0.24	13.	85	25.5	0			
Spinel	0.02	0.0	0	9.3	2.97	0.	01	24.7			3)	
1) br	ight blue fluo	rescence	-	²) pale pu	urple fluor	escence	³) cor	responds to	$3.29\% {\rm Fe_2O}$	3		
Tab	ole 3. Lattice	constants	s determi	ned with	a Jagodz	inski-type (Twinier can	nera (error	is $\pm 0.01\%$	(-55
	X-rav Num	ber of	D I	irect latt	ice constar	nts	Cell	Re	ciprocal latt	ice constan	ts	
	Nr. observ	ations	a_0 (Å)	$b_0(Å)$	$\mathbf{c}_0\left(\mathrm{\AA}\right)$	β	$\nabla (\Lambda^3)$	a* (A ⁻¹)	b* (A ⁻¹)	c* (A ⁻¹)	β*	
Diopside 1a	369 2	0	9.775	8.951	5.257	105.66°	443.0	0.10624	0.11172	0.19754	74.34°	
1 1b	379 1	6	9.777	8.954	5.258	105.66°	443.2	0.10622	0.11169	0.19750	74.34°	
la+b	ŝ	6	9.776	8.952	5.258	105.66°	443.1	0.10623	0.11171	0.19752	74.34°	
2	378 1	9	9.755	8.930	5.253	105.79°	440.4	0.10653	0.11198	0.19785	74.21	
33	474 1	<i>∞</i>	9.754	8.931	5.253	105.78°	440.4	0.10653	0.11197	0.19782	14.22	
4a	$\frac{316}{21}$		9.750	8.927	5.251		439.7	0.10660	0.11202	0.19793	74.160	
4 b	375 2	2	9.751	8.920	0.202	100.84	439.1	10001.0	0.11204	06/61.0	01.17	
4a+b	- - -	_ ,	9.750_{3}	8.926_{0}	5.251_4	105 833	439.70	0.10660	$5 0.11203_2$	0.19793	74.107	
0.0	476	- 1	9.752	8.926	0.202	105.54	439.8	0.10650	0111005	010702	14.180	
9	475 I	 ດ	9.70T	8.924	0.201	109.62	439.1	00001.0	0.11200	celet.0	01.11	
I remoute with diopside 4	325 2		9.844	18.053	5.279	104.75°	907.2	0.10505	0.05539	0.19489	75.25°	
Scheelite	346 1	3	5.2450		11.376		312.95	0.19066		0.08791		
Spinel	322 2	0	8.089				529.3			0.12360		
	(2 f	ilm)										

.

Geological Observations in the Bergell Area (SE-Alps)

353

H.-R. Wenk and R. Maurizio

morphic pyramidal forms, 1/2'' in size was found in a quartz vein in Val Sissone. It seems to be a product of hydrothermal activity possibly related to the formation of molybdenite in the Forno granite. This hydrothermal activity produced a system of quartz, aplite and pegmatite dykes which penetrate the granite and caused alteration in metamorphic rocks in the vicinity (WENK, 1970a).

The authors wish to thank Frank Brown for the chemical analyses, and Ugo Ruinelli and Barbara Hiss for the contribution of specimens from their personal collections.

References

SMPM = Schweiz. Mineralog. u. Petrogr. Mitteilungen

- DEER, W. A., R. A. HOWIE and J. ZUSSMANN (1963): Rock-forming Minerals, vol. 2. Longmans, London.
- DÉVERIN, L. (1937): Kalksilikatmineralien, Bergell. SMPM 17, 531.
- GYR, T. (1967): Geologische und petrographische Untersuchungen am Ostrande des Bergeller Massivs. Diss. ETH, Zürich.
- LEUNG, S. (1969): Deformation Structures of Diopside. Ph. D. thesis, Berkeley.
- NIGGLI, P., J. KOENIGSBERGER and R. L. PARKER (1940): Die Mineralien der Schweizer Alpen, Vol. 1. Wepf, Basel.
- RUTSTEIN, M. S. and R. A. YUND (1969): Unit-cell parameters of synthetic diopsidehedenbergite solid solutions. Amer. Mineral. 54, 238-245.
- SCHWANDER, H. and E. WENK (1967): Studie mit der Röntgen-Mikrosonde an basischen Plagioklasen alpiner Metamorphite. SMPM 47, 225–234.
- STAUB, R. (1920): Gesteinsmetamorphose in Graubünden. Vjschr. natf. Ges. Zürich 65, 1–54.
- TROMMSDORFF, V. (1966): Progressive Metamorphose kieseliger Karbonatgesteine in den Zentralalpen zwischen Bernina und Simplon. SMPM 46, 431–460.
- WEIBEL, M. and T. LOCHER (1964): Die Kontaktgesteine im Albigna- und Fornostollen. (Nördl. Bergeller Massiv.) SMPM 44, 157–185.
- WENK, E. (1962): Plagioklas als Index-Mineral in den Zentralalpen. SMPM 42, 139-152.
- --- (1963): Klinohumit und Chondrodit in Marmoren der Tessiner Alpen und der Disgrazia-Gruppe. SMPM 43, 287–294.
- WENK, E. and F. KELLER (1969): Isograde in Amphibolitserien der Zentralalpen. SMPM 49, 157–177.
- WENK, E., H. SCHWANDER and V. TROMMSDORFF (1967): Optische Orientierung zweier Anorthite aus metamorphen Gesteinen. SMPM 47, 219-224.
- WENK, H.-R. (1970a): Geologische Beobachtungen im Bergell. I. Gedanken zur Genese des Bergeller Granits. Rückblick und Ausblick. SMPM 50, 321–348
- (1970b): Lattice parameters of calcic amphiboles (in preparation).
- (1970c): Submicroscopic twinning in lunar and experimentally deformed pyroxenes. Contrib. Mineral. Petrol.

Manuscript received April 15, 1970.

354