Zeitschrift:	Schweizerische mineralogische und petrographische Mitteilungen = Bulletin suisse de minéralogie et pétrographie
Band:	66 (1986)
Heft:	3
Artikel:	The metamorphic evolution of garnet-cordierite-silimanite gneisses of NW Spitsbergen (Svalbard)
Autor:	Klaper, Eva M.
DOI:	https://doi.org/10.5169/seals-50897

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

Download PDF: 03.07.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Schweiz. mineral. petrogr. Mitt. 66, 295-313, 1986

The metamorphic evolution of garnet-cordierite-sillimanite gneisses of NW Spitsbergen (Svalbard)

by Eva M. Klaper¹

Abstract

Metapelitic gneisses occurring within the Caledonian migmatite complex of NW Spitsbergen (Svalbard archipelago) are most typically represented by schistose biotite-bearing cordierite—garnet—sillimanite rocks.

A consistent pressure-temperature range of 5 to 6 kbar and 650 °C increasing to 700 °C from the south to the north has been deduced for the last metamorphic equilibration using standard geo-thermobarometry with water-saturated cordierite. Calculated with dry cordierite the pressure reaches 4 to 5 kb at the same temperature of metamorphism.

The textural and chemical features are interpreted in terms of a set of (retrograde) mineral reactions. From the textural evidence as well as from the deduced mineral reactions it may be concluded that the rocks were at higher metamorphic conditions prior to the PT-conditions calculated from five published geothermobarometers.

Although the maximum PT-conditions reached are unknown, it is suggested that they were close to granulite facies conditions with low water activity. The H_2O required to form the late stage assemblages containing hydrated minerals such as biotite and muscovite was presumably released by the crystallizing migmatitic melts in the quartzo-feldspathic portions of the rocks. Therefore, the water activity increased significantly during the initial decompression and cooling of the rocks.

The well preserved portion of the PT-path described in this study is interpreted as a part of essentially one main Caledonian metamorphic cycle culminating in regional upper amphibolite facies metamorphism, "in situ" partial melting and migmatization. Additional heating may be postulated in connection with the intrusion of syntectonic gray granites in the northern part of the study area. Weak late heating is related to the intrusion of the post-tectonic Hornemantoppen Batholith.

Keywords: Metapelites, migmatites, geothermobarometry, Caledonian, Spitsbergen.

Introduction

During the summer of 1983 a Swiss expedition visited NW Spitsbergen, Svalbard. As a member of this expedition I was given the opportunity to carry out fieldwork.

Present address: Geologisches Institut, Universität Bern, Baltzerstr. 1, CH-3012 Bern, Switzerland.

¹ Geologisches Institut, ETH-Zentrum, CH-8092 Zurich, Switzerland.

The studied area is located north of 79°30' north and west of 12° east and includes Magdalenefjorden, Smeerenburgfjorden and Vasahalvoya. The region visited consists of a Caledonian migmatite and gneiss complex. Granites, migmatitic and metapelitic gneisses form the vast majority of rock types exposed in this area which had been systematically studied by a number of expeditions of the Polar Institute of Norway. A compilation of most available data has been given by HJELLE and OHTA (1974).

The primary purpose of this study is to describe the observed textures and the mineral chemistry of metapelitic cordierite – garnet – sillimanite gneisses occurring in the lower Hekla Hoek succession of northwestern Spitsbergen. The metamorphic conditions under which these rocks equilibrated will be deduced from phase petrology and standard geothermobarometry.

Geological setting

The Caledonian gneiss and migmatite area of northwestern Spitsbergen is characterized by the predominance of various banded gneisses and granitic migmatites. Amphibolites, hornblende gneisses as well as calcsilicate marbles



Fig. 1 Map of sample localities. Locality numbers correspond to those in Table 1. Geology: migmatitic gneisses and granites (white) and post-tectonic hornblende-biotite granodiorite intrusion (dashed).

and dolomitic marbles occur as small lenses and bands within the migmatite complex. This metasedimentary rock sequence represents part of the lower Hekla Hoek succession of Precambrian age (OHTA 1974a). A large post-tectonic hornblende-biotite granodiorite pluton (Hornemantoppen Batholith) intruded the central part of this area (Fig. 1).

A complex deformation history with three main phases of folding has been described by HJELLE and OHTA (1974). The general trend of early formed fold axes is from north to south, while later minor folds show east-west trending fold axes. Metamorphism reached upper amphibolite facies during the main phases of Caledonian deformation and was followed by recrystallisation which may be correlated with later deformation phases (OHTA 1974b). Formation of an open, regional N-S trending anticlinorium with south plunging fold axis (HJELLE 1979) in NW Spitsbergen is associated with later migmatization and synchronous formation of grey granites. Late stage chloritization can be related to the intrusion of the Hornemantoppen granodiorite body.

Description of rock types and mineral textures

The most common rock types considered in this study are dark biotite-rich metapelitic schists or gneisses and lighter, also biotite-bearing gneissose rocks with cordierite recognizable in hand specimen. The biotite-rich samples are strongly foliated and banded while the lighter coloured rocks usually display only a weakly to moderately developed foliation.

The mineral assemblages found in metapelitic gneisses are listed in table 1. Garnet: In most samples garnet displays irregular and lobate outlines (Fig. 2c). The grainsize typically reaches 1-3 mm. Idiomorphic crystals are extremely rare. Cordierite may replace garnet pseudomorphically. Observed inclusion minerals in garnet are quartz, biotite, plagioclase and, in one sample (Spi 66), also cordierite. Some samples contain web-like garnets which reach diameters of 2-4 cm. Polyphase garnets are rare, but do exist in the northern and western, tectonically deeper parts of the studied area (Fig. 2e, 2f). In some samples garnets are broken apart with muscovite growing in the gaps (Fig. 2b).

Cordierite: Cordierite is very abundant in some rocks. It either forms the rock matrix together with quartz and plagioclase (Fig. 2b, 2d, 2f) or occurs as a late phase replacing garnet (Fig. 2c). Cordierite may be overgrown by fibrolitic sillimanite and is often cracked and pinitized or almost completely transformed to a sericitic felt. Cordierite also forms reaction rims around spinel (Fig. 2a).

Sillimanite: Sillimanite occurs in two different textural types a) as small prismatic needles closely associated with biotite, b) overgrowing cordierite (Fig. 2c) which may replace garnet. Sometimes, sillimanite occurs as radial fibers on

Samp SPI SPI SPI SPI	le 1	Locality	Qtz	Mu	Bi	Gt	5111	Crd	Pla	So	other
SPI SPI SPI SPI	1	standwyddyn ins pedini					3111	er a		90	0 en er
SPI SPI SPI		2	×	16.5 d	×				o		Amph, Chl
SPI SPI	2	2	×	х	х	х			×		Chl,Serizite
SPI	4	2	×	×	×				×		Amph, Chl
COT	5*	2	×	0	×	×			×		
SPI	6	2	×	0	×	×			×		Chl,Ep
SPI	7	9	×	0	×		×	×	×		Chl
SPI	8	9	x	0	· X		×	×	×		Kfs,Chl
SPI	9	9	×	¢	×	×	×		×		Chl
SPI	12*	9	×	×	×	×	×	×	×		Kfs,Chl
SPI	14	10	×		×			×	×		Chl
SPI	15*	8	x		×	×			×		Chl,Amph
Spi	16	8	×	×	×		×	×	×		Kfs
SPI	18*	1	×		×	×			×		Chl
SPI	20	11	×		×				×		Kfs
SPI	21	14	×	0	×	×			×		Kfs
SPI	25	14	×		х	×			×		Chl
SPI	27	14	×	0	×				×		Kfs,Chl
SPI	29	14	×	0	×				×		Kfs
SPI	30	14	×	×	O				×		Kfs,Chl
SPI	33*	14	х		×	×			×		Kfs
SPI	34	14	×		×	×			×		Kfs
SPI	36*	19	×	0	×	×		×	0		Chl
SPI	37*	12	×	0	×	×2	×	×			Chl
SPI	38*	16	×		×	×	×	×	×	0	Kfs,Chl,Ky
SPI	39*	16	×	o	×	0	0	×	o		Chi
SPI	40	19	×	×	0	×		×	×		
SPI	43*	15	×		×	×			×		
SPI	47	17	×		×			×	×		Kfs,Chl
SPI	49	18	×	×	×		×	×	×		Kfs,Chl
SPI	54	3	×		×	×			×		Chi
SPI	55	3	×		×				×		Chi
SPI	59*	4	×		×	×2			×		Kts,Chl
SPI	60	4	×	×	×		×				Chi
SPI	64	7	×	×	×	×	×		×		Chi
SPI	65	7	×		×	×	×		×		Chi
SPI	66*	7	×	×	×	×	×	×	×		Krs, Chi
Oht	403		х	×	×	×			×	×	LO .
Oht	403				×	×		×	212	×	co, Ampri
Oht	440		×		×			×	×	×	0

Tab. 1 Mineral assemblages of metapelitic gneisses.

Oht = Ohta (1974a), x: essential mineral, o: accessory mineral, 2: two or more generations, r: retrograde

*: Samples analyzed on the electron microprobe

Abbrev	Almandine	Crd	Cordierite	Mu	Muscovite
Amoh	Amphibole	Ep	Epidote	Plg	Plagioclase
An	Anorthite	Gt	Garnet	Qtz	Quartz
Bi	Biotite	Kfs	K-feldspar	Sill	Sillimanite
Chl	Chlorite	Ky	Kyanite	Sp	Spinel
Co	Corundum				

completely pinitized cordierite. It is difficult to decide whether these sillimanite fibers were formed from muscovite or decomposed to muscovite.

Spinel: Green spinel was observed as a modally subordinate mineral. Spinel only occurs as inclusion in cordierite (Fig. 2a) and in the neighbourhood of sillimanite overgrowing cordierite.

Biotite: The modal abundance of biotite varies strongly with the rock type, but it is a major constituent in all samples. Biotite was found in contact with all other minerals.

Muscovite: Muscovite is a rather late mineral replacing cordierite (Fig. 2f) and often coexists with garnet (Fig. 2b).

298

Metamorphic evolution of gneisses of NW Spitsbergen



Fig. 2 Typical textures from metapelitic rocks:

299

			(Garnet			*	Bi	otite*		
Sample Comment	12/7 rim	12/6 core	38/2 rim	38/6 core	66/6 rim	66/2 core	12/1	38/1	38/15 i**	66/11	66/2 i
Si02	37.41	37.59	37.21	38.35	37.09	37.75	34.32	34.91	36.19	34.23	35.33
T102	21 16	20 86	00.15	00.00	01 04	20 07	3.89	2.29	4.39	5.20	3.57
A1203	22.10	20.80	22.15	20.99	21.34	20.67	19.09	18.02	18.08	17.86	18.27
MoO	3 52	4 22	2 12	34.90	5 76	33.91	22.44	23.32	20.93	22.80	22.79
MaQ	2 68	2 79	3 40	2.98	1 52	2 50	6 53	8 00	8 09	5 46	7 08
CaO	1.26	1.39	0.76	0,93	0.94	0.92	0.50	0.00	0.00	5.40	7.00
Na ₂ O							0.14	0.33	0.22	0.17	0.16
κ ₂ Ō							10.15	7.25	8,78	9.34	9.30
Total	99.92	100.62	99.70	100.48	100.08	99.43	96.56	95.12	96.68	96.25	97.29
Formula	12 (0)					22 (0)					
Si	3 015	3 017	2 931	3.014	3.006	3.054	5.267	5.428	5 441	5.329	5 392
Ti	0.040	0.047		0.02.	0.000		0.449	0.268	0.496	0.609	0.410
AI	2.010	1.973	2.057	1.944	2.039	1.971	3.453	3.303	3.204	3.278	3.286
Fe++	2.284	2.266	2.243	2.293	2.266	2.294	2.880	3.032	2.631	2,968	2.908
Mn	0.240	0.287	0.141	0.156	0.395	0.240			0.020		
Mg	0.322	0.334	0.399	0.349	0.184	0.301	1.494	1.854	1.813	1.267	1.610
Ca	0.109	0.120	0.064	0.077	0.082	0.080			o		
Na						•	0.042	0.099	0.064	0.051	0.047
к 							1.987	1.438	1.684	1.855	1.810
Total	7.98	8.00	7.84	7.83	7.97	7.94	15.57	15.42	15.33	15.38	15.46
A] m	77.3	75.4	78.8	79.8	77.4	78.7					
Spess	8.1	9.5	5.0	5.4	13.5	8.2					
Pyr	10.9	11.1	14.0	12.1	6.3	10.3					
Crear	37	1 0	2 2	27	28	27					

Tab. 2 Representative microprobe analyses of Spitsbergen minerals.

	PI	LAGIOCLA	SE	K-FEL(DSPAR	SILLIN	IANITE	SPINEL		CORDIERI	TE
Sample	12/2	38/3	66/9	12/4	38/1	12/1	38/1	38/2	12/1	38/4	66/8
Si02 Al203 Fe0 FE203	59.14 25.58	61.99 23.37	62.63 22.59	63.66 18.42	66.36 18.06	36.18 59.12 1.18	36.80 59.19	0.00 56.35 33.20	47.59 32.29 11.01	49.05 32.33 9.32	48.53 31.91 10.27
MgÕ CaO Na ₂ O K ₂ O Cr ₂ O ₃ ZnO	6.81 6.97 0.31	5.87 7.97 0.24	5,48 7.60 0.16	0.46 1.40 15.71	0.00 2.94 11.82	0.23	0.07	1.70 3.32	6.24 0.09 0.04	7.74 0.18 0.00	5.82 0.40 0.06
Total	98.81	99.44	98.46	99.65	99.18	96.71	96.06	99.00	97.26	98.61	96.99
Formula		12 (0)		8 ((0)	5 (0)	16 (0)		18 (0)	
Si Al Fe++ Fe+++	2.662 1.357	2.763 1.228	2.806 1.193	2.965 1.011	3.033 0.973	1.015 1.953 0.028	1.033 1.955	0.000 7.690 3.219 0.169	5.010 4.007 0.969	5.052 3.925 0.803	5.090 3.945 0.901
Mg Ca Na K Cr Zn	0.328 0.608 0.018	0.280 0.689 0.014	0.263 0.660 0.009	0.023 0.126 0.933	0.000 0.260 0.689	0.010	0.003	0.530 0.156 0.284	0.979 0.018 0.005	1.188 0.034 0.000	0.910 0.081 0.008
Total	4.97	4.97	4.93	4.93	4.96	3.01	2.99	11.99	10.99	11.00	10.94
An Ab Or	34.4 63.7 1.9	28.5 70.1 1.4	28.2 70.8 1.0	2.1 11.6 86.2	0.0 27.4 72.6						

* Total includes minor amounts of F and Cl, ** i: inclusion in garnet

Chlorite: Chlorite is clearly a late alteration product and replaces biotite in some samples.

Mineral chemistry

The chemical composition of the minerals was obtained by using an automated ARL microprobe equipped with a Tracor Northern energy dispersive detector system and five crystal X-ray spectrometers at the Institut für Mineralogie und Petrographie at ETH Zurich. Analyzed samples are marked with a subscript * on Table 1 and representative mineral analyses are given in Table 2. The extent of MgFe-1 substitutions is shown in the histograms in Fig. 3.



Fig. 3 Iron-magnesium variation histograms for biotite, garnet, cordierite and spinel (SPI 38). Smallest dash represents one analysis. r: rim, c: core, i: inclusion in garnet.

Garnet: The single phase garnets show no or only a very weak chemical zonation (Fig. 4) with some garnet cores slightly enriched in pyrope and depleted in almandine component. Most garnets (except SPI 39 and 59) show no zonation with respect to the grossular component and show only a slight variation in the spessartine component which may be enriched in the outermost parts of the garnets. This enrichment in manganese can best be observed in garnets which display characteristic resorption features and have very irregular outlines. This pattern of Mn enrichment has been reported to be a typical resorption feature by GRANT and WEIBLEN (1971) and DEBETHUNE et al. (1975).

In some samples (e.g. SPI 37, SPI 59) two growth phases of garnets can be observed (Fig. 2e and f). The analyzed polyphase garnets may or may not show a zonation. There is, however, no discontinuity in the chemical composition from core to rim. Examples are given in Figure 4. No zoning can be observed in polyphase garnets from cordierite bearing samples (e.g. SPI 37) from the northern, higher temperature part of the area studied. WOODSWORTH (1977) has shown that prograde garnet zoning profiles may be smoothed or homogenized at high grade. Such homogenization could have taken place at the temperatures attained (700°C) during metamorphism (see below). However, polyphase garnets from biotite gneiss samples from the south-western part of the area (e.g. SPI 59) show a clear zoning pattern with respect to their Ca, Fe and Mn zoning.





Fe/(Fe+Mg) ratios vary within narrow limits in each sample and between 0.75 to 0.94 on the whole. Some samples show a bimodal iron-magnesium distribution (Fig. 3) in garnet which is related to differing core and rim compositions.

Biotite: Fe/(Fe+Mg) ratios of 0.5-0.75 were found in all analyzed samples (Fig. 3). Biotite inclusions in garnet have lower Fe/(Fe+Mg) ratios than the matrix biotites.

Cordierite: Cordierites are chemically homogeneous with Fe/(Fe+Mg) ratios between 0.4 and 0.5, irrespective of local mineral assemblage or grain size. In some samples cordierite shows a strong alteration to yellow pinite. Most analyses show up to 0.50 wt% Na₂O. The water content in cordierite has not been determined.

Spinel: All spinels analyzed are essentially hercynite—Mg-spinel solutions with up to 13 mol.% spinel endmember. Gahnite (up to 7 mol.%), magnetite (up to 3 mol.%) and chromite (up to 2 mol.%) components total less than 12 mol.%.

Plagioclase: The anorthite content of plagioclases varies about 10 mol.% within each sample but shows no systematic zoning within each grain and lies between An(20) and An(35) in the different samples analyzed. No regional changes in plagioclase composition were observed.

Alkalifeldspar: The composition of K-feldspar (microperthite) ranges from Or(70) to Or(90).

White mica: Most analyzed muscovites contain 4 to 7% celadonite solid solution.

Chlorite: The chlorite which is a late stage mineral and formed as alteration product from biotite is of ripidolitic composition.

Geothermometry and geobarometry

The widespread mineral assemblage cordierite – biotite – garnet – plagioclase – sillimanite – quartz can be used to deduce estimates of pressure and temperature conditions under which these rocks equilibrated. The calibrations of the equilibria listed below were used for PT-calculations (exchange component notation adopted from THOMPSON (1982).

(1)	MgFe-1(bi) = MgFe-1(gt)	Thompson (1976)
		INDARES and MARTIGNOLE (1985)
(2)	MgFe-1(crd) = MgFe-1(gt)	Thompson (1976)
(3)	an(plag) = gross(gt) + sill + qtz	ARANOVICH and PODLESSKIY (1980)
3		GHENT et al. (1979)
(4)	Fe-crd(crd) = alm(gt) + sill + qtz	Thompson (1976)
		NEWTON and WOOD (1979)

The coexistence of K-feldspar and plagioclase in sample SPI 38 permits the temperature estimation from the additional equilibrium

(5) NaK-1(K-fsp) = NaK-1(plag)

STORMER and WHITNEY (1977).

Temperatures calculated from equilibrium 1 are shown in Fig. 5 and those from equilibrium 2 in Fig. 6. The scatter of the calculated garnet-biotite rim temperatures is of about 100°C around mean temperatures of 650 to 700°C in the southern part of the studied area and 700 to 750°C in the central and northern part. To deduce statistically valid mean temperatures, the average value from several samples has been taken for the northern and southern part of the studied area. Garnet-cordierite temperatures generally overlap well with garnet-biotite temperatures but show a slightly smaller scatter. Garnet core temperatures calculated from biotite or cordierite inclusions in garnet (e.g. SPI 18,



Fig. 5 Calculated temperatures from garnet-biotite Fe-Mg exchange thermometer at 5 kbar. Calibration of THOMPSON (1976). r: rim, c: core.



Fig. 6 Calculated temperatures from garnet-cordierite Fe-Mg exchange thermometer at 5 kbar. Calibration of THOMPSON (1976). r: rim, c: core.

36, 37, 38, 66) are about 50°C higher than rim temperatures. There are, however, exceptions (e.g. SPI 12, 39) which show the opposite trend with higher rim temperatures.

The Ca content in the garnets is rather low and can, therefore be ignored but the rather high Mn content in the garnets indicates that the data have to be interpreted very carefully (NEWTON and HASELTON 1981). Temperatures up to 800 °C are, however, unrealistically high, because these rocks do not contain the assemblage orthopyroxene + quartz and the calculated phase relations also contradict such high temperatures. Temperatures calculated according to the calibration of INDARES and MARTIGNOLE (1985) which accounts for deviations from ideality in both garnet and biotite solid solution, are about 50 °C lower than the values given above according to the THOMPSON (1976) calibration and slightly decrease the scatter in the data to about 60 °C. The former data are consistent with temperature estimates of 640 to 680 °C from dolomitic marbles from the same area (BUCHER 1981).

At an average metamorphic temperature increasing from the south towards

the north from 675 to 725°C, garnet-plagioclase pressures of 4 to 5 kbar (at 700°C) have been deduced according to the calibration of ARANOVICH and POD-LESSKIY (1980) as well as GHENT et al. (1979).

Plagioclase inclusions in garnet allow the deduction of garnet core pressures assuming that an alumosilicate (sill or ky) was also present in the assemblage when the garnet cores formed. This assumption may be valid because there are rocks showing sillimanite patches pseudomorphous after kyanite, which indicates that kyanite was present at an earlier stage of metamorphism. Kyanite relicts have also been reported from neighbouring areas (OHTA 1978). Some samples with a suitable assemblage (e.g. SPI 66) indicate retrograde garnet growth with core pressures of up to 2 kb higher than garnet rim pressures at decreasing metamorphic temperatures (see above). However, there are two samples (SPI 12, 39) indicating garnet growth with increasing temperature but there is no information from inclusion assemblages about the pressure under which the garnet cores formed.¹

In contrast to equilibrium 3 which is strongly temperature dependent, the assemblage cordierite + garnet + sillimanite + quartz represents an excellent geobarometer. The water content of cordierite, however, plays an important role in the positioning of equilibrium 4. Assuming dry cordierite, using the calibration of NEWTON and WOOD (1979), the regional pressures are between 4 and 5 kbar, which is consistent with the calibration of ARANOVICH and PODLESSKIY (1980) and GHENT et al. (1979). Assuming H₂O saturated cordierite the pressures are between 5 and 6 (-7) kbar which is consistent with the calibration of THOMPSON (1976).



Fig. 7 Calculated pressures from the assemblages garnet + plagioclase according to ARANOVICH and PODLESSKIY (1980) (A) and garnet + cordierite + sillimanite + quartz according to THOMPSON (1976) (T) and NEWTON and WOOD (1979) (NW) for sample SPI 12. Temperatures given according to THOMPSON (1976).

¹ The pressure change indicated by the garnet zoning has been calculated with the program PTPATH (written by F. Spear) and suggests a decompression with heating path of metamorphism for these samples.





Coexisting K-feldspar-plagioclase pairs in sample SPI 38 revealed solvus temperatures of 720°C at 5 kbar using the sanidine calibration of STORMER and WHITNEY (1977). Microcline temperatures are 80–90°C higher.

Temperatures calculated from calibrations 1 and 2 and pressures calculated from equilibrium 3 and 4 are represented in Fig. 7 for one typical sample (SPI 12).

The results for the thermobarometry are summarized on Fig. 8 for four of the samples separately. The pressure-temperature conditions have been plotted and permit the selection of a reasonably well defined PT-range for the last supposed chemical equilibration of the area studied.

Interpretation of textures and mineral reactions

A commonly observed textural relation involving cordierite + partially resorbed garnet + sillimanite + quartz suggests that cordierite does not represent a cogenetic mineral with garnet. The observed strong resorption of retrograde garnet and its replacement by cordierite can be explained as the result of the reaction

$$2 gt + 4 sill + 5 qtz \rightarrow 3 crd$$
(4)

The equilibrium conditions of this cordierite-producing reaction have been used in the previous section as a geobarometer and are shown on Fig. 9 for sample SPI 38 forming a low grade boundary to the estimated PT-conditions.

All reactions calculated in this section are based on the data of HELGESON et al. (1978). The free energy data of almandine, hercynite and Fe-cordierite given in DROOP and BUCHER (1984) as well as their activity models (BUCHER and DROOP, 1983) were used for the calculations.

The observed assemblage garnet + K-feldspar + sillimanite + biotite + quartz indicating simultaneous growth of biotite and sillimanite replacing garnet can be explained by the reaction

$$1 \text{ gt} + 1 \text{ kfs} + 1 \text{ H}_2\text{O} \rightarrow 1 \text{ sill} + 1 \text{ bi} + 2 \text{ qtz}$$
(6)

The retrograde reaction 6 leading to the simultaneous formation of sillimanite + biotite is shown on Fig. 9 to give a low temperature boundary to the independently inferred PT-conditions. The reaction curve is calculated for sample SPI 12 and $a_{H_2O} = 1$. The assumption for $a_{H_2O} = 1$ is based on two observations: only PT-estimates using wet cordierite give a consistent PT-box (Fig. 9) and, secondly, the position of the reaction curve indicates that all H₂O stored in a probable melt phase has been expelled to enable re-hydration reactions.

Simultaneous growth of cordierite and biotite observed in some samples (e.g. SPI 38) may be explained by hydration reactions of the type

Klaper, E.M.



Fig. 9 Summary of the metamorphic conditions. Dashed box: PT-conditions from thermobarometry. Granite solidus after WYLLIE (1977), equilibrium conditions of reactions 6 (SPI 12), 8 (SPI 38). The low temperature limit for the reaction assemblage orthopyroxene (OPX) + quartz = anthophyllite (ANT) which has been calculated using the dataset of HELGESON et al. (1978), has not been reached.

$6 \text{ gt} + 4 \text{ kfs} + 4 \text{ H}_2\text{O} \rightarrow 3 \text{ qtz} + 3 \text{ crd} + 4 \text{ bi}$	(7a)
(in quartz saturated rocks)	
9 gt + 5 kfs + 3 sill + 5 $H_2O \rightarrow 6$ crd + 5 bi	(7b)
(in guartz deficient rocks)	

All these textures indicate that cordierite formation may have resulted from decompression and during a period where re-hydration reactions were active. The conclusion of decompression formation of cordierite is further supported by spinel + cordierite textures (Fig. 2a). Spinel mantled by cordierite has been interpreted as the educt mineral of reaction 8:

$$5 \operatorname{qtz} + 2 \operatorname{sp} \rightarrow 1 \operatorname{crd}$$
 (8)

Cordierite formation, therefore, requires decompression and the rocks were at higher pressures prior to equilibration at PT-conditions calculated from thermobarometry. The reaction curve calculated for water-saturated cordierite passes through the determined PT-box (Fig. 9) and suggests high H_2O activity for the retrograde equilibration phase. This interpretation is again supported by the occurrence of kyanite bearing rocks west of the studied area and of rocks with sillimanite patches pseudomorphous after kyanite (HJELLE, in OHTA, 1974a). Uncertain kyanite relicts have been observed in one sample (SPI 38). This supports the suggestion that the rocks also underwent the reaction

$$1 \text{ ky} \rightarrow 1 \text{ sill}$$
 (9)

This reaction also indicates that metamorphic pressures reached higher peak values than the ones determined in this study.

The PT-conditions deduced for garnet equilibration are not consistent with the stable coexistence of muscovite + quartz. Experimental work (HOLDAWAY and LEE 1977) also demonstrates that cordierite, almandine and muscovite are not stable together. However, when almandine exists as an excess phase due to high Mn and/or Ca component in garnet these minerals may coexist (OSBERG 1971, OKRUSCH 1971). Texturally, the observed muscovite is clearly a late, retrograde mineral overgrowing cordierite or garnet and has probably formed from the following retrogressive hydration reaction

$$1 \text{ kfs} + 1 \text{ sill} + 1 \text{ H}_2\text{O} \rightarrow 1 \text{ mu} + 1 \text{ qtz}$$
(10)

and possibly also according to the reactions

$$1 \text{ gt} + 2 \text{ kfs} + 3 \text{ H}_2\text{O} \rightarrow 1 \text{ mu} + 1 \text{ bi} + 3 \text{ qtz}$$
(11)

$$3 \operatorname{crd} + 8 \operatorname{kfs} + 14 \operatorname{H}_2 O \rightarrow 6 \operatorname{mu} + 2 \operatorname{bi} + 15 \operatorname{qtz}$$
(12)

The equilibrium conditions for reaction 10 are shown on Fig. 9 labelled as mu + qtz – out curve for unit water activity.

The PT-conditions calculated from thermobarometry are above the conditions of the H₂O saturated muscovite-granite solidus which is shown on Fig. 9 according to WYLLIE (1977). Partial melting is in fact indicated by the widespread migmatitic textures in quartzo-feldspathic gneisses of NW Spitsbergen. The reactions involving K-feldspar formulated on the previous pages may thus involve a melt phase rather than solid K-feldspar. This melt phase, containing plagioclase and quartz as well as minor amounts of mafic components, probably stored a large amount of H₂O. This H₂O has been given off by the earlier prograde dehydration reactions such as dehydration of muscovite, biotite and possibly staurolite. Along the cooling and uplift path the H₂O stored in the melt was released and excess H₂O expelled from the migmatites increased the water activity as soon as the rocks came close to the muscovite-granite solidus and was then built into the newly formed minerals.

309

Summary and geological interpretation

Metapelitic gneisses from NW Spitsbergen typically contain cordierite + garnet + biotite + sillimanite + plagioclase + k-feldspar + quartz \pm spinel. The PT-range deduced from these rocks by thermobarometry indicate isobaric conditions of 5 to 6 kbar (assuming H₂O saturated cordierite) at a temperature between 650 to 700 °C). A slight increase in the average metamorphic temperature of about 50 °C from the south to the north can be observed.

Figure 10 indicates a possible PT-path of the rocks of the lower Hekla Hoek succession. Only a small portion of this path is well constrained. This part is represented by decompression with increasing (early garnets) or decreasing (late garnets) temperatures, respectively. An additional phase of heating may be postulated because of sillimanite fibers overgrowing partly pinitized cordierite in the northern part of the studied area. This heating can possibly be correlated with the intrusion of grey granites which can be observed prior to the Hornemantoppen intrusion. A weak late heating related to this intrusion is responsible for late stage chloritization of biotite.

It is clear from the observed textures, relict minerals and deduced mineral reactions that the inferred conditions of metamorphism could not have been the maximum conditions experienced by these rocks and that they have endured



Fig. 10 Possible PT-path for metapelitic rocks of the lower Hekla Hoek formation in northwestern Spitsbergen. Isotopic age data taken from HJELLE (1979) and GAYER et al. (1966).

both, higher pressures and possibly also higher temperatures prior to the PTconditions calculated from standard geothermobarometry. If, as suggested, the rocks have been at higher PT-conditions at some earlier stages during metamorphism, this implies that conditions inside the PT-field for granitic melt formation prevailed (Fig. 9), because thermobarometry on biotite, cordierite and plagioclase inclusions in most garnet porphyroblasts suggest a retrograde growth path almost parallel to the kyanite-sillimanite transformation curve.

At metamorphic peak conditions of the rocks, probably a low water activity prevailed. The H_2O required to make the high degree of equilibration at 5-6 kbar and 700 °C possible was presumably released by crystallizing migmatitic melts present in the quartzo-feldspathic gneisses.

Although the maximum conditions reached are unknown and earlier stages in the history of the rocks are poorly preserved, it seems reasonable to assume that conditions close to granulite facies have been reached by the lower Hekla Hoek rocks of NW Spitsbergen. It is also significant that corundum as additional petrogenetic indicator mineral has been reported by OHTA (1974b) at one locality occurring in a layered gneiss sequence in the assemblage co + sp + crd+ bi + plag + qtz.

A metamorphic pressure of about 4 kbar at 640–680 °C was estimated for the positions of equilibrium curves and invariant points in PT-space deduced from siliceous dolomitic marble assemblages within the same area (BUCHER 1981). This pressure of 4 kbar is in agreement with pressures calculated from metapelitic rocks assuming dry cordierite. A pressure of 4 kbar is also compatible with the alumosilicate triplepoint data of HOLDAWAY (1971). Based on the data of RICHARDSON et al. (1971), OHTA (1974b) deduced a higher pressure of 5.5 kbar at roughly the same temperature (650–700 °C) which is consistent with the calculations for wet cordierite in this study.

Acknowledgements

A travel grant by the "Schweizerische Naturforschende Gesellschaft" is gratefully acknowledged.

I thank all members of the expedition for their support. Particularly, I am indepted to B. and R. Eichin for their company in the field and to the leader of the expedition, Dr. L. Schmutz who also helped collecting the samples. I appreciate the comments on an earlier version of this manuscript by Dr. J. Ridley and Dr. G.T.R. Droop. The critical review of Dr. Y. Ohta is gratefully acknowledged.

References

ARANOVICH, L. YA. and PODLESSKIY, K. K. (1980): The garnet-plagioclase barometer. Doklady, Earth Sci. Sec. 251, 101-103.

BUCHER-NURMINEN, K. (1981): Petrology of chlorite-spinel marbles from NW-Spitsbergen (Svalbard). Lithos 14, 203-213.

- BUCHER-NURMINEN, K. and DROOP, G.T.R. (1983): The metamorphic evolution of garnet-cordierite-sillimanite gneisses of the Gruf-Complex, Eastern Pennine Alps. Contrib. Mineral. Petrol. 84, 215-227.
- DEBETHUNE, P., LADURON, D. and BOQUET, J. (1975): Diffusion processes in resorbed garnets. Contrib. Mineral. Petrol. 50, 197-204.
- DROOP, G.T.R. and BUCHER-NURMINEN, K. (1984): Reaction textures and metamorphic evolution of sapphirine-bearing granulites from the Gruf Complex, Italian central Alps. J. Petrol. 25, 766-803.
- GANGULY, J. (1979): Garnet and clinopyroxene solid solutions and geothermometry based on Fe-Mg distribution coefficients. Geochem. Cosmochem. Acta 43, 1021-1029.
- GAYER, R.A., GEE, D.G., HARLAND, W.B., MILLER, J.A., SPALL, H.R., WALLIS, R.H. and WINSNES, T.S. (1966): Radiometric age determinations on rocks from Spitsbergen. Norsk Polar Institutt, Skrifter 137, 1-39.
- GHENT, E.D., ROBBINS, B.D. and STOUT, M.Z. (1979): Geothermometry, geobarometry and fluid compositions of metamorphosed calc silicates and pelites, Mica Creek, British Columbia. Am. Mineralog. 64, 874-885.
- GRANT, J.A. and WEIBLEN, P.W. (1971): Retrograde zoning in garnet near the second sillimanite isograd. Am. J. Sci. 270, 281-297.
- HELGESON, H.C., DELANY, J.M., NESBITT, H.W. and BIRD, D.K. (1978): Summary and critique of the thermodynamic properties of rock-forming minerals. Am. J. Sci. 278-A, 1–229.
- HJELLE, A. (1974): The geology of Danskoya and Amsterdamoya. In: Hjelle, A. and Ohta, Y. (eds.): Contribution to the geology of northwestern Spitsbergen. Norsk Polar Institutt, Skrifter 158, 5-38.
- HJELLE, A. (1979): Aspects of the geology of northwestern Spitsbergen. Norsk Polar Institutt, Skrifter 167, 37-62.
- HJELLE, A. and OHTA, Y., eds. (1974): Contribution to the geology of northwestern Spitsbergen. Norsk Polar Institutt, Skrifter 158, 1-107.
- HOLDAWAY, M.J. (1971): Stability of andalusite and the aluminium silicate phase diagram. Am. J. Sci. 271, 97-131.
- HOLDAWAY, M.J. and LEE, S. M. (1977): Fe-Mg cordierite stability in high grade pelitic rocks based on experimental, theoretical and natural observations. Contrib. Mineral. Petrol. 63, 175-198.
- INDARES, A. and MARTIGNOLE, J. (1985): Biotite-garnet geothermometry in the granulite facies. The influence of Ti and Al in biotite. Am. Mineralog. 70, 271–278.
- NEWTON, R.C. and HASELTON, H.T. (1981): Thermodynamics of the garnet-plagioclase-Al₂SiO₅-quartz geobaromter. In: NEWTON, R.C., NAVROTSKY, A. and WOOD, B.J. (eds.): Thermodynamics of minerals and melts. Springer Verlag, Heidelberg.
- NEWTON, R.C. and WOOD, B.J. (1979): Thermodynamics of water in cordierite and some petrological consequences of cordierite as a hydrous phase. Contrib. Mineral. Petrol. 68, 391-405.
- OHTA, Y. (1974a): Tectonic development and bulk chemistry of the rocks from the Smeerenburgfjorden area, Spitsbergen. In: HJELLE, A. and OHTA, Y. (eds.): Contribution to the geology of northwestern Spitsbergen. Norsk Polar Institutt, Skrifter 158, 69–106.
- OHTA, Y. (1974b): Geology and structure of the Magdalenefjorden area, Spitsbergen. In: HJELLE, A. and OHTA, Y. (eds.): Contribution to the geology of northwestern Spitsbergen. Norsk Polar Institutt, Skrifter 158, 39-68.
- OHTA, Y. (1978): Caledonian metamorphism in Svalbard, with some remarks on the basement. Polarforschung 48, 78-91.
- OKRUSCH, M. (1971): Garnet-cordierite-biotite equilibria in the Steinach aureole, Bavaria. Contrib. Mineral. Petrol. 32, 1-23.
- OSBERG, P.H. (1971): An equilibrium model for Buchan-type metamorphic rocks, south central Maine. Am. Mineral. 56, 569-576.

RICHARDSON, S. W., GILBERT, M. C. and BELL, P. M. (1968): Kyanite-sillimanite equilibrium between 700°C and 1500°C. Am. J. Sci. 266, 513-541.

STORMER, J.C. and WHITNEY, J.A. (1977): Two feldspar geothermometry in granulite facies metamorphic rocks. Sapphirine granulites from Brazil. Contrib. Mineral. Petrol. 65, 123-133.

THOMPSON, A.B. (1976): Mineral reactions in pelitic rocks: II. Calculations of some PTX(Fe,Mg) phase relations. Am. J. Sci. 276, 425-454.

THOMPSON, J. B. (1982): Composition space: an algebraic and geometric approach. In: FERRY, J. M. (ed.): Characterization of metamorphism through mineral equilibria. Rev. Mineral. 10, 1–31.

WOODSWORTH, G.J. (1977): Homogenisation of zoned garnets from pelitic schists. Can. Mineral. 15, 230-242.

WYLLIE, P.J. (1977): Crustal anatexis: an experimental review. Tectonophysics 43, 41-71.

Manuscript received December 8, 1986; revised manuscript accepted January 23, 1987.