

Geochemistry of meta-lamprophyres from the Central Swiss Alps

Autor(en): **Oberhänsli, Roland**

Objektyp: **Article**

Zeitschrift: **Schweizerische mineralogische und petrographische Mitteilungen
= Bulletin suisse de minéralogie et pétrographie**

Band (Jahr): **66 (1986)**

Heft 3

PDF erstellt am: **11.09.2024**

Persistenter Link: <https://doi.org/10.5169/seals-50898>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Geochemistry of meta-lamprophyres from the Central Swiss Alps

by *Roland Oberhänsli*¹

Abstract

Geochemical investigations show that lamprophyres behave as nearly closed chemical systems under metamorphic conditions; except for volatiles, uranium and trace elements concentrated in feldspars (Sr, Ba). Trace element analyses show an enrichment of mobile elements (Sr, K, Rb, Ba) compared to a primitive basaltic composition. Cr and Ni distribution compared to Mg and Fe distribution indicates that the evolution of lamprophyric magmas starts from evolved basaltic melts. On the one hand REE patterns indicate that lamprophyric and calc-alkalic granitic magmas cannot be cogenetic. On the other hand, the REE patterns do not reflect primitive mantle compositions.

A mantle origin is proposed for the lamprophyres of the Swiss Alps. Due to dehydration of a subducted slab of oceanic crust, the overlying lithospheric mantle wedge was metasomatized. At deeper levels, melting of the subducted oceanic crust occurred combined with partial melting of the upper mantle. Mixing of melts of the metasomatized mantle with partial melts from the down-going plate could have produced lamprophyric magmas. Stagnation of such magmas at different levels below and within the crust yields diverse lamprophyre magmas, e.g. minettes, vogesites, etc. Later tectonic activity, a change from compressional to tensional stress fields (probably due to isostatic uplift of the orogenic calc-alkalic batholiths and thrusting) allowed the emplacement of the lamprophyric dikes.

Keywords: Geochemistry, lamprophyres, Central Alps, Switzerland.

Introduction

The origin of lamprophyric dikes is still very controversial. Extensive reviews of the vast literature on the genesis of lamprophyres are presented by WIMMENAUER (1973) and ROCK (1984). Among the many current genetic ideas and models three major opinions emerge:

1. Models involving magmatic processes, such as differentiation, fractional crystallization and/or partial melting:

¹ Mineralogisch-petrographisches Institut der Universität, Baltzerstr. 1, CH-3012 Bern.

- Lamprophyres represent remobilized mafic residuals of granitic, anatectic or tholeiitic melts without participation of new primitive basaltic magma.
2. Models considering complete or selective contamination, assimilation or hybridization with or without volatile enrichment at different crustal levels: Lamprophyres represent reaction products of a basaltic magma with K-rich material in the crust.
 3. Models preferring partial melts of unaffected parents from deeper levels as well as metasomatically enriched lithospheric mantle: Lamprophyres are products of more or less primitive partial melts.

BEGER (1923) presented a world-wide compilation of lamprophyres with 382 analyses, of which 52 were from the Central Swiss Alps. At that time Beger referred to the Gotthard region as one of the best known and best investigated areas with respect to lamprophyric dike rocks. This statement was mainly based on the classical studies of WEBER (1904), GRUBENMANN (1919) and SONDER (1921). Beger's assessment stands in remarkable contrast to the extensive recent compilation by ROCK (1984) in which not a single reference to lamprophyres of the Central Swiss Alps is to be found. Whether this shift in emphasis is due solely to the fact that modern research on the rocks has been largely confined to unpublished theses (KÜPFER, 1977; SCHALTEGGER, 1984; STEINER, 1984), or whether the metamorphic overprint has deterred those interested in the petrogenesis of lamprophyres is not clear. However, the extent of alteration and the geochemical consequences of the Alpine metamorphism on these rocks have been poorly documented until very recently. An extensive data collection by OBERHÄNSLI (1985) and the aforementioned theses provide the basis for a reassessment. The aim of this paper is to discuss these geochemical data on metamorphosed lamprophyres, especially from the area of the Central Swiss Alps belonging on the Hercynian orogenic cycle which escaped earlier Hercynian metamorphism. It is suggested here that the limited degree of bulk chemical change during metamorphism of lamprophyres does permit the use of geochemical data in tracing the primary origin of these rocks, even in metamorphic terranes. Such a model is developed for the suite of meta-lamprophyres from the Central Swiss Alps.

Material and methods

Mafic dikes and lamprophyres occur throughout the Swiss Alps in the crystalline massifs and their gneissic envelopes as well as in the crystalline Penninic and Austroalpine nappe systems. With few Tertiary exceptions they mark the latest (or last) magmatic event of the Hercynian orogenic cycle. A compilation of lamprophyre occurrences from the Swiss Alps is given in figure 1 and table 1.

Lamprophyres were investigated along a N-S profile from the Schwarzwald to the Ticino. Lamprophyres were sampled in the unmetamorphosed massifs e.g. in the Vogesen (Vosges) and the Schwarzwald (Black Forest) and were examined together with meta-lamprophyres from the increasingly overprinted external (Aarmassif) and internal (Gotthardmassif) massifs and the crystalline Penninic nappes (Ticino) from the Alps. Chemical analyses from lamprophyres of the Punteglias area (samples A 16-26) are taken from the unpublished Ph. D. thesis of T. KÜPFER (1977).

For comparison the distribution of meta-lamprophyres along an E-W traverse from the Mont Blanc to the lower Engadine window was also studied.

Sample locations are given in a tectonic sketch map (Fig. 2) showing Vogesen, Schwarzwald, Aar- and Gotthardmassif, as well as the Penninic realm of the Ticino. Short petrographic descriptions are compiled in table 2.

Mineralogical and metamorphic aspects of the meta-lamprophyres from the Central Swiss Alps shall be discussed elsewhere (OBERHÄNSLI, in prep.).

USE OF NOMENCLATURE

In the Vogesen and the Schwarzwald the terms lamprophyre, semilamprophyre and anchibasalt as defined by WIMMENAUER (1973), as well as the terms minette, vogesite, spessartite and kersantite will be used. Where recognition is still possible in the Alps, the terms (minette, vogesite, etc.) are used despite of weak metamorphic overprinting (terms like meta-minette will not be introduced) and these rocks will also be referred to as lamprophyric dikes. With increasing deformation and metamorphic recrystallization, the distinction between the rock-types such as minette and vogesite is no longer possible. For such rocks the term meta-lamprophyres will be used. From a petrographical point of view, these meta-lamprophyres are chlorite-sericite schists or biotite gneisses.

ANALYTICAL METHODS

Bulk rock, major and trace elements were determined by X-ray fluorescence as described by DIETRICH et al. (1976) and NISBET et al. (1979).

The XRF analyses were performed with automatic Philips sequential spectrometers (PW 1450) at the Eidgenössische Landwirtschaftliche Versuchsanstalt Liebefeld, Köniz and at the Mineralogical Institute of the University of Fribourg. Trace elements were analysed on pressed powder pills by XRF at the Centre d'Analyse of the Geological Institute of the University of Lausanne.

Rare earth elements (REE) were analysed in collaboration with the Institute of Radiochemistry of the University of Bern, by radiochemical neutron activation analysis at the Federal Institute of Reactor Research, Würenlingen.




Fig. 1 Distribution of mafic dikes, lamprophyres and porphyritic diabases in the Central Alps. Data from literature and this study.




Fig. 2 Sample locality index map covering the Vogesen (V), the Schwarzwald (S), the Aar- (A) and Gotthard-massif (G) and the Ticino (T).

Tab. 1 Lamprophyre occurrences in the Swiss Alps, compiled from the literature. Detailed references can be obtained from the author by request.

Schwarzwald *****					
lamprophyre	Laufenburg Kaisten Leuggern	bi-gneiss granite	Niggli P., 1912 Suter H., 1924 * Büchi et al., 1984 * Müller W.H., 1984 Nagra, 1984 *	Laufenburg Nagra drill holes	BI PLG KFSP AP ab ser mu chl cc ep/czo sph mgt
=====					
Aarmassif *****					
minette (malchite)	Lauchernalp	banded amphibolites, biotite gneisses,	Zraggen P., 1975	Lötschental	KFSP BI DI -> ACT bi ser ep czo sph
spessartite	Schaffberg Kleines Nesthorn Distelberg Beichpass	northern gneisses of central Aargranite	Ledermann H., 1946 * Hügi et al., 1985		PLG HBL ZO EP CHL ->ACT
lamprophyre	Oberaletsch Aletschhorn	amphibolites	Niggli et al., 1930 *	Aletschhorn	
lamprophyre kersantite spessartite vogesite	SE Blatten Massaschlucht S Sparrhorn Grisighorn (lake)	augen gneisses central Aargranite, qtz dioritic gneisses	Labhard T., 1965 Steck A., 1966	Massa valley, N Naters Belalp, Grisighorn, N Naters	BI PLG HBL EP ab bi ep sph cc act gr
kersantite	N Bettmeralp	granitic gneisses	Zbinden P., 1950	Bettmeralp, Goms	
lamprophyre	Märjelensee	central Aargranite	Niggli et al., 1930 *	Aletsch glacier	
kersantite	Grünhörnligrat Finsteraarhorn Oberaarhorn Löffelhorn	northern Aargranite central Aargranite	Wyss R., 1932	Finsteraarhorn	BI KFSP AB-OLIG QZ EP ZO AP ZIR SPH ser
spessartite kersantite	S Steinlauhorn	central Aargranite, gneisses	Abrecht J., 1975	Oberhasli valley	PLG KFSP BI -> hbl ser
kersantit	Kalter Kehr	southern gneisses	Niggli C., 1965 Velde D. *	Obergoms, SW Gletsch	bi ep ab qz
lamprophyre	Trübtenbach Grimselsee Nollen (Grimsel) N Todtseeli Nägeliigrätli Belvedere (Furka) Furkahorn S Lochstock (Ross- mettlen) Schöllenen	central Aargranite, gneisses S central Aargranite	Fehr W., 1923, 22, 26 Hügi E., 1907 Huber M., 1922 Minder W., 1932 Niggli P., 1924 * Stalder A., 1964 (*) Nagra, 1981	between Grimsel and Andermatt	BI -> mu chl ep sph cc
	Trübtensee Trübtenjoch	southern Grimselgranite	Fehr W., 1926 Wyss R., 1932		BI KFSP AB-OLIG QZ EP/ZO AP ZIR SPH ser
	Urnerloch Teufelsboden	gneisses S of Grimsel- granite			
lamprophyre	Dammastock E Flanke	central Aargranite	Liechti H., 1933 *	Göschernalp valley	
lamprophyre	Sustenpass	Silberbergserie	Schaltegger U., 1984 *	Sustenpass	DI BI SP (OL) KFSP PLG CC AP EP act ser chl mu cc ep ilm
hornblende- minette	Winterberg W Flanke	central Aargranite	Fischer O., 1905 *	Voralp valley, Göschenen	BI HBL AP MGT mica ep ab sph
minette	Erstfeldertal	Blocks !!	Sauer A., 1905 Lotze R., 1914	Erstfeld valley	BI PYX KFSP KFSP BI DI -> chl
lamprophyre	SBB Kraftwerkstollen Amsteg	central Aargranite	Hügi E., 1923	Amsteg	
(1)	(2)	(3)	(4)	(5)	(6)
(1) Rock type		(3) Country rock	(5) Area		
(2) Locality		(4) Authors	(6) Mineralogy		

* chemical analysis
BI magmatic mineral
bi metamorphic mineral

Tab. 1 (continued).

lamprophyre	Maderanertal	central Aargranite	Sigrist F., 1947	Maderaner valley	
kersantite	Klüsertal-Fellital	central Aargranite	Pflugshaupt P., 1927 *	W Bristenstock	BI HBL PLG sph ser ilm
lamprophyre spessartite kersantite	Stöckligrat S Wicheltal Crispalt Giufstöckli Culmatsch Val Mila Piz Ault Piz Gendusas Val Calvaniev Val Flaci	southern Aargranite Giufsyenite southern gneisses	Weber F., 1904 * Niederer J., 1932 Huber W., 1948	Vorderrhein valley	HBL BI EP KFSP PLG MGT act sph ep ab ser BI PLG HBL QZ ep ser sph
spessartite kersantite	Val Gliems Cuolm Tgietschen Piz Cambrialas W Clavadi	granitic gneisses	Eugster H., 1951 Böhm C., 1986 Weber F.	Val Russein	PLG BI HBL QZ EP CHL ser act chl
lamprophyre	Piz Posta Biälla Crap Grond Val Ufiern	monzodiorite Puntegliasgranite	Wehrli L., 1896 Weber F., 1924 Küpfer T., 1974, 77 *	Val Punteglias	BI PLG KFSP DI QZ HBL PX ab ser bi act ep/zo sph
kersantite	Kreuzbach	syenite	Hügi T., 1941	Vättis	BI PLG KF SPH EP CC chl ser mgt
=====					
Gotthardmassif *****					
lamprophyre	Kummenhorntobel Stock SE Münster Distelgrat S Pizzo Gallina	Gotthard gneisses, 2-mica-gneisses	Oberholzer W., 1955	Obergoms	PLG HBL -> bi act chl
lamprophyre	Val Prosa, Nufenen	Bi-plg-gneisses	Schmidt & Preiswerk, 1908	E Nufenenpass, Bedretto valley	
lamprophyre	W Nufenenpass Gross Mutthorn Gerental Pne di Manio Alpe di Manegorio Rotondohütte Rottällhorn Ronggergrat Cavannapass Passo Lucendro Pizzo Lucendro Fibbia	Rotondogranite Scoresciagneiss Fibbia granitegneiss Prato series Gamsboden granitegneiss	Sonder R., 1921 * Fischer E., 1923 Eichenberger R., 1924 Eichenberger R., 1926 * Hafner S., 1958 Hafner S., 1975	N Bedretto valley, W of Gotthard pass	bi ab ep/zo qz act sph mu BI PLG mu ep/zo bi sph act ab BI KFSP MGT DI -> HBL sph ser chl bi ab
lamprophyre spessartite kersantite	Winterhorn Mätteli Alpe di Rodont Monte Prosa Gotthard Hospiz Fieud	aplitic granite Gamsbodengneiss bi-plg-gneiss	Walndziok P., 1906 Kriege L., 1916 Grubenmann U., 1919 * Sonder R., 1921 * Ambühl E., 1929 Koenigsberger J., 1930 Hofmänner F., 1964 *	Gotthard pass area	BI ab zo mgt mu bi hbl sph BI bi hbl act gr ab mu ep
lamprophyre kersantite	Passo Sella Unterapital Cadlimo Plauncacotschna	micaceous gneisses	Fritsch K., 1873 Kriege L., 1916 Ambühl E., 1929 Zweifel H., 1954	Gotthard area, E of the pass	BI act bi mu sph
spessartite kersantite	Tgiern Toma Piz Miez	bi-gneisse, amphibolite granodiorite	Huber H., 1943	Val Medel	BI PLG HBL zo mu ser act chl cc
lamprophyre	Val Cristallina	Cristallina granite	Holst W., 1913	Lukmanier area	BI PLG KFSP MGT ep ser ab
intermediate dykes	Rosbodenstock Piz Maler Val Nalps Piz Pazzola	bi-plg-gneisses	Niggli E., 1944 * Niggli E., 1948 * Arnold A., 1970	Vorderrhein valley	HBL BI PLG QZ ep act ab ser sph chl HBL PLG BI ab ?bi
=====					
(1)	(2)	(3)	(4)	(5)	(6)
(1) Rock type	(2) Locality	(3) Country rock	(4) Authors	(5) Area	(6) Mineralogy

Tab. 1 (continued).

Penninic nappes *****					
lamprophyre TERTIARY	N Colle Pallasina	metasediments	Dal Piaz G.V., 1979	Combinzone Ayas valley, N Italy	HBL BI KFSP DI AP SPH OL
lamprophyre	Poncione dei Laghetti	granodiorite	Ramsey & Allison, 1979 Steiner H., 1984 *	Maggia nappe Mattorello, S Airolo	bi act kfsp qz cc ep
lamprophyre		bi-plg-gneiss	Kriege L., 1916 Bossard L., 1929	Lukmanier nappe, Lago Ritom, S Airolo	
minette	Alp Tobel	Rofnaporphyre	Ruetschi G., 1903 *	Suretta nappe, Ausserferrera,	bi mu par' or
Austroalpine nappes *****					
lamprophyre	Mt. Morion	Arollagneiss	Stutz A., 1940 *	Dent Blanche nappe, Valpelline valley	PLG HBL BI ser zo
lamprophyre TERTIARY	Colle Pallasina	gneiss minuti	Dal Piaz G.V., 1979 *	Sesia zone, Ayas valley, N Italy	HBL BI KFSP DI AP SPH OL
lamprophyre	N Piz Crevasalvas N Piz Nalar Piz Caldras Piz d'Err Jenatsch area Piz Giumels S Val Alvra (Cresta Mora) N Piz Lagrev N & S Julier pass St. Moritz SE Piz Julier Piz Albana Piz Palaschin (Tsheppa)	granite diorite	Eugster & Frey, 1927 Cornelius H., 1935 Bühler C., 1983 *	Err nappe Bernina nappe	PLG PYX BI AP ILM ab act HBL PYX act ab
lamprophyre vogesite spessartite kersantite	Mortel N Piz Corvatsch Corvatsch W grat S Mortelhütte Alp Ota Roseggletscher Piz Chalchagn Muottas di Pontresina Berninastrasse Bovalhüttenweg	granite granite porphyre	Grubenmann U Staub R., 1915 * Münger R., 1982 * Müller D., 1982 *	Bernina nappe	HBL BI ILM act ep ser chl ab ?bi ILM -> sph mu ab ser act stilp chl
camptonite	Crasta Languard	Languardkristallin	Schuppli H., 1921 Scheidegger B., 1984 *	Languard nappe	?OL CPX HBL BI PLG ser bi Na-amph chl
vogesite	Alp Laret Belezza, Ardez-Fetan	Tasnagranite	Züst O., 1905 Grubenmann U., 1909 *	Tasna nappe	
Hbl vogesite	Griankopf	mica gneisses	Grubenmann U., 1909 *	Ützaldecke	KFSP HBL AUG BI chl mgt ep cc
Bergell Intrusion *****					
lamprophyre TERTIARY	Val Trubinasca Cima di Vazzeda	granodiorite marbles	Staub R. (Niggli et al., 1930 *) Wenk et al., 1977 *	NW Bergell	BI HBL EP AP SPH ILM
Southern Alps *****					
lamprophyre	Pallanza Mergozzo	schisti dei Laghi bi-plg-gneisses	Boriani et al., 1977	SW Lago Maggiore	
lamprophyre	Quinto Contone Cadenazzo S Gubiasca S Indemini Astano SE Sessa	bi-plg-gneisses Ceneri zone	Kelterborn P., 1923 Bearth P., 1932 * Bächlin R., 1937 Spicher A., 1940 Gräter P., 1951 Reinhard M., 1964 * Wenger C., 1983	Sotto Ceneri	PLG HBL AUG BI CC QZ AP SPH chl
(1)	(2)	(3)	(4)	(5)	(6)
(1) Rock type	(2) Locality	(3) Country rock	(4) Authors	(5) Area	(6) Mineralogy

Geochemistry

IDENTIFICATION OF META-LAMPROPHYRES

Alpine deformation and metamorphism render recognition of lamprophyric dike rocks difficult. Especially difficult is the distinction of meta-lamprophyres from porphyritic and diabasic dikes. Therefore, geochemical methods have been additionally used to identify strongly overprinted meta-lamprophyres.

The geochemical and mineralogical criteria introduced by WIMMENAUER (1973) and ROCK (1984) were employed to identify and classify lamprophyric rocks. In addition to these two screens, MANSON'S (1967) chemical criteria for basalt discrimination were applied. Because Wimmenauer's lamprophyre classification scheme is partly based on Manson's basalt discrimination criteria, there is generally a good correlation between both methods. Rock's modal, mineralogical, chemical and normative screens differ in many respects. For example, most samples from the highly metamorphosed region of the Ticino are not meta-lamprophyres according to the criteria of ROCK (1984), although they would be classified as semilamprophyres according to WIMMENAUER (1973). By contrast, the pyroxene spessartites from Punteglias (samples A 20 to A 26) do not violate Rock's criteria and would be classified as lamprophyres but they are basaltic after Manson's and anchibasaltic after Wimmenauer's classification.

Lamprophyres and meta-lamprophyres plotted in the triangle of normative Or-Pla-Maf (WIMMENAUER, 1973) show great variations (Fig. 3). Many samples fall into the fields of anchibasalts and diabases on the one hand, or into the fields of porphyrites and semilamprophyres on the other hand, instead of into the lamprophyre field.




Fig. 3 Norm (CIPW) classification for lamprophyres after Wimmenauer (1973). OR: orthoclase, AB: albite, AN: anorthite, OL: olivine, DI: diopside, HY: hypersthene, S: semilamprophyres and porphyrites, A: anchibasalts and diabases.




Fig. 4 Cr vs Ni distribution in lamprophyres and meta-lamprophyres. Different areas show varying compositional fields, reflecting the regional diversity of different magmatic suites.

None of the criteria used for different screens are conclusive for the identification of completely recrystallized meta-lamprophyric rocks. WIMMENAUER and HAHN-WEINHEIMER (1966) found that Cr and Ni contents of lamprophyres from the Schwarzwald and Vogesen are very high (Cr 440 ppm, Ni 100 ppm). Therefore, high concentrations of Cr and Ni were taken to identify meta-lamprophyres amongst the mafic dike rocks with high contents in alkalis from the Central Alps. The results are shown in figure 4. The mafic to lamprophyric dike rocks from different areas seem to have their particular Cr and Ni values, indicating a characteristic, slight to strong enrichment. For example samples from Schwarzwald are clearly higher in Cr than those from the Vogesen. The Aarmassif samples are intermediate and show two distinct populations: one with low Cr (< 3–180 ppm) and Ni (< 3–60 ppm) values representing anchibasaltic and semilamprophyric dikes (recognizable in part by their relic magmatic mineralogy) and another population enriched in Cr (200–700 ppm) and Ni (40–180 ppm), representing meta-lamprophyres. In the Gotthardmassif, dike rock samples have generally significantly lower Cr values compared to Vogesen, Schwarzwald and Aarmassif. Again, two populations representing lamprophyres (Cr ca. 200 ppm) and semilamprophyres (Cr ca. 40 ppm) can be observed. From this diagram one might conclude that every region (Vogesen, Schwarzwald, Aarmassif and Gotthardmassif) has its own characteristic pattern of Cr and Ni. These patterns might reflect the influence of igneous fractionation processes. However, in Alpine lamprophyric dikes and meta-lamprophyres it is not possible to assume Cr and Ni fingerprints for single intrusive complexes within one region, as has tentatively been done for restricted areas in the Schwarzwald (MÜLLER, 1982). In general, the high Cr concentrations (> 200 ppm) found in alkalic mafic dikes in the Alps are a strong indication for

the lamprophyric nature of dike rocks containing relics of magmatic biotite or hornblende.

Porphyritic, semilamprophyric or anchibasaltic dikes tend to show significantly lower Cr concentrations. This is also confirmed by data from the Austroalpine Bernina nappe (EIKENBERGER, 1984).

However, recognition and identification of the original character of meta-lamprophyres, e.g. minette, vogesite, kersantite and spessartite, on the basis of bulk rock analyses is rather difficult. Mean values of major elements for different minettes and kersantites were compiled by MÜLLER (1982). The most obvious difference between the two rock types is their Na₂O to K₂O ratio which is below 0.5 for minettes and above 0.5 for kersantites. In the metamorphic terranes of the Central Alps however, distinction between minettes, vogesites, kersantites, spessartites using this ratio is not reliable.

BULK CHEMICAL CONSEQUENCES OF METAMORPHISM

The discussion on recognition of meta-lamprophyres by geochemical methods showed that Cr is one of the elements almost unaffected by metamorphism.

As to uranium, recent investigations on lamprophyres from the Southern Schwarzwaldmassif report relatively high contents (BÜCHI et al., 1984). The U enrichment in lamprophyres from the Southern Schwarzwald with concentrations of 14–19 ppm U (BÜCHI et al., 1984), is confirmed in drill holes from northeastern Switzerland (Tab. 2). The U concentrations in other samples from the Schwarzwald as well as from the Vogesen, however, are lower and contain 5 to 10 ppm U. In the course of U-prospecting by routine scintillometer methods in the Alps, meta-lamprophyres have been found to have low U contents (HÜGI and LABHART, pers. comm). In meta-lamprophyres from the Alps, U concentrations are below 5 ppm, with very few exceptions from the Aarmassif, whereas the Mont Blanc samples show a higher U content (> 13 ppm). The distribution of U and its variation seems to be a very local phenomenon which can be attributed in part to metamorphic depletion.

Trace elements included in alkalifeldspars represent another class of mobile elements susceptible to metamorphic and/or metasomatic effects. The Ba-concentration, although variable in primary lamprophyres, is indeed higher in nonmetamorphic samples. Lamprophyres from the Vogesen and Schwarzwald, Mont Blanc and the Northern Aarmassif contain more than 1000 ppm of Ba, whereas the Ba concentration in meta-lamprophyres from the Southern Aarmassif, Gotthardmassif and the Ticino is below 1000 ppm (Tab. 2).

Similarly, the distribution of Sr shows a depletion of Sr due to metamorphism. Nonmetamorphic samples from Vogesen, Schwarzwald and the northern Aarmassif have Sr contents above 400 ppm (up to 660 ppm) whereas

samples from the Gotthardmassif and the Ticino show mean values of Sr of 200 ppm and 150 ppm respectively. Rb seems not to be redistributed, showing comparable concentrations (ca. 200–240 ppm) over the investigated areas. U, Sr and Ba are the only elements that show the effect of metamorphic overprinting; all others vary randomly (Tab. 2). STEINER (1984) discusses the geochemical small scale behavior near the margins of lamprophyric dikes under metamorphic conditions.

Keeping in mind that lamprophyres show considerable syn- to post-magmatic autohydrothermal alteration which results in great chemical variations, the overprint of regional metamorphism is not chemically discernable in meta-lamprophyres from the Central Alps. The regional metamorphism of lamprophyres may thus tentatively be considered a largely isochemical process.

MAJOR AND TRACE ELEMENTS

For reasons outlined above a specific discussion of each meta-lamprophyre type is not possible. Chemically semilamprophyres and anchibasaltic rocks do differ significantly from lamprophyres. Semilamprophyres are higher in SiO_2 (Fig. 5), lower in FeO, MgO, TiO_2 , MnO and CaO (Tab. 2) compared to meta-




Fig. 5 Potassium silica diagram showing the separation of the semilamprophyres from lamprophyres and meta-lamprophyres. C: calc-alkalic, K: high-K calc-alkalic, S: shoshonitic, U: ultrapotassic.

lamprophyres. The main compatible trace elements Cr, Ni and Co, as well as most of the incompatible trace elements Zr, Sr and Ba, are also lower than in meta-lamprophyres. Whereas potassium values are comparable, sodium is enriched compared to the meta-lamprophyres. The anchibasaltic rock group is higher in CaO and Na₂O and significantly lower in K₂O concentration compared to the meta-lamprophyres. The porphyrite group does not plot as a separate rock type in the chemical discrimination diagrams. Distinction between meta-lamprophyres and porphyrite is made only on the basis of mineralogical aspects i.e. microscopically and/or macroscopically visible matrix feldspar. The meta-lamprophyres range from calc-alkaline to shoshonitic and ultrapotassic compositions. Similar to dike rocks from the northwestern Alps in Italy (VENTURELLI et al., 1984), these ultrapotassic rocks are lower in Al and higher in P, Zr, Rb, K, Th and U than the calc-alkaline and shoshonitic rocks.

Fig. 6 shows a high Ni/MgO ratio and indicates a distribution corresponding to a trend line of fractional crystallization of olivine from a primitive picritic




Fig. 6 Ni vs MgO diagram showing a line of olivine fractionation after HART and DAVIS (1978) starting from a primitive melt with 12% MgO. Ni/Mg ratios (Ni ppm/MgO %) vary between 5 and 35, they are generally higher in ultrapotassic rocks.

tic basalt with approximately 12 wt.% MgO (HART and DAVIS, 1978). These authors showed that it is possible to produce such picritic basalt melts by some 5% partial melting of a model mantle peridotite composed of some 79% ol, 20% opx and 10% cpx. The high Cr, Ni, MgO values of the meta-lamprophyres investigated here and their high mg-number also point to an origin by partial melting of a primitive mantle.

The Th/Nb values are generally high (> 0.5) and are thus comparable to values of lavas from active margins (Th/Nb > 0.2; BAILEY, 1981). The enrichment of LREE (Fig. 8) and Zr are comparable to within-plate patterns.

Normalization of calc-alkalic, shoshonitic as well as ultrapotassic meta-lamprophyres against a N-type MORB (WOOD et al., 1979; sample 409-2) show bossed patterns (Fig. 7) similar to those of calc-alkalic volcanic arc basalts, showing enrichment of Sr, K, Rb and Ba. The enrichment of Ce is typical for




Fig. 7 Spider diagram for MORB normalized trace elements (MORB after WOOD et al., 1979). The patterns display an enrichment of mobile elements due to partial melt of primitive material as well as Nb, P, Ti anomalies indicating subduction processes. For explanation see text.

high-K calc-alkalic rocks (PEARCE, 1983). The weaker enrichment of Nb, P and Ti suggests an influence of subduction processes (PEARCE, *op. cit.*). In figure 7 patterns of lamprophyres and meta-lamprophyres are very similar and again no effect of metamorphism can be detected.

RARE EARTH ELEMENTS

Rare earth element (REE) contents are given in table 3. Chondrite-normalized REE distribution (SUN and NESBITT 1977, 1978) of lamprophyres and meta-lamprophyres show a pattern (Fig. 8) where light REE are enriched by factors of 60 to 300. No Eu anomaly can be detected. Heavy REE and to some extent also light REE produce gently inclined plateaus. The patterns follow a general alkalibasaltic trend (CHAUVEL and JAHN, 1984). $(La/Yb)_N$ values are low and vary from 4 to 20.

For comparison, REE patterns of lamprophyres compiled by ROCK (1984) are plotted together with data from the Schwarzwald (MÜLLER, 1982, 1984) and the Alps (Fig. 9). These lamprophyres show steeper patterns with LEE enrichment varying from 300 to 1000 for La and HREE depletion compared to the lamprophyres from the Schwarzwald and the Alps. $(La/Yb)_N$ values are higher and vary between 70 and 90. The higher HREE in lamprophyres from the Schwarzwald and meta-lamprophyres from the Alps may correlate with the very elevated amphibole content. Amphibole is the only mineral with high HREE concentrations occurring in these rocks. Garnet, another mineral with HREE enrichment is not a primary phase in lamprophyre.




Fig. 8 Chondrite normalized REE distribution pattern showing an enrichment of LREE of 60 to 300 times. Two "plateaus" are visible for LREE and HREE. The HREE plateau possibly is a reflection of the very high amphibole content of Alpine lamprophyres and meta-lamprophyres.




Fig. 9 Chondrite normalized REE distribution pattern of the Alps compared with data from the Schwarzwald (MÜLLER, 1982) and data given in a compilation by ROCK (1984). The Schwarzwald as well as the Alpine samples are less enriched in LREE and show higher HREE contents than lamprophyres from other localities.

Magmatic evolution

Since REE data of lamprophyres from the Schwarzwald and the Central Swiss Alps differ considerably from published data of lamprophyres elsewhere (Fig. 9), some ideas on the genesis of these calc-alkalic lamprophyres from the Hercynian orogenic cycle are presented here.

Trace element distributions (Fig. 7) show that a considerable mantle component as well as variable portions of a second component, probably subduction-related, might be involved in the generation of lamprophyric magmas. The MgO content indicates a small amount of partial melt released from a peridotite (AHERN and TURCOTTE, 1979; THOMPSON et al., 1984) and the enrichment in Cr and Ni can be assigned to the portion of partial peridotite melt. The general trace element enrichment fits well with data reported from alkalic xenoliths with Cr-spinell and high Ni content in olivine. As pointed out by VENTURELLI et al. (1984), however partial melting of primordial mantle cannot account for all of the trace element ratios. Admixture of other source-material is required.

Petrological investigations of the meta-lamprophyres from the Central Swiss Alps allowed to deduce pressures of 20 to 30 kb and temperatures between 950 and 1150 °C from the magmatic relics (OBERHÄNSLI, 1985 and in prep.). These physical conditions represent a situation in the upper mantle at 60 to 100 km depth.

The model presented below is based on the following petrogenetic considerations:

- Below the site of production of calc-alkalic magmas partial melting of portions of the upper lithospheric mantle wedge due to a possible partial melt of deeply subducted (> 100 km) oceanic crust seems possible (WYLLIE and SEKINE, 1982; SEKINE and WYLLIE, 1982; PEARCE, 1983).
- The composition of a melt produced from the subducted crust might be similar to that of a rhyodacite (NICHOLLS and RINGWOOD, 1973).
- Mixing of such a siliceous melt with the newly produced partial melts of an overlying mantle peridotite can lead to local concentrations of potassium and separations of the resulting magma (SEKINE and WYLLIE, 1982).
- Portions of these alkalic melts could remain at the bottom of the continental lithosphere until they are activated by tectonic or thermal events (WYLLIE and SEKINE, 1982).

The geochemical data discussed here and further geological data presented elsewhere (OBERHÄNSLI, 1985 and in prep.) are consistent with the following genetic model of Hercynian lamprophyric magmas beneath the Central Alps.

Lamprophyres, according to this model, formed by complex processes of melting of oceanic crust at great depth (> 100 km) and in different proportions, producing acid, rhyodacitic melts of variable compositions. These diverse products provoke partial melting in the overlying upper mantle. Variable percentages of melting of this earlier metasomatized mantle (metasomatized above the down going slab by fluids due the dehydration of the slab) produce partial mantle melts of varying compositions. Mixing of the two melts and the possibility that potassium enriched calc-alkalic melts could stagnate at the bottom of the continental lithosphere provides a clue to the diversity of lamprophyres e.g. minettes, vogesites, spessartites and kersantites.

The production and/or resorption of biotite and/or hornblende in lamprophyric magmas is controlled by fluid content, fluid and total pressure as well as temperature.

Tectonic activity rather than changes in the thermal pattern are thought to have led to the emplacement of lamprophyric dikes. Continued thrusting, isostatic uplift and/or volume reduction by cooling (shrinking) of the earlier intruded calc-alkalic batholiths had the potential to produce deep fault systems through which the lamprophyric magmas intruded. Thrustlike faults and compressional stresses at shallow crustal levels, due to concave upward flexures formed in response to denudation and isostatic uplift, while at the same time deeper crustal levels, according to SPENCER (1985), lay in a large scale extensional stress field.

Tab. 2 Bulk rock analyses of lamprophyres, meta-lamprophyres, semilamprophyres, porphyrites, anchibasalts and gneisses from Vogesen (V), Schwarzwald (S), Aar- (A), Mont Blanc- (M) and Gotthardmassifs (G) and Ticino (T) and petrographic sample descriptions.

Sample Nr.	V 1	V 2	V 3	V 4	V 5	V 6	V 7	V 8	V 9	V 10	V 11	V 12	V 13	V 14
Weight %														
SiO ₂	45.31	44.48	52.31	55.98	58.78	53.74	47.06	48.64	53.49	57.98	52.63	45.59	63.61	57.80
TiO ₂	1.29	1.27	1.40	1.15	0.78	0.98	1.64	1.65	1.22	0.78	1.47	1.36	0.71	1.15
Al ₂ O ₃	12.61	12.02	14.16	16.99	15.26	18.76	13.71	13.89	14.52	15.16	15.35	13.63	14.48	17.54
Fe ₂ O ₃	2.79	2.60	2.48	2.23	1.51	4.60	4.32	5.13	3.54	1.75	2.19	4.28	2.20	2.12
FeO	3.63	3.85	3.77	4.16	3.75	4.06	3.54	2.67	3.36	3.67	5.19	2.67	1.82	3.58
MnO	0.17	0.21	0.16	0.10	0.12	0.19	0.17	0.16	0.11	0.12	0.09	0.33	0.06	0.08
MgO	7.30	8.50	5.46	3.98	5.54	3.89	6.84	6.29	7.52	5.64	8.45	7.30	2.64	4.17
CaO	7.94	8.44	6.16	3.84	4.92	5.69	6.74	7.16	6.13	5.25	2.04	6.83	2.84	1.94
Na ₂ O	1.70	2.08	3.82	3.08	3.16	5.03	2.65	2.63	2.86	3.23	1.51	2.18	3.22	3.52
K ₂ O	5.90	3.82	3.11	3.92	3.10	0.71	3.99	4.06	3.09	3.31	4.99	5.17	5.07	4.19
P ₂ O ₅	2.32	2.42	0.94	0.33	0.28	0.19	0.78	0.80	0.54	0.26	0.56	1.16	0.41	0.33
P ₂ O ₄	1.35	2.10	2.29	2.18	2.39	1.90	2.78	3.04	2.50	2.55	3.51	1.75	0.88	3.14
CO ₂	7.43	7.93	3.10	0.08	0.16	0.14	4.35	4.54	0.26	0.13	1.09	7.66	0.08	0.04
Total	99.57	99.72	99.16	98.02	99.75	99.88	98.57	100.66	99.14	99.83	99.07	99.91	98.02	99.60
ppm :														
Ba	6621	4964	2167	1287	506	302	2025	1572	806	493	1634	3418	2227	-
Rb	140	96	75	279	139	37	132	148	116	133	302	179	201	-
Sr	1556	1204	998	571	389	493	426	511	691	365	269	850	996	-
Pb	62	18	7	29	8	21	24	29	<6	<6	9	64	71	-
Th	46	37	15	6	<5	<5	<5	9	7	<5	10	24	24	-
U	5	10	4	<1	3	6	<1	6	10	<1	4	6	9	-
Nb	86	82	29	7	19	7	23	25	25	12	15	34	17	-
La	413	330	173	76	18	<15	130	149	61	<15	161	265	108	-
Ce	579	507	271	148	77	67	216	216	143	69	199	364	190	-
Nd	146	146	85	45	27	19	74	78	47	15	74	122	44	-
Y	36	30	26	35	26	37	28	37	32	21	32	34	34	-
Zr	1016	937	382	235	176	147	387	394	238	166	325	543	392	-
V	166	148	151	121	102	119	210	232	150	84	136	197	96	-
Cr	320	336	174	19	247	11	380	383	396	227	292	398	78	-
Ni	175	216	113	16	117	18	236	212	197	107	89	187	36	-
Co	36	35	38	12	20	13	36	46	42	17	23	48	23	-
Cu	127	95	28	36	42	7	42	19	15	40	6	24	44	-
Zn	201	293	114	74	90	118	163	181	81	82	116	189	61	-
Ga	18	16	20	19	16	23	21	21	21	14	19	19	20	-
Sc	21	18	14	20	14	18	21	23	21	12	29	24	15	-
Total	11770	9518	4884	3035	2036	1463	4574	4291	3099	1857	3744	6989	4686	0

Tab. 2 (continued).

Sample Nr.	S 1	S 2	S 3	S 4	S 5	S 6	S 7	S 8	S 9	S 10	S 11	S 12	S 13	S 14	S 15	S 16
Weight %																
SiO ₂	45.99	53.25	54.43	52.47	59.85	58.05	54.47	58.85	60.21	53.48	49.80	60.13	62.25	52.42	59.72	49.55
TiO ₂	1.32	0.98	0.84	1.11	1.45	1.48	1.16	1.03	0.97	1.33	1.40	1.04	1.44	1.09	1.03	0.79
Al ₂ O ₃	15.74	14.26	13.66	14.50	13.78	13.75	13.97	13.98	15.94	13.64	14.20	16.34	15.10	13.20	14.99	12.61
Fe ₂ O ₃	8.72	2.08	2.31	1.39	2.59	2.89	1.67	2.54	1.82	2.69	2.67	2.92	3.61	2.37	1.80	2.41
FeO	2.34	4.35	4.64	5.43	3.49	3.75	4.25	2.91	3.19	3.73	2.82	2.35	2.32	4.35	3.77	4.67
MnO	0.08	0.12	0.12	0.12	0.06	0.07	0.10	0.09	0.11	0.10	0.42	0.08	0.06	0.13	0.10	0.13
MgO	7.45	8.17	9.72	8.09	6.68	6.72	8.79	6.05	3.61	8.04	6.28	3.86	4.07	9.00	4.50	10.68
CaO	3.33	4.29	6.16	5.71	1.58	1.55	3.62	3.07	4.28	3.07	5.76	2.41	1.58	5.23	2.37	5.80
Na ₂ O	0.22	2.30	1.96	2.91	1.51	1.22	1.05	2.51	3.17	1.75	1.36	2.70	2.44	1.58	1.79	1.71
K ₂ O	6.63	3.80	2.95	3.06	5.98	5.97	4.69	4.97	3.68	5.27	6.49	5.48	4.67	3.95	4.23	3.31
P ₂ O ₅	1.07	0.62	0.49	0.55	0.88	0.91	0.73	0.97	0.45	1.08	1.16	0.59	0.60	0.80	1.29	0.67
P ₂ O ₄	3.82	2.58	1.67	3.24	2.47	2.82	4.17	2.17	1.35	2.56	2.15	1.63	1.87	2.77	3.05	4.86
CO ₂	1.32	1.58	0.18	0.75	0.11	0.11	2.15	0.15	1.06	1.07	2.91	0.05	0.05	2.18	0.40	2.20
Total	98.03	98.38	99.13	99.33	100.43	99.29	100.82	99.29	99.84	97.81	97.42	99.58	100.06	99.07	99.04	99.39
Ppm :																
Ba	5982	2126	1562	1002	2491	1927	2324	2139	1247	1911	1966	1553	1171	1716	1339	3040
Rb	197	165	128	141	187	204	212	187	136	341	401	295	289	298	325	137
Sr	247	357	391	304	506	481	282	374	690	284	359	443	177	316	116	447
Pb	43	18	15	11	23	20	9	32	12	19	27	27	15	97	13	24
Th	6	15	7	16	24	23	10	14	11	41	43	30	19	41	36	27
U	<1	7	4	8	7	7	11	9	4	9	20	14	9	20	21	10
Nb	13	17	14	16	29	27	22	29	11	39	41	25	24	28	25	16
La	220	66	19	40	161	131	93	90	69	72	92	100	99	73	73	51
Ce	282	125	106	115	189	147	141	102	145	110	144	242	100	239	129	59
Nd	68	34	33	47	56	51	42	28	49	39	51	98	28	109	31	31
Y	37	36	35	33	36	40	39	48	32	44	51	39	52	40	41	36
Zr	311	251	229	264	576	569	283	350	273	406	428	326	401	293	274	180
V	223	176	169	144	280	196	173	177	136	219	227	145	198	194	182	183
Cr	447	626	883	474	586	496	745	340	798	557	798	296	444	1021	123	890
Ni	325	179	268	152	81	78	262	95	34	133	146	106	114	273	50	166
Co	39	46	43	31	42	18	47	35	24	46	39	30	40	49	49	36
Cu	27	33	33	35	23	16	47	25	41	25	21	33	30	52	79	13
Zn	448	83	104	78	125	113	200	84	108	189	121	83	138	236	105	269
Ga	25	18	17	16	19	19	18	18	17	18	17	20	20	19	19	13
Sc	29	29	28	22	32	31	28	26	19	34	36	24	35	36	28	33
Total	8969	4407	4088	2949	5473	4594	4988	4202	3163	4536	5028	3929	3403	5150	3058	5661

Tab. 2 (continued).

Sample Nr.	A 2	A 3	A 4	A 5	A 6	A 7	A 8	A 9	A 10	A 11	A 12	A 13	A 14	A 15	A 16	A 17
Weight %																
SiO ₂	53.28	49.27	57.97	55.12	45.71	46.57	47.63	53.34	56.42	57.72	57.33	54.41	51.72	48.33	53.60	53.50
TiO ₂	1.09	1.73	0.75	1.46	1.20	1.48	1.16	1.13	0.92	0.91	0.91	0.81	1.39	0.99	1.00	1.10
Al ₂ O ₃	15.35	15.97	17.08	17.08	15.29	14.66	14.99	17.02	16.19	16.17	17.36	16.82	15.47	12.48	12.80	14.10
Fe ₂ O ₃	2.97	2.89	2.54	4.64	2.80	2.84	1.00	3.24	3.16	1.79	3.27	4.31	2.81	1.91	1.20	1.30
FeO	4.09	7.14	2.62	3.78	4.46	3.84	6.30	4.53	2.82	4.09	3.49	2.73	4.83	5.35	5.20	4.50
MnO	0.12	0.17	0.11	0.15	0.14	0.12	0.16	0.13	0.12	0.13	0.13	0.14	0.25	0.13	0.11	0.18
MgO	8.71	6.38	3.28	3.19	6.56	7.98	7.61	5.14	4.78	6.00	3.87	5.27	8.03	8.84	8.40	7.40
CaO	3.55	7.67	5.88	7.12	8.20	8.27	8.46	7.46	5.55	3.35	5.86	7.36	4.57	6.59	7.20	8.40
Na ₂ O	2.59	3.64	2.80	3.23	2.31	1.66	2.85	3.11	2.93	3.35	1.19	1.98	2.16	1.46	3.10	2.90
K ₂ O	4.21	1.64	3.22	1.44	3.64	5.05	2.90	1.69	3.55	4.52	4.44	3.63	5.47	5.18	4.20	2.70
P ₂ O ₅	0.34	0.58	0.20	0.34	1.22	1.11	1.17	0.30	0.38	0.36	0.23	0.19	0.56	1.10	1.40	1.40
P ₂ O ₄	3.44	2.91	1.95	2.54	3.61	2.79	3.40	2.84	1.55	1.84	1.92	1.67	1.45	2.48	1.00	2.00
CO ₂	0.13	0.55	0.17	0.13	3.81	2.23	1.94	0.06	0.03	0.07	0.03	0.05	0.82	3.64	0.10	0.65
Total	99.87	100.54	98.97	100.22	98.95	98.70	99.57	99.99	98.40	100.30	100.03	99.37	99.53	98.48	99.31	100.13
ppm :																
Ba	608	390	826	686	2030	2222	1653	437	898	737	1029	807	517	3610	5000	3000
Rb	93	100	253	71	159	185	83	85	152	237	238	296	522	394	150	70
Sr	680	642	862	470	887	646	1068	743	747	457	491	764	454	983	850	1475
Pb	<6	9	<6	11	22	12	9	25	<6	7	22	25	37	42	18	36
Th	<5	<5	<5	<5	<5	10	22	<5	<5	5	<5	<5	11	16	-	-
U	<1	3	<1	4	4	3	5	<1	<1	2	3	3	10	2	-	-
Nb	9	15	<3	13	55	74	49	8	10	12	12	6	20	42	-	-
La	47	66	<15	33	128	163	118	<15	<15	42	19	<15	43	128	-	-
Ce	144	123	103	159	287	312	241	147	108	113	149	131	151	215	-	-
Nd	68	73	53	84	110	123	97	73	23	34	71	71	78	75	-	-
Y	29	37	20	36	39	33	32	33	28	31	33	27	30	41	-	-
Zr	204	223	94	185	343	482	315	155	186	188	156	116	219	281	355	495
V	124	146	110	223	168	152	161	184	107	131	139	142	82	191	-	-
Cr	273	102	32	31	212	275	212	78	104	118	58	145	359	717	470	480
Ni	121	51	24	18	90	141	107	23	38	41	34	135	176	161	130	95
Co	28	39	17	14	25	34	37	17	15	26	26	26	27	42	21	35
Cu	32	36	17	35	58	51	53	25	8	8	14	17	16	55	42	39
Zn	87	107	100	117	95	82	78	105	88	120	66	128	219	108	80	90
Ga	17	15	19	20	16	17	15	19	17	18	20	19	23	17	-	-
Sc	22	23	13	26	27	21	20	28	10	12	16	19	17	35	-	-
Total	2586	2200	2543	2236	4755	5038	4375	2185	2539	2339	2596	2777	3011	7155	7116	5815

Tab. 2 (continued).

Sample Nr.	A 18	A 19	A 20	A 21	A 22	A 23	A 24	A 25	A 26	A 27	A 29	A 30	A 31	A 32	A 33	A 34
SiO ₂	54.00	53.60	53.10	52.50	53.20	54.40	49.40	50.10	49.80	53.00	49.80	51.88	58.37	56.67	59.85	55.03
TiO ₂	0.88	1.50	0.92	0.71	0.62	1.40	1.80	0.93	1.80	1.11	1.24	1.00	0.84	1.01	0.86	1.21
Al ₂ O ₃	16.50	12.60	16.20	15.30	14.70	17.70	16.50	14.80	16.20	17.34	16.17	15.94	15.26	16.21	14.99	17.58
Fe ₂ O ₃	1.70	1.50	2.00	1.30	2.10	2.10	1.80	2.60	2.00	5.10	4.75	3.49	1.96	2.53	2.27	1.89
FeO	5.00	4.80	5.50	6.00	5.10	5.00	7.40	5.90	7.40	2.95	2.40	3.41	3.99	4.95	4.21	5.84
MnO	0.09	0.08	0.14	0.12	0.13	0.13	0.19	0.20	0.22	0.13	0.14	0.14	0.15	0.14	0.15	0.16
MgO	7.80	8.40	8.60	10.20	9.30	4.50	8.40	10.20	7.70	5.96	4.39	8.90	5.67	6.11	5.50	5.25
CaO	5.20	5.60	5.50	5.10	7.80	5.40	5.30	7.90	5.80	8.89	6.72	6.54	3.55	4.33	3.09	6.19
Na ₂ O	1.90	2.00	2.60	3.00	2.00	4.40	3.50	2.60	3.80	2.74	5.38	0.79	3.94	1.58	2.75	2.27
K ₂ O	3.30	7.50	2.70	2.10	1.90	2.20	1.60	1.80	1.60	1.18	1.58	5.12	3.78	4.31	4.19	3.69
P ₂ O ₅	0.24	0.92	0.24	0.22	0.20	0.32	0.50	0.23	0.49	0.22	0.65	0.24	0.32	0.27	0.35	0.30
H ₂ O ⁺	2.90	1.60	2.80	3.50	3.30	1.80	3.60	2.80	2.70	2.71	2.22	3.24	2.02	3.03	1.73	1.45
CO ₂	0.11	0.30	0.15	0.30	0.66	0.10	0.10	0.76	0.10	0.12	3.65	0.41	0.07	0.04	0.46	0.06
Total	99.62	100.40	100.45	100.35	101.01	99.45	100.09	100.82	99.61	101.45	99.09	101.10	99.92	101.18	100.40	100.92

ppm :

Ba	2575	1700	1025	370	240	550	950	360	560	231	240	1348	378	1211	375	568
Rb	190	270	175	165	110	150	100	135	105	65	76	249	264	202	299	214
Sr	780	620	600	320	340	680	710	360	360	809	312	279	99	205	322	388
Pb	16	19	15	12	10	38	8	12	22	9	<6	6	<6	7	7	18
Th	-	41	-	-	-	-	-	-	-	<5	<5	<5	<5	<5	5	<5
U	-	-	-	-	-	-	-	-	-	4	1	5	<1	1	4	5
Nb	-	-	-	-	-	-	-	-	-	8	14	9	12	16	17	13
La	-	116	-	-	-	-	-	-	-	<15	<15	<15	27	56	39	39
Ce	-	220	-	-	-	-	-	-	-	58	181	163	145	165	99	162
Nd	-	-	-	-	-	-	-	-	-	28	99	75	74	77	34	89
Y	-	-	-	-	-	-	-	-	-	30	37	26	30	34	33	41
Zr	190	580	170	105	110	200	240	110	200	102	242	137	192	194	195	185
Y	-	-	-	-	-	-	-	-	-	212	82	115	122	181	171	191
Cr	300	360	350	640	550	85	190	710	180	115	131	533	247	313	270	113
Ni	95	185	85	250	215	10	100	195	105	26	55	102	59	102	59	53
Co	46	38	48	63	52	39	70	64	72	28	29	36	28	27	23	30
Cu	35	40	24	37	31	22	27	62	20	24	36	19	9	49	8	16
Zn	70	80	85	95	55	95	240	70	140	77	77	87	95	197	104	154
Ga	-	-	-	-	-	-	-	-	-	20	18	22	18	20	25	22
Sc	-	25	-	-	-	-	-	-	-	27	16	23	20	27	22	24
Total	4297	4294	2577	2057	1713	1869	2635	2078	1764	1873	1646	3234	1819	3084	2111	2325

Tab. 2 (continued).

Sample Nr.	G 1	G 2	G 3	G 4	G 5	G 6	G 7	G 8	G 9	G 10	G 11	G 12	G 13	G 14
Weight %														
SiO ₂	65.78	51.59	53.71	62.98	55.27	54.72	52.95	55.70	53.82	50.46	60.46	55.17	49.23	49.23
TiO ₂	0.80	1.14	1.19	1.05	1.08	0.94	1.13	1.60	1.06	0.83	1.06	0.80	0.47	0.82
Al ₂ O ₃	15.19	16.29	15.63	15.79	14.78	16.07	16.74	15.75	16.11	14.98	15.98	16.42	15.02	15.26
Fe ₂ O ₃	1.10	1.95	2.76	2.08	1.80	3.03	2.50	1.98	2.44	3.56	2.89	2.30	1.71	2.42
FeO	4.58	6.48	7.88	3.70	8.21	4.32	5.55	5.20	4.47	3.19	2.80	3.79	2.55	5.59
MnO	0.09	0.16	0.16	0.10	0.17	0.15	0.14	0.12	0.12	0.15	0.11	0.13	0.12	0.15
MgO	2.70	6.87	5.70	1.83	5.34	5.71	5.63	3.43	4.90	7.06	1.67	3.97	3.82	9.22
CaO	1.91	7.28	2.38	3.26	2.90	6.35	4.21	4.86	6.32	6.10	4.99	4.70	2.88	7.26
Na ₂ O	2.45	2.40	1.86	4.15	1.76	0.18	3.72	2.30	4.09	0.55	3.66	2.13	3.09	0.22
K ₂ O	3.21	2.87	5.48	3.16	4.99	6.80	4.06	5.23	1.93	6.49	3.69	5.15	2.81	5.54
P ₂ O ₅	0.22	0.27	0.29	0.36	0.27	0.21	0.34	0.76	0.34	0.19	0.38	0.30	0.22	0.16
H ₂ O ⁺	1.90	1.72	2.03	1.00	1.95	1.50	1.55	2.03	1.10	2.36	1.07	2.31	1.74	3.06
CO ₂	0.11	0.32	0.67	0.07	1.32	0.58	1.80	1.49	2.87	3.52	0.65	3.17	1.12	0.80
Total	100.04	99.34	99.74	99.53	99.84	100.56	100.32	100.45	99.57	99.44	99.41	100.34	99.79	99.73
ppm :														
Ba	647	405	420	848	467	735	578	891	430	450	897	643	190	570
Rb	148	133	278	134	254	331	179	209	98	341	225	240	297	268
Sr	242	509	174	217	177	73	109	283	245	70	303	182	177	73
Pb	14	6	11	<6	10	24	9	17	6	6	19	634	14	7
Th	12	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5	9	<5	<5
U	<1	<1	<1	<1	<1	4	1	1	1	<1	<1	<1	10	<1
Nb	11	8	11	19	11	12	11	37	11	9	25	8	38	8
La	51	<15	71	25	86	20	66	119	<15	15	42	26	<15	<15
Ce	78	81	93	101	115	84	82	178	99	111	192	91	21	129
Nd	46	29	49	35	61	26	49	72	34	61	97	35	<10	75
Y	31	32	31	38	33	33	29	51	30	34	43	26	21	28
Zr	191	142	150	366	136	135	186	454	187	117	428	163	89	117
V	97	182	182	73	216	161	106	164	105	138	110	145	85	190
Cr	79	231	223	19	223	196	101	85	103	226	30	36	204	479
Ni	42	37	36	18	37	46	34	44	39	42	26	20	38	129
Co	26	33	<8	<8	8	31	28	25	17	12	<8	13	9	30
Cu	20	9	10	6	11	8	11	<4	9	15	23	11	19	15
Zn	108	92	96	60	94	107	122	116	104	100	114	111	69	95
Ga	20	20	27	18	28	17	19	21	19	18	23	19	19	18
Sc	15	20	35	11	38	16	16	19	13	27	14	18	12	33
Total	1878	1969	1897	1988	2005	2059	1736	2786	1550	1792	2611	2430	1312	2264

Tab. 2 (continued).

Sample Nr.	T 1	T 2	T 3	T 4	T 5	T 6	T 7	T 8	T 9	T 10	T 11	T 12	T 13	T 14	T 15	T 16	T 20
Weight %																	
SiO ₂	50.08	61.40	62.21	61.81	64.55	57.16	65.04	65.37	67.14	55.79	68.21	64.77	70.97	64.25	63.83	71.12	54.40
TiO ₂	0.96	0.60	0.63	0.63	0.53	0.72	0.53	0.50	0.71	0.78	0.84	0.52	0.31	0.65	0.65	0.75	1.20
Al ₂ O ₃	14.29	18.20	18.23	18.00	16.33	19.93	16.47	16.33	15.13	15.47	14.95	15.62	15.17	16.87	17.00	13.27	16.60
Fe ₂ O ₃	2.87	0.56	0.87	0.30	0.96	1.62	1.22	1.31	1.07	0.62	0.87	1.61	0.68	0.98	1.18	4.83	1.20
FeO	5.28	3.63	3.01	3.51	2.43	4.04	2.38	2.18	3.77	7.06	3.88	1.81	1.01	3.14	2.97	0.00	6.30
MnO	0.17	0.10	0.08	0.08	0.07	0.11	0.08	0.08	0.10	0.20	0.10	0.10	0.12	0.10	0.07	0.08	0.10
MgO	9.85	1.57	1.23	1.18	1.48	1.90	1.44	1.65	2.01	8.06	1.98	2.88	0.78	1.91	1.94	1.26	7.10
CaO	6.85	2.88	1.85	1.81	3.28	3.37	3.84	2.55	1.60	4.78	2.13	2.28	1.24	3.94	4.11	1.93	4.55
Na ₂ O	0.50	4.44	6.81	6.86	3.67	4.44	3.11	1.71	2.29	0.32	2.90	6.70	5.06	3.09	3.14	3.29	1.10
K ₂ O	5.15	4.54	2.77	2.76	4.10	3.90	4.28	6.51	4.17	4.43	2.31	2.03	3.39	3.51	3.17	2.67	4.70
P ₂ O ₅	0.28	0.20	0.22	0.21	0.19	0.20	0.18	0.17	0.24	0.14	0.25	0.19	0.12	0.15	0.15	0.29	0.30
H ₂ O ⁺	2.04	0.73	0.82	1.85	0.80	1.26	0.79	1.09	1.38	1.56	1.25	0.84	0.58	0.88	0.87	0.00	1.80
CO ₂	1.52	0.19	0.32	0.15	0.48	0.24	0.36	0.16	0.09	0.05	0.09	1.01	0.08	0.05	0.06	0.00	0.10
Total	99.84	99.04	99.05	99.15	98.87	98.89	99.72	99.61	99.70	99.26	99.84	100.36	99.51	99.52	99.14	99.49	99.45
ppm :																	
Ba	610	300	136	365	411	330	327	376	666	324	1103	155	452	456	554	-	307
Rb	224	311	199	329	234	257	266	344	185	273	80	115	96	180	174	-	488
Sr	131	125	94	91	218	178	119	86	166	128	206	202	187	207	296	-	39
Pb	14	13	9	9	13	10	9	9	9	9	16	9	9	11	<6	-	12
Th	<5	6	<5	<5	<5	6	<5	7	<5	<5	9	<5	<5	<5	<5	-	21
U	2	<1	<1	<1	<1	3	1	<1	<1	<1	3	3	1	<1	<1	-	13
Nb	12	16	11	4	7	29	8	5	11	10	13	7	4	10	11	-	29
La	36	30	27	20	17	41	<15	26	42	44	50	<15	<15	21	15	-	57
Ce	168	43	41	41	43	80	44	45	78	83	77	41	42	44	43	-	61
Nd	90	26	25	<10	26	26	<10	27	26	27	46	25	<10	26	26	-	<10
Y	33	29	32	11	20	49	24	21	27	28	27	19	14	19	18	-	60
Zr	147	160	168	112	146	159	143	147	176	119	228	146	129	114	118	-	207
V	158	69	36	63	54	75	44	61	92	171	97	38	21	69	88	-	333
Cr	598	16	11	16	11	16	12	<10	68	444	89	11	<10	16	21	-	268
Ni	176	20	20	20	20	20	20	20	30	161	34	20	20	20	<3	-	112
Co	39	12	8	14	12	15	12	19	23	44	22	13	13	21	21	-	35
Cu	13	60	19	11	21	50	14	9	33	9	109	15	18	13	23	-	<4
Zn	151	238	97	130	88	171	95	58	85	135	104	93	28	99	91	-	98
Ga	19	28	30	27	23	37	24	20	20	19	21	24	18	20	22	-	20
Sc	17	11	9	10	7	14	6	8	13	26	14	8	5	11	14	-	35
Total	2638	1513	972	1273	1371	1566	1168	1288	1750	2054	2348	942	1057	1357	1535	0	2195

Tab. 2 (continued).

Description of samples and localities		D: drillhole, I: tunnel		mac: macroscopically visible	
Nr.	Locality	sample description			
V 1	Wackenbach	minette, massy compact dark grey	S 12	Laufenburg	T 646.80/268.20
V 2	Wackenbach	mac: bi qz minette, massy compact grey brown	S 13	Laufenburg	I 646.80/268.20
V 3	S Rothau	mac: bi qz-oiicelli kfsp-insets minette, finegrained massy dark	S 14	Laufenburg	I 646.80/268.20
V 4	Sperberbächle	grey with salmon red points mac: qz-oiicelli	S 15	Kaisten	D 644.64/265.62
V 5	Sperberbächle	vogesite, dark massy compact grey	S 16	Leuggern	D 657.63/271.21
V 6	Eftermatten Lilsbach	vogesite with fine margin, very compact massy grey green	A 2	Madraner Tal	703.85/183.75
V 7	Andlau	mac: plg resorbed px few bi spessartite, compact massy grey	A 3	Tscharren	700.45/178.25
V 8	Andlau	mac: plg kernsantite finegrained, redish grey	A 4	Ezlhütte	794.90/174.85
V 9	Tannenber bei Saales	mac: qz-oiicelli bi resorbed px/ol minette, redish grey compact	A 5	Kleiner Mutsch	697.80/174.20
V 10	Tannenber bei Saales	mac: bi px spessartite massy compact grey green	A 6	Sustenpass	677.25/175.75
V 11	Henaumont ob Markkirch	mac: bi px qz-oiicelli lamprophyre, massy compact, grey green	A 7	Sustenpass	T 677.25/176.00
V 12	Col de la Grosse Pierre	mac: bi px qz-oiicelli kernsantite, feldspar rich medium grained greenish grey	A 8	E Sustenpasshöhe	677.80/177.05
V 13	Petites Alpes	mac: kfsp bi plg lamprophyre, finegrained dark grey	A 9	Trift	672.18/170.17
V 14	Col de la Grosse Pierre	mac: kfsp bi lamprophyre with gniss insets, massy compact grey brown to redish	A 10	Göschenen	688.20/170.65
S 1	Todmoos	minette, dark grey homogeneous	A 11	Göschenen	688.20/170.65
S 2	Lochmühle	mac: bi pyx minette, fine to medium grained	A 12	Schöllenen	678.75/167.25
S 3	Murgtal	mac: bi resorbed px/ol lamprophyre, fine to medium grained	A 13	Urnerloch	688.25/166.60
S 4	Tiefenstein	mac: bi plg kfsp (porphyrite) lamprophyre, massy greenish grey	A 14	Urnerloch	688.25/166.60
S 5	Albtal	mac: bi qz-oiicelli kernsantite, compact massy brownish grey	A 15	Vättis, Kreuzbach	752.50/198.10
S 6	Albtal	mac: bi qz-oiicelli lamprophyre, fine to medium grained grey, compact	A 16	Punteglias	716.06/182.80
S 7	Albtal	mac: bi kfsp resorbed cpx/ol lamprophyre, with chilled margin, black finegrained compact massy	A 17	Punteglias	716.03/182.76
S 8	Bantlisboden	mac: bi kfsp matrix, dark red brown spherulitic parts, bi overgrows, spherulites	A 18	Punteglias	715.26/180.97
S 9	Laufenburg	mac: bi, kfsp cpx/ol pseudomorphs lamprophyre compact, greenish grey	A 19	Punteglias	715.35/182.35
S 10	Laufenburg	mac: bi kfsp lamprophyre, dark grey compact redish grey, masy xenoliths of greenstones	A 20	Punteglias	715.31/182.51
S 11	Laufenburg	mac: bi kfsp lamprophyre, massy compact dark grey, fluidal textures	A 21	Punteglias	715.06/182.58
		mac: qz-oiicelli lamprophyre, compact massy redish	A 22	Punteglias	715.22/182.61
		mac: bi, totally resorbed cpx/ol spherulitic, groundmass devitrified, bi overgrows spherulites			
		mac: bi, totally resorbed cpx/ol lamprophyre, compact massy redish			
		spherulitic, groundmass devitrified, bi overgrows, spherulites			
		mac: bi, totally resorbed cpx/ol lamprophyre, massy dark grey to brown redish spots			
		mac: bi fsp lamprophyre, dark grey green with dark green spherulites			
		mac: bi chl plg kernsantite, compact greenish black			
		mac: bi chl fsp minette			
		mac: bi px/ol pseudomorphs minette			
		mac: bi px/ol pseudomorphs minette			
		mac: bi px/ol pseudomorphs minette			
		mac: chl schisc, green chlorite schist			
		mac: chl bi-pseudomorphs lamprophyric dike, finegrained green chl-bi-schist			
		mac: chl ser bi pyr ep/fsp clods lamprophyre, bi-chl-gneiss, grey white			
		mac: bi fsp lamprophyre, ep-chl-bi-gneiss, green			
		mac: ep fsp chl pyr lamprophyric dike, epidote chlorite gneiss, compact grey gniss			
		mac: bi fsp lamprophyric dike, biotite gneiss, compact grey with pygmatic fsp veins			
		mac: bi dike, chl-gneiss, granulous green grey			
		mac: plg ep chl bi chlorite gneiss, compact			
		mac: chl ep fsp chlorite biotite gneiss, brown grey			
		mac: bi fsp qz insets dike, silver grey dark sericite schist			
		mac: bi fsp ep lamprophyric dike, chlorite sericite schist, finegrained silver grey			
		mac: chl ser bi fsp lamprophyric dike, bi-chl-ser-schist, grey silvery gniss with fsp veins			
		mac: bi chl ser lamprophyric dike, bi-chl-schist, silver grey			
		mac: bi chl ser fsp-veins kernsantite, compact grey dark gneissaceous			
		mac: bi chl? qz-insets fsp? pyroxene-minette, greenish grey compact to finegrained			
		mac: bi px pseudomorphs pyroxene-kernsantite, grey green massy			
		mac: bi pyx amphibole-kernsantite, grey green massy			
		mac: bi anph pyroxene-minette, greenish grey compact, finegrained			
		mac: bi px pseudomorphs amphibole-kernsantite, grey green			
		mac: bi amph pyroxene-spessartite, grey green finegrained, compact			
		mac: hbl px pseudomorphs pyroxene-spessartite, grey green			

Tab. 2 (continued).

A 23	Punteglias	715-05/181.36	mac: hbl px pseudomorphs spessartite, grey green finegrained mac: hbl px pseudomorphs	G 4	Mätteli	686.05/160.60	mac: bi chl ser plg dike, grey black biotite gneiss mac: bi plg qz
A 24	Punteglias	715-38/181.71	mac: hbl px pseudomorphs spessartite, grey green finegrained mac: hbl px pseudomorphs	G 5	Gottthardroad	687-80/158.70	lamprophyric dike, grey to black biotite gneiss mac: bi mu fsp
A 25	Punteglias	714-98/182.19	mac: hbl px pseudomorphs spessartite, grey green finegrained mac: hbl px pseudomorphs	G 6	Gottthardroad	685.70/157.60	dike, massy pseudodioritic biotite epidote gneiss, black and white dots mac: qz bi fsp ep
A 26	Punteglias	712.13/180.54	mac: hbl px pseudomorphs spessartite, grey green finegrained mac: hbl px pseudomorphs	G 7	Gottthardroad	687.10/154.50	biotite gneiss, green to black biotite schist, silvery grey with white fsp inlets, dark bi clods mac: fsp bi chl ser
A 27	Laß Serein	705-05/176.10	mac: ep fsp bi 7amph dike, ep-bi-schist, green grey mac: bi chl cc clods	G 8	Gottthardroad	T 687.15/158.10 (5520m S)	white fsp inlets, dark bi clods mac: bi mu plg
A 29	Reichfirn	637-20/142.20	mac: bi chl cc clods lamprophyric dike, clear massy grey mac: chl ser bi fsp ep	G 9	Tremolaroad	686.60/155.80	biotite amphibole gneiss, brown grey - dike, pseudodioritic gneiss with black and white dots mac: amph fsp ser
A 30	Aletschhorn	642-70/146.00	mac: chl ser bi fsp ep lamprophyre, ser-chl-schist, phyllitic, grey, orenulated, mac: bi fsp ser	G 10	Gottthardroad	686.57/155.76	compact massy grey lamprophyric dike, mac: bi qz fsp
A 31	Oberaletschhütte	641.10/141.70	mac: chl ser bi fsp ser lamprophyric dike, compact grey gneiss mac: qz-insets pyr	G 11	Bedretto	T 679 /154 Window (5160m)	compact massy grey lamprophyric dike, mac: bi chl ser
A 32	Grubengletscher	663.00/162.80	mac: chl ser bi fsp ser lamprophyric dike, compact grey gneiss mac: qz-insets pyr	G 12	Rosbodenstock	693.10/165.50	chlorite biotite schist, grey brown mac: chl bi ser
A 33	Grimsel	669.10/158.85	mac: chl ser bi fsp-insets dike, chl-ser-schist, dark grey phyllitic mac: chl ser bi fsp-insets	G 13	Rosbodenstock	693.10/165.50	dike, clear silver grey ser-chl-schist mac: chl ser
A 34	Grimsel Egg	668-80/158.35	mac: bi ser fsp qz-olcelli dike, brown grey bi-ser-gneiss mac: bi ser fsp qz-olcelli	G 14	Lukmanierroad	707.29 /165.40	dike, clear grey green bi-amph-schist mac: chl ser bi plg
A 35	Grimsel Erlen	668-68/157.80	mac: chl ser bi? pyr qz-olcelli dike, grey green chlorite schist mac: chl ser bi? pyr qz-olcelli	T 1	Laghetta	688.40/147.39	dike, coarse chl-bio-amph-gneiss mac: bi chl ep plg
A 36	Grimsel Erlen	668-68/157.80	mac: chl plg bi dike, bi-chl schist, greyish green mac: chl plg bi	T 2	Viadotto di Monte	D 702.75/149.25	biotite gneiss mac: bi plg qz
A 37	Grimsel Hospiz	668.50/158.20	mac: chl plg ep lamprophyric dike, bi-chl-gneiss, greenish silver grey mac: chl plg ep	T 3	Viadotto di Monte	D 702.75/149.25	two mica gneiss mac: bi mu plg qz
A 38	Grimsel Hospiz	668.50/158.20	mac: chl plg ep lamprophyric dike, grey green schistose gneiss mac: bi chl ser	T 4	Viadotto di Monte	D 702.75/149.25	biotite gneiss mac: bi mu plg qz
A 39	Zuleitungsstollen Grimsel	T 668 /158 (903m)	mac: chl ser bi amph foliated lamprophyre, amph-bi-gneiss, very finegrained, strongly foliated mac: chl ser bi amph	T 5	Viadotto di Monte	D 702.75/149.25	biotite gneiss mac: bi plg qz
A 40	E Grimselsee	668-70/157.70	mac: chl bi ser lamprophyric dike, chl-bi-gneiss, greenish grey with spherulites mac: chl bi ep/fsp clods qz-olcelli	T 6	Viadotto di Monte	D 702.75/149.25	two mica gneiss mac: bi mu plg qz
A 41	Verbindungsstollen Grimsel	T 668 /158 F III	mac: chl bi ep/fsp clods qz-olcelli dark greenish grey mac: chl bi pyr ep/fsp clods	T 7	Viadotto di Monte	D 702.75/149.25	two mica gneiss mac: bi mu plg qz
A 42	N Grimselroad	669.50/157.30	mac: chl bi pyr ep/fsp clods lamprophyric dike, bi-chl-ser gneiss mac: chl ser bi	T 8	W Faido	705.70/148.10	two mica gneiss mac: bi mu plg qz
A 43	Grimselpass	669.05/157.60	mac: chl ser bi lamprophyric dike, ser-bi-chl schist, dark grey black phyllitic mac: chl bi ser	T 9	Sta. Petronella Bisoca	718.40/134.65	two mica gneiss mac: bi mu plg qz
A 44	Grimselpass	669.05/157.60	mac: chl bi ser lamprophyric dike, ser-chl-bi-schist, silvery grey to greenish mac: chl bi ser	T 10	Quarry Iragna	717.50/132.60	biotite gneiss, dark redish to black with black bi flakes mac: bi plg
M 1	Station Glacier des Pelerins	669.05/157.60	mac: chl bi ser lamprophyre, dark grey green schist mac: chl bi ser	T 11	Biascina Block	708.80/142.92	biotite gneiss mac: bi plg qz
M 2	Below Glacier des Pelerins	669.05/157.60	mac: chl bi ser lamprophyre, dark grey gneiss mac: chl bi mgt qz-olcelli	T 12	Biascina	708.80/142.92	biotite amphibole gneiss mac: bi plg amph qz
G 1	Chämleten	686.50/163.20	mac: chl bi mgt qz-olcelli biotite plagioclase gneiss mac: bi chl mu	T 13	Biascina	708.80/142.92	biotite gneiss mac: bi plg qz
G 2	Mätteli	686.15/161.25	mac: bi chl mu lamprophyre or mafic inclusion, bi- ep-gneiss, grey black, pyrogenic folds mac: bi chl qz segregations	T 14	Piotte di Bodio	713.40/137.72	biotite gneiss mac: bi plg qz
G 3	Mätteli	686.10/160.75	mac: bi chl qz segregations lamprophyric rimfacies of a quartz- porphyre, dark grey biotite gneiss with feldspar clods and veins mac: bi chl qz	T 15	Piotte di Bodio	713.40/137.72	biotite gneiss mac: bi plg qz
				T 16	Grimosina	711.10/138.65	biotite gneiss mac: bi plg qz
				T 20	Val Ambra		biotite gneiss, dark redish with biotite flakes mac: bi plg

:REE(ppm)	La	Ce	Nd	Sm	Eu	Tb	Yb	Lu
: V 4	: 43.36	: 133.12	: 63.69	: 14.61	: 2.26	: .935	: 3.23	: .28
: V 8	: 53.6	: 185.9	: 83.2	: 16.98	: 2.9	: .91	: 2.2	: .11
: S 6	: 31.48	: 78.24	: 74.4	: 11.2	: 2.28	: 1.02	: 2.77	: .31
: S 13	: 17.2	: 61.1	: 31.8	: 7.8	: 1.29	: .99	: 5.2	: .38
: A 7	: 74.6	: 218.9	: 90.8	: 16.9	: 3.2	: .93	: 2.4	: .26
: A 13	: 28.9	: 83.7	: 42.9	: 11.	: 1.7	: .69	: 1.9	: .34
: A 19	: 116.	: 220.	: -	: -	: 4.7	: -	: 2.7	: .3
: G 3	: 13.7	: 49.4	: 88.2	: 7.9	: 1.29	: .61	: 2.63	: .21
: G 11	: 30.25	: 105.7	: 46.8	: 11.3	: 2.6	: 1.2	: 3.9	: .4
: T 1	: 21.4	: 78.2	: 46.5	: 9.6	: 1.5	: .67	: 2.7	: .55
: T 11	: 11.3	: 35.7	: 18.2	: 3.6	: .75	: .41	: 1.6	: .2

Tab. 3 Rare earth element analyses from lamprophyres and meta-lamprophyres from Vogesen, Schwarzwald and Central Swiss Alps.

Acknowledgements

I am thankful to Prof. E. Niggli who has guided my interest to these complex and, in many aspects, intriguing rocks. It is a pleasure to acknowledge the help of Martin Engi, Ivan Mercolli, Urs Raz and Franz Schenker whose contributions to fruitful discussions, cohabitation during field-work and assistance in work is invaluable to me. Prof. Tj. Peters and J. Meyer provided data on the NAGRA drillholes from northeastern Switzerland. Further, I thank E. Reusser, P. Ulmer and R. Vocke for discussions, H. R. Pfeifer for trace element analyses and U. Krähenbühl for REE analyses. I especially thank Lisa Dietrich and Jill Engi who corrected the English.

References

- AHERN, J.L. and TURCOTTE, D.L. (1979): Magma migration beneath an ocean ridge. *Earth and Planet. Sci. Lett.* 45, 115-122.
- BAILEY, J.C. (1981): Geochemical criteria for a refined discrimination of orogenic andesites. *Chem. Geol.* 32, 139-154.
- BEGER, P.J. (1923): Der Chemismus der Lamprophyre. In: P. Niggli (ed.), *Gesteins- und Mineralprovinzen Vol. 1, Einführung*, 217-577. Borntraeger, Berlin.
- BÜCHI, E., BULETTI, M. und NIGGLI, E. (1984): Neue Aufschlüsse im schweizerischen Anteil des Schwarzwälder Grundgebirges. *Schweiz. mineral. petrogr. Mitt.* 64, 49-66.
- CHAUVEL, C. and JAHN, B.M. (1984): Nd-Sr isotope and REE geochemistry of alkali basalts from the Massif Central, France. *Geochem. Cosmochim. Acta* 48, 93-110.
- DIETRICH, V., OBERHÄNSLI, R. and WALPEN, P. (1976): Röntgenfluoreszenzanalyse der Silikatgesteine. Report Inst. Krist. Petr. ETH-Zürich.
- EIKENBERG, J. (1984): Geologie und Petrographie der Bernina. Teil VIII: Val da Fain. Diploma ETH-Zürich, unpubl.

- GRUBENMANN, U. (1919): Lamprophyrische Gesteine im zentralen Gotthardgranitgneis aus der Umgebung des St.-Gotthard-Hospizes. *Vrtljschr. natf. Ges. Zürich* 64, 418-433.
- HART, S. R. and DAVIS, K. E. (1978): Nickel partitioning between olivine and silica melt. *Earth and Planet. Sci. Lett.* 40, 203-219.
- KÜPFER, TH. (1977): Mineralogisch-petrographische und geochemische Untersuchung der Syenodiorite und der Ganggesteine im Puntegliasgebiet. Ph. D. thesis Univ. Bern, unpubl.
- MANSON, V. (1967): Geochemistry of basaltic rocks: Major elements. In: Basalts, eds. H. H. HESS and A. POLDERVAART, 215-270.
- MÜLLER, H. (1982): Petrographie und Geochemie von Lamprophyren des Schwarzwaldes. Diploma Univ. Freiburg i. Br., unpubl.
- MÜLLER, H. (1984): Die Lamprophyre des Schwarzwaldes. *Fortschr. Miner.* 62, 106-111.
- NICHOLS, I. A. and RINGWOOD, A. E. (1973): Effect of water on olivine stability in tholeiites and production of silica-saturated magmas in the island arc environment. *J. Geol.* 81, 285-300.
- NISBET, E. B., DIETRICH, V. and ESENWEIN, A. (1979): Routine trace element determination in silicate minerals and rocks by X-ray fluorescence. *Fortschr. Mineral.* 57, 264-279.
- OBERRHÄNSLI, R. (1985): Mineralogy and geochemistry of meta-lamprophyres from the Central Swiss Alps. Habilitationsschrift Univ. Bern, unpubl.
- PEARCE, J. A. (1983): Role of the sub-continental lithosphere in magma genesis at active continental margins. In: Continental Basalts and Mantle Xenoliths, eds.: Hawkesworth and Norr; Shiva Geology Series, Shiva, Nantwich, 230-249.
- ROCK, N. M. S. (1984): Nature and origin of calc-alkaline lamprophyres: minettes, vogesites, kersantites and spessartites. *Trans. R. Soc. Edinburgh: Earth Sci.* 74, 193-227.
- SCHALTEGGER, U. (1984): Geologie und Petrographie der Gneiszone von Erstfeld in der Umgebung des Sustenpasses, Aarmassiv. Diploma Univ. Bern, unpubl.
- SEKINE, T. and WYLLIE, P. J. (1982): Phase relationships in the system $KAlSiO_4$ - Mg_2SiO_4 - SiO_2 - H_2O as a model for hybridization between hydrous siliceous melts and peridotite. *Contr. Mineral. Petrol.* 79, 368-374.
- SONDER, R. (1921): Untersuchungen über den Differentiationsverlauf der spät-paläozoischen Granitintrusionen im zentralen und westlichen Gotthardmassiv. *Schweiz. mineral. petrogr. Mitt.* 1, 5-71.
- SPENCER, J. E. (1985): Pliocene low-angle normal faulting and dike emplacement, Homer Mountain and surrounding areas, southeastern California and southernmost Nevada. *Bull. Geol. Soc. Amer.* 96, 1140-1155.
- STEINER, H. (1984): Mineralogisch-petrographische, geochemische und isotopengeologische Untersuchungen an einem Meta-Lamprophyr und seinem grano-dioritischen Nebengestein (Matorello-Gneis) aus der Maggia-Decke. *Schweiz. mineral. petrogr. Mitt.* 64, 261-272.
- SUN, S. S. and NESBITT, R. W. (1977): Chemical heterogeneity of Archean mantle, composition of the earth and mantle evolution. *Earth and Planet. Sci. Lett.* 35, 429-448.
- SUN, S. S. and NESBITT, R. W. (1978): Petrogenesis of Archean ultrabasic and basic volcanics: evidence from rare-earth elements. *Contr. Mineral. Petrol.* 65, 301-325.
- THOMPSON, R. N., MORRISON, M. A., HENDRY, G. L. and PARRY, S. J. (1984): An assessment of the relative roles of crust and mantle in magma genesis: an elemental approach. *Phil. Trans. R. Soc. London A* 310, 549-590.
- VENTURELLI, G., THROPE, R. S., DAL PIAZ, G. V., DEL MORO, A. and POTTS, P. J. (1984): Petrogenesis of calc-alkaline, shoshonitic and associated ultrapotassic Oligocene volcanic rocks from the Northwestern Alps, Italy. *Contrib. Mineral. Petrol.* 86, 209-220.
- WEBER, FR. (1904): Über den Kali-Syenit des Piz Giuf und Umgebung. *Beitr. geol. Karte Schweiz* NF 14.
- WIMMENAUER, W. (1973): Lamprophyre, Semilamprophyre und anchibasaltische Ganggesteine. *Fortschr. Mineral.* 51, 3-67.

- WIMMENAUER, W. and HAHN-WEINHEIMER, P. (1966): Geochemische und petrographische Kriterien für die Herkunft von Kersantiten und Minetten. *N. Jb. Mineral. Mh.* 7, 201–220.
- WOOD, D.A., TARNEY, J., VAREL, J., SAUNDERS, A.D., BOUGAULT, H., JORON, J.L., TREUIL, M. and CANN, J.R. (1979): Geochemistry of Basalts drilled in the North Atlantic by IPOD Leg 49; Implications for mantle heterogeneity. *Earth and Planet. Sci. Lett.* 42, 77–97.
- WYLLIE, P.J. and SEKINE, T. (1982): The formation of mantle phlogopite in subduction zone hybridization. *Contr. Mineral. Petrol.* 79, 375–380.

Manuscript received March 2, 1987; revised manuscript accepted March 25, 1987.